

8.10 The significance of ocean deoxygenation for ocean megafauna

Sheanna Steingass¹ and Yasuhiko Naito²

¹*Oregon State University, Department of Fish and Wildlife, 2030 SE Marine Science Drive, Newport, Oregon, 97365, USA*

²*National Institute of Polar Research, 10-3 Midori-cho, Tachikawa-shi, Tokyo, 190-8518, Japan*

Summary

- Marine mammal distribution is primarily driven by prey availability. Therefore, community-wide impacts on gilled species affect the behaviour of marine mammals. Coastal hypoxia is increasing in areas of critical marine mammal habitat. There are approximately 48 species of marine mammal in the hypoxia-affected regions of the Northern California Current System, Black Sea, Baltic Sea, and Gulf of Mexico.
- Coastal hypoxia events lead to shifts in distribution, mobility, predator avoidance, and mortality of gilled animals. Severe or prolonged hypoxia can lead to shifts in food webs, with the potential for impacting foraging success of marine mammals. Increased ocean warming, and the resultant decreases in oxygen availability suggest an imminent pattern of worsening hypoxia worldwide.
- These patterns may lead to increased pressure for marine mammal species that are already threatened or endangered. Conversely, increased rates of predation on gilled species incapacitated or spatially compressed by hypoxia may benefit certain marine mammals. Direct links between coastal hypoxia and marine mammals may be difficult to quantify, but abundant nearshore marine mammals are excellent potential study species to begin understanding the various implications.
- Ocean deoxygenation drives offshore oxygen minimum zone (OMZ) expansion and shoaling. This expansion may positively affect the foraging efficiency of northern elephant seals, due to (1) their increased ability to feed on inactive prey, (2) decrease of diving costs in terms of both time and energy expenditure, and subsequent increased prey search time in the bottom phase of foraging dives, and (3) elevated prey density related to the compression of vertical prey distribution. Increased dependency on the prey in the OMZ (about 40% of total their prey) may lead to species composition changes within the mesopelagic community. Increased foraging efficiency by elephant seals may lead to further population increases, potentially resulting in large disturbance and shifts in the functioning of the mesopelagic ecosystem through top-down and trophic cascade effects.
- Sperm and beaked whales, typical deep divers in the mesopelagic zone, foraging on squids which inhabit the upper mesopelagic zone and often rest in the OMZ. Although foraging and vertical migration of squids remain unclearly defined, it is suggested that future shoaling of the OMZ will cause the vertical compression of this 'resting zone', consequently giving a foraging advantage to sperm whales and beaked whales feeding on squid.
- Increased foraging efficiency in habitats where the OMZ shoaling occurs will enhance the role of deep diving mammals in nutrient cycling, via altered nutrient pumping to surface waters from the OMZ, leading to increased productivity and nutrient flux to depths. This increased flux could ultimately affect oxygen consumption as a result of microbial respiration and nitrification in the oxygen limited zone, leading to further expansion of the OMZ.



Sheanna Steingass and Yasuhiko Naito

Ocean hypoxia effect	Potential consequences
Decreasing oxygen concentrations in nearshore region produce community-wide impacts on gilled species.	<ul style="list-style-type: none"> • Altered foraging behaviour and spatial habitat use of marine mammals. There are approximately 48 species of marine mammals in the hypoxia-affected regions of the Northern California Current System, the Black Sea, the Baltic Sea, and the Gulf of Mexico. • Shifts in distribution, mobility, predator avoidance, schooling behaviour, and mortality of gilled animals. Severe or prolonged hypoxia can lead to shifting marine community composition, and may potentially impact foraging success of marine mammals. • Increased pressure for marine mammal species, that are already threatened or endangered, to adapt to a changing regime. Conversely, increased predation success on gilled species incapacitated or spatially compressed by hypoxia may benefit certain marine mammals.
Decreasing oxygen concentrations and shoaling in offshore regions impact resting zones of mesopelagic prey.	<ul style="list-style-type: none"> • Increased shoaling may positively affect the foraging efficiency of marine mammals due to (1) their increased ability to successfully feed on inactive or incapacitated prey, (2) decreased diving costs (shallower dives) and increased allotment of search time in the bottom phase of foraging dives, and (3) elevated prey densities related to the vertical and horizontal compression of prey fields. • Increased dependency on the prey in the OMZ (northern elephant seals: about 40%) may lead to species composition and functional changes within the mesopelagic community. • Increased foraging efficiency may lead to population increases (i.e. northern elephant seal), potentially resulting in large functional shifts within the mesopelagic ecosystem. • Future shoaling of the OMZ may provide foraging advantages to sperm whales and beaked whales feeding on squid within these zones. • Increased foraging efficiency will enhance the role of deep-diving mammals in the nutrient cycling of the OMZ via nutrient pumping, leading to increased productivity and nutrient influx to depths. This increased flux could ultimately increase oxygen consumption via microbial respiration and nitrification in the oxygen limited zone, leading to further expansion of the OMZ.

8.10.1 Introduction

Loss of oxygen in the water column may not physiologically affect air-breathing marine mammals, but these animals respond to changes in prey availability as driven by oceanographic factors, including ocean deoxygenation, which makes the threat of increased global deoxygenation to marine mammals as yet unclear. The extent of ocean deoxygenation varies temporally and spatially according to the variability of the environment. Deoxygenation in nearshore areas is very complex due to the influence of land-based inputs and human activities. Severe deoxygenation in the environment has the capacity to affect marine mammals due to changes in prey abundance caused by hypoxic conditions (Chan et al., 2008; Diaz & Rosenberg, 2008; Steingass & Horning, 2017). Offshore deoxygenation occurs with

vertical shoaling of the oxygen minimum zone (OMZ) at a decadal scale as a consequence of global warming (Bograd et al., 2008; Emerson et al., 2004; Ito et al., 2016; Keeling et al., 2010; Stramma et al., 2010, 2012), and this may gradually affect the foraging behaviour of marine mammals.

Although it is difficult to provide a detailed explanation of the relationships between ocean deoxygenation and marine mammals, it is possible to provide (1) the general status of nearshore marine mammals in relation to deoxygenation, and (2) provide a scenario of how pelagic deep diving marine mammals could react to ocean deoxygenation adopting the elephant seal, (*Mirounga angustirostris*) (Figure 8.10.1A) and sperm whales, (*Physeter macrocephalus*) (Figure 8.10.1B) as models of offshore marine mammals.

8.10.2 Nearshore deoxygenation

In recent years, coastal hypoxia has increased in spatial extent, prevalence, and duration on a global scale (Diaz & Rosenberg, 2008). Nearshore coastal ecosystems represent a large percentage of fisheries productivity (Chan et al., 2008). However, ecosystem-scale consequences of coastal hypoxia are just beginning to be examined. Little is known regarding the impacts of deoxygenation events on marine predators, particularly mammals. While not directly reliant on the oxygen in the water column, marine mammals are susceptible to alterations in distribution, availability, behaviour, and mortality of their gilled prey. These effects have been initially examined through conceptual models simulating marine mammal foraging during hypoxia, (Steingass & Horning, 2017), ecosystem monitoring (Craig et al., 2001), and modelling mammal behaviour during related oceanographic studies (Hazen et al., 2013). However, these effects have not been well-studied in the wild.

There are an estimated 48 species of marine mammal (Table 8.10.1) in the hypoxia-affected regions of the Northern California Current System, Black Sea, Baltic Sea, and Gulf of Mexico that may be vulnerable to impacts from ocean deoxygenation.

Table 8.10.1 Marine mammal species (n = 48) inhabiting areas impacted by major hypoxic zones. (Barlow et al., 2010; Becker et al., 2012; Carretta et al., 2017; Ciesielski et al., 2004; Das et al., 2003; Klinowska, 1991; Koschinski, 2001; Natoli et al., 2008; Van de Vijver et al., 2007; Verfuß et al., 2007; Viaud-Martinez et al., 2008; Wells & Scott, 1999; Würsig, 2017) GoM = Gulf of Mexico; CCS = California Current System; BLT = Baltic Sea; BLK = Black Sea

Scientific Name	Common Name	Classification	Region
<i>Balaenoptera acutorostrata</i>	Minke whale	Cetacea	CCS GoM
<i>Balaenoptera borealis</i>	Sei whale	Cetacea	CCS GoM
<i>Balaenoptera edeni</i>	Bryde's whale	Cetacea	CCS GoM
<i>Balaenoptera musculus</i>	Blue whale	Cetacea	CCS GoM
<i>Balaenoptera physalus</i>	Fin whale	Cetacea	CCS GoM
<i>Berardus bairdii</i>	Baird's beaked whale	Cetacea	CCS
<i>Delphinus capensis</i>	Long-beaked common dolphin	Cetacea	CCS GoM
<i>Delphinus delphis</i>	Short-beaked common dolphin	Cetacea	BLK CCS GoM
<i>Eschrichtius robustus</i>	Gray whale	Cetacea	CCS
<i>Eubalaena glacialis</i>	Northern Atlantic right whale	Cetacea	GoM
<i>Feresa attenuata</i>	Pygmy killer whale	Cetacea	GoM
<i>Globicephala macrorhynchus (not confirmed)</i>	Short-finned pilot whale	Cetacea	CCS GoM
<i>Globicephala melas</i>	Long-finned pilot whale	Cetacea	GoM

8.10.2.1 Trends and impacts - nearshore

Marine mammal distribution is primarily driven by prey availability. Therefore, community-wide impacts on gilled species affect the behaviour of marine mammals. Coastal hypoxia is increasing in areas of critical marine mammal habitat (Chan et al., 2008; Conley et al., 2011; Rabalais et al., 2009; Zaitsev, 1992). The severity of these events varies seasonally, but events have recently reached anoxia (<0.5 ml O₂ L⁻¹) or near-complete loss of oxygen, to the point of being unable to sustain gilled animals. In such cases, mass mortality of prey could reduce the ability of mammals to find food.

Coastal hypoxia events lead to shifts in distribution, mobility, predator avoidance, and increased mortality of gilled animals. Severe or prolonged hypoxia can lead to shifts in food web function (Rabalais et al., 2009), with the potential for impacting foraging success of marine mammals. These effects are predicted to increase in the future (Checkley Jr. & Barth, 2009; Gilbert et al., 2010; Grantham et al., 2004; Pierce et al., 2012; Rabalais et al., 2009; Steingass & Horning, 2017).

<i>Grampus griseus</i>	Risso's dolphin	Cetacea	CCS GoM
<i>Kogia breviceps</i>	Pygmy beaked whale	Cetacea	CCS GoM
<i>Kogia sima</i>	Dwarf sperm whale	Cetacea	CCS GoM
<i>Lagenodelphis hosei</i>	Fraser's dolphin	Cetacea	GoM
<i>Lagenorhynchus obliquidens</i>	Pacific white-sided dolphin	Cetacea	CCS
<i>Lissodelphis borealis</i>	Northern right whale dolphin	Cetacea	CCS
<i>Megaptera novaeangliae</i>	Humpback whale	Cetacea	CCS GoM
<i>Mesoplodon</i>	Mesoplodont beaked whales	Cetacea	CCS
<i>Mesoplodon bidens</i>	Sowerby's beaked whale	Cetacea	GoM
<i>Mesoplodon densirostris</i>	Blainville's beaked whale	Cetacea	GoM
<i>Mesoplodon europaeus</i>	Gervai's beaked whale	Cetacea	GoM
<i>Orcinus orca</i>	Killer whale	Cetacea	CCS GoM
<i>Peponocephala electra</i>	Melon-headed whale	Cetacea	GoM
<i>Phocoena phocoena</i> <i>Phocoena phocoena relicta</i> <i>Phocoena phocoena vomerina</i>	Harbour porpoise	Cetacea	BLT BLK CCS
<i>Phocoenoides dalli</i>	Dall's porpoise	Cetacea	CCS
<i>Physeter macrocephalus</i>	Sperm whale	Cetacea	CCS GoM
<i>Pseudorca crassidens</i>	False killer whale	Cetacea	GoM
<i>Stenella attenuata</i>	Pantropical spotted dolphin	Cetacea	GoM
<i>Stenella clymene</i>	Clymene dolphin	Cetacea	GoM
<i>Stenella coeruleoalba</i>	Striped dolphin	Cetacea	CCS GoM
<i>Stenella frontalis</i>	Atlantic spotted dolphin	Cetacea	GoM
<i>Stenella longirostris</i>	Spinner dolphin	Cetacea	GoM
<i>Steno bredanensis</i>	Rough-toothed dolphin	Cetacea	GoM
<i>Tursiops truncatus</i>	Common bottlenose dolphin	Cetacea	BLK CCS GoM
<i>Ziphius cavirostris</i>	Cuvier's beaked whale	Cetacea	CCS GoM
<i>Arctocephalus townsendi</i>	Guadalupe fur seal	Pinnipedia	CCS
<i>Callorhinus ursinus</i>	Northern fur seal	Pinnipedia	CCS
<i>Eumatopias jubatus</i>	Steller sea lion	Pinnipedia	CCS
<i>Halichoerus grypus</i>	Grey seal	Pinnipedia	BLT
<i>Mirounga angustirostris</i>	Northern elephant seal	Pinnipedia	CCS
<i>Phoca hispida</i>	Ringed seal	Pinnipedia	BLT
<i>Phoca vitulina</i>	Harbour seal	Pinnipedia	BLT CCS
<i>Zalophus californianus</i>	California sea lion	Pinnipedia	CCS
<i>Enhydra lutris</i>	Sea otter	Mustelidae	CCS
<i>Trichechus manatus</i>	West Indian manatee	Sirenia	GoM




Figure 8.10.1 (A) Elephant seal (*Mirounga angustirostris*). © MZPHOTO.CZ / Shutterstock.com; (B) Sperm whale (*Physeter macrocephalus*). © wildest animal / Shutterstock.com.

8.10.2.2 Ecosystem consequences – nearshore

A number of fish species consumed by marine mammals, and affected by hypoxia, are also important fisheries species. Domenici et al. (2000, 2002) showed that swim speeds, schooling behaviour, and predator avoidance of Atlantic herring (*Clupea harengus*) (Figure 8.10.2) are all affected by hypoxia. During the most severe oxygen depletion, swim speed decreased until schooling was completely disrupted, leading to decreased predator avoidance capabilities. In lesser sandeel (*Ammodytes tobianus*), moderate hypoxia increased burrowing. As hypoxia continued to increase, the number of fish in the water column doubled, and diel burrowing cycles were disrupted by severe hypoxia (Behrens & Steffensen, 2007; Behrens et al., 2010). All of these behavioural modifications point to increased susceptibility to predation.

Greater understanding of the effects of coastal hypoxia on marine mammals can be obtained through rigorous monitoring of diet, behaviour, spatial distribution and abundance. Direct links between coastal hypoxia and marine mammals remain difficult to quantify, but abundant nearshore marine mammals are likely excellent study species to begin unravelling this issue.

8.10.2.3 Implications of continuing ocean deoxygenation

Increased ocean warming, and the resultant decreases in oxygen retention ability of sea water suggest an imminent pattern of worsening hypoxia worldwide. This pattern may lead to increased pressure on marine mammals that are already threatened or endangered. Conversely, increased rates of predation on gilled

species incapacitated or spatially compressed by hypoxia may benefit certain marine mammals. While initially beneficial for opportunistic or generalist species of mammals, hypoxia-driven ecosystem shifts or faunal collapse may result in irreversible trends limiting marine mammal distribution and populations growth and resilience.

8.10.3 Offshore deoxygenation

8.10.3.1 Deoxygenation and foraging behaviour of offshore marine mammals

The capacity of sea water to retain dissolved oxygen universally decreases with increasing temperature (Keeling et al., 2010). Ocean warming simultaneously reduces the water density and stratifies the upper water column; this leads to decreased oxygen transport to subsurface waters and subsequent reduction in the flux of oxygen into the interior ocean (Keeling et al., 2010). These mechanisms suggest offshore deoxygenation may not directly affect offshore marine mammals foraging in shallow waters, such as baleen whales. However, decadal climatic changes have caused deoxygenation events which present as shoaling of the upper boundary of the OMZ or OMZ expansion, in global marine systems, including the North Pacific, Atlantic, eastern tropical Pacific, and Arabian Sea regions (Andreev & Watanabe, 2002; Bograd et al., 2008; Emerson et al., 2004; Gilly et al., 2013; Ito et al., 2016; Keeling et al., 2010; Stramma et al., 2010, 2012). Although the definition of OMZ varies with the physiological tolerance of organisms and ecosystems to hypoxia (e.g. $<20 \mu\text{mol O}_2 \text{ kg}^{-1}$ or $0.5 \text{ ml O}_2 \text{ L}^{-1}$ in the Pacific region), OMZs have generally expanded globally in the last 50 years (Gilly et al., 2013). Furthermore oxygen levels continue to decline, for example Stramma et al. (2008) describe a



Figure 8.10.2 Large adult male killer whale stalking herring in shallow water off Norway. © Nature Picture Library / Alamy stock photo.

rate of decline ranging from $0.09 - 0.34 \mu\text{mol O}_2 \text{ kg}^{-1} \text{ year}^{-1}$ within the 300 - 700 m depth range in the tropical ocean regions of the eastern Atlantic and the equatorial Pacific.

While there are limited qualitative and quantitative data regarding how marine mammals react to changes in prey conditions in relation to OMZ expansion, there have been consistent changes in mesopelagic foraging behaviour of the northern elephant seal in the eastern North Pacific where the OMZ has expanded (shoaled) (Andreev & Watanabe, 2002), showing evidence that OMZ expansion enhances foraging efficiency of an air-breathing predator. There is also evidence of changes in the diving behaviour of squid-feeding beaked and sperm whales, species that may act as model species to provide simple predictions of how these whales may react to ocean deoxygenation through squid feeding.

8.10.4 Ecological and biogeochemical consequences for northern elephant seals from ocean deoxygenation

Northern elephant seals migrate twice a year, undertaking post-breeding and post-moulting migrations to the north-eastern Pacific (from 38° to 60° N and from the

coast to 172.5° E; Le Boeuf et al., 2000; Robinson et al., 2012), during which they dive to depths of 400 - 600 m (maximum depth: 1735 m; Robinson et al., 2010). Elephant seals dive continuously both day and night, diving to shallower depths at night and deeper depths during the daytime, showing an apparently diel pattern (Figure 8.10.3).

Daytime foraging in the OMZ and OLZ (oxygen limited zone; $20 - 60 \mu\text{mol}$; Gilly et al., 2006) accounts for about 40% of their total feeding, as measured by jaw motion suggesting the importance of daytime foraging despite increased dive depths and thereby reduced foraging efficiency. The high dependency of elephant seals on prey found within the OMZ and OLZ during the day may relate to increased efficiency of capturing prey residing in the OMZ and OLZ at >600 m depth in order to avoid predation pressure in the oxygen rich zone. Gilled animals within the OMZ demonstrate lower metabolic rates making their capture easier (Childress & Seibel, 1998; Naito et al., 2013; Seibel, 2011). Although the mechanistic details are not fully understood, marine hypoxia allows seals to consume large amounts of prey in the OMZ and OLZ due to increased foraging efficiency (roughly estimated total and daytime amounts (i.e. from OMZ): $210,000 \text{ t year}^{-1}$ and $84,000 \text{ t year}^{-1}$




Figure 8.10.3 Typical diel pattern of diving and foraging behaviour based on jaw motion activity of northern elephant seal in the north-eastern Pacific, and typical vertical distribution of oxygen in the region. NFZ and DFZ denote the Night-time Foraging Zone and the Daytime Foraging Zone respectively. These zones of foraging were determined by the depth ranges between average minimum and maximum depth of jaw motion events in every hour during post breeding migration of a seal (Total no. of foraging dives: 6408, Seal ID: PB-1015). Boxes in the vertical oxygen distribution profile indicates depth and oxygen content ranges of NFZ and DFZ.

respectively based on the daily consumption rate: 3% of body mass (Naito et al., 2013), average body mass: 200 kg (assumed value), estimated number of foraging days: 200 (Le Boeuf et al., 2000), approximate population size: 175,000 (Weber et al., 2000) which underlies the important role of seals in structuring the mesopelagic ecosystem. Further shoaling of the OMZ and OLZ in the future may enhance seal foraging efficiency due to (1) increase of inactive prey, (2) decreased diving cost enabling an increase of prey search time during the dive bottom phase, and (3) increased prey density due to vertically-compressed oxygenated habitat (Gilly et al., 2013). These may further increase the dependency of feeding in the expanded OMZ during the day as compared to night-time foraging, which as a result may decrease. A large modification of the day/night dependency ratio (i.e. from the present approximate day to night ratio of 4:6) may trigger an ecological disturbance, ultimately changing the species composition of the deep-sea community as prey composition may differ between day and night (Naito et al., 2017). Enhanced foraging efficiency could elevate the population fitness and breeding rate of elephant seals, leading to a potential long-term increase in population and thereby top-down modification of their terrestrial breeding habitats. However, any future increase of this population in relation to ocean deoxygenation is uncertain, as carrying capacity at breeding sites may also limit

population trends. While long term population increases of elephant seals has been reported (6% increase year⁻¹; Weber et al., 2000), how this increase relates to ocean deoxygenation remains to be confirmed.

The unexpectedly high dependency of elephant seals on prey in the OMZ may have an unprecedented effect on nutrient cycling in relation to OMZ expansion. While an important function of nutrient pumping to the surface water by baleen whales has been reported (Lavery et al., 2010; Roman & McCarthy, 2010), no biological pump information from the OMZ has been reported. Elephant seals, as deep-water foragers, may serve a similar



Figure 8.10.4 Dall's porpoise *Phocoenoides dalli* in south-east Alaska. © Minden Pictures / Alamy stock photo.



Figure 8.10.5 Cuvier's beaked whale (*Ziphius cavirostris*). © Nature Picture Library / Alamy stock photo.

function as nutrient cyclers. Foraging within the OMZ is particularly important in biogeochemical cycling, since prey extraction from the OMZ and subsequent nutrient delivery to the surface in the form of a faecal plume and urine is relatively large allowing seals to play a driving role in this cycle. The pump effect of feeding by elephant seals and other marine mammals enhances the productivity of surface waters (Lavery et al., 2010). Although the quantitative details are not known, it is predicted that increased future productivity will lead to an increase in the export of organic matter to subsurface waters, as a result of deoxygenation in subsurface waters (Gilly et al., 2013) driven by an increase in oxygen consumption by microbial respiration and nitrification, combined with the stratification and surface water warming effects.

8.10.5 Deoxygenation and foraging behaviour of toothed whales

Toothed whales (odontocetes) are dependent upon the mesopelagic zone as their major foraging zone (Belta, 2015); pelagic porpoises and dolphins (small Delphinidae) (Figure 8.10.4) tend to forage in the upper part of the mesopelagic zone, while whales of the Physeteridae, Ziphiidae and large Delphinidae (Figure 8.10.5) dive very deep to forage in the lower part of

the mesopelagic zone (Aoki et al., 2007; Belta, 2015; Clarke, 1996; Miller et al., 2004; Ohizumi et al., 2003; Tyack et al., 2006). Among toothed whales, diving and foraging behaviour of sperm whales and some beaked whales have been investigated extensively, and the results indicate that they hunt and consume squids using an echolocation system in the lower part of the mesopelagic zone, often foraging at deeper depths than elephant seals (Aoki et al., 2007; Blanco & Raga, 2000; Clarke, 1996; Kawakami, 1980; Miller et al., 2004; Ohizumi et al., 2003; Santos et al., 2000; Tyack et al., 2006). This suggests that a key to understanding sperm whale response to ocean deoxygenation is through quantitative understanding of squid behaviour in relation to the OMZ. Although squid forage in the euphotic zone, they are well-adapted to the low oxygen environment and utilize the OMZ to forage and rest (Gilly et al., 2006). These resting squid may give whales a large advantage in foraging. Davis et al. (2006) monitored diving behaviour of sperm whales and the jumbo squid, *Dosidicus gigas*, which rests in the upper boundary of the OMZ (>300 m depth) after feeding on myctophid fish in warmer shallow waters in the Gulf of California. Consistent dive depths of sperm whales foraging for jumbo squid suggested that the OMZ expansion compresses the distributional range of squids allowing whales to forage upon them

more efficiently. Compared to sperm whales within the shallow OMZ in the Gulf of California, other populations of sperm whales often dive to deeper depths (400-1000 m) in zones with no apparent OMZ, including the North Atlantic, Gulf of Mexico, Ligurian Sea, Kumano coast of Japan, and Ogasawara Islands (Aoki et al., 2017; Watwood, 2006). These differences in foraging depth suggest that sperm whales and beaked whales may in fact experience a large advantage to foraging within areas of OMZ shoaling, due to greater efficiency in feeding upon squid. As with elephant seals, increased efficiency of foraging by toothed whales could lead to significant disturbance and disruption to the marine ecosystem through top-down effects leading to a trophic cascade (Estes & Palmisano, 1974) of ecosystem function in the future.

Toothed whales may have a similar deep biological pump function which affects biogeochemical cycling as in the case of the northern elephant seal, but the details of this pump function are as yet not clearly understood. Further investigations detailing the relationship between marine mammals and mesopelagic deoxygenation are necessary to enable a clear understanding of how ocean warming is affecting the deep sea ecosystem.

8.10.6 References

- Andreeve, A., & Watanabe, S. (2002). Temporal changes in dissolved oxygen of the intermediate water in the subarctic North Pacific. *Geological Research Letters*, 29, 14, 1680. <https://doi.org/10.1029/2002GL015021>
- Aoki, K., Amano, M., Yoshioka, M., Mori, K., Tokuda, D., & Miyazaki, N. (2007). Diel diving behavior of sperm whales off Japan. *Marine Ecology Progress Series*, 349, 277-287. <https://doi.org/10.3354/meps07068>
- Barlow, J., Henry, A., Redfern, J.V., Yack, T., Jackson, A., Hall, C., ... Ballance, L. (2010). Oregon, California and Washington line-transect and ecosystem (ORCAWALE) 2008 Cruise Report.
- Becker, E.A., Forney, K.A., Ferguson, M.C., Barlow, J., & Redfern, J. (2012). Predictive Modeling of Cetacean Densities in the California Current Ecosystem Based on Summer/fall Ship Surveys in 1991-2008. US Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, Southwest Fisheries Science Center - 499.
- Belta, A. (Ed.) (2015). *Whales, Dolphins, and Porpoises: a natural history and species guide*. The University of Chicago Press, Chicago, 288pp. <https://doi.org/10.7208/chicago/9780226183220.001.0001>
- Behrens, J.W., & Steffensen, J.F. (2007). The effect of hypoxia on behavioural and physiological aspects of lesser sandeel, *Ammodytes tobianus* (Linnaeus, 1785). *Marine Biology*, 150, 1365-1377. <https://doi.org/10.1007/s00227-006-0456-4>
- Behrens, J.W., Petersen, J.K., Ærtebjerg, G., & Steffensen, J. (2010). Influence of moderate and severe hypoxia on the diurnal activity pattern of lesser sandeel *Ammodytes tobianus*. *Journal of Fish Biology*, 77, 538-551. <https://doi.org/10.1111/j.1095-8649.2010.02697.x>
- Blanco, C., & Raga, J.A. (2000). Cephalopod prey of two *Ziphius cavirostris* (Cetacea) stranded on the western Mediterranean coast. *Journal of the Marine Biological Association of the United Kingdom*, 80, 381-382. <https://doi.org/10.1017/S0025315499002064>
- Bograd, S.J., Castro, C., Di Lorenzo, E., Palacios, D.M., Bailey, H., Gilly, W., & Chavez, F.P. (2008). Oxygen declines and the shoaling of the hypoxic boundary in the California Current. *Geophysical Research Letters*, 35, L1 2607. <https://doi.org/10.1029/2008GL034185>
- Carretta, J.V., Forney, K.A., Lowry, M.S., Barlow, J., Baker, J., Johnston, D., ... Mattila, D.K. (2017). US Pacific marine mammal stock assessments: 2016. NOAA NMFS/SWFSC.
- Chan, F., Barth, J., Lubchenco, J., Kirincich, A., Weeks, H., Peterson, W., & Menge, B. (2008). Emergence of anoxia in the California current large marine ecosystem. *Science*, 319, 920-920. <https://doi.org/10.1126/science.1149016>
- Checkley, Jr. D.M., & Barth, J.A. (2009). Patterns and processes in the California Current System. *Progress in Oceanography*, 83, 49-64. <https://doi.org/10.1016/j.pocean.2009.07.028>
- Ciesielski, T., Wasik, A., Kuklik, I., Skóra, K., Namieśnik, J., & Szefer, P. (2004). Organotin compounds in the liver tissue of marine mammals from the Polish coast of the Baltic Sea. *Environmental Science & Technology*, 38, 1415-1420. <https://doi.org/10.1021/es034790z>
- Childress, J.J., & Seibel, B.A. (1998). Life at stable low oxygen levels: Adaptations of animals to oceanic oxygen minimum layers. *Journal of Experimental Biology*, 201, 1223-1232.
- Clarke, M.R. (1996). The role of cephalopods in the world's oceans: an introduction. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 351, 979-983. <https://doi.org/10.1098/rstb.1996.0088>
- Conley, D., Carstensen, J., Aigars, J., Axe, P., Bonsdorff, E., Eremina, T., ... Zillen, L. (2011). Hypoxia Is Increasing in the Coastal Zone of the Baltic Sea. *Environmental Science & Technology*, 45, 6777-6783. <https://doi.org/10.1021/es201212r>
- Craig, J.K., Crowder, L.B., Gray, C.D., McDaniel, C.J., Henwood, T. A., & Hanifen, J.G. (2001). Ecological effects of hypoxia on fish, sea turtles, and marine mammals in the northwestern Gulf of Mexico. In N.N. Rabalais & R.E. Turner (Eds.). *Coastal hypoxia: consequences for living resources and ecosystems*. American Geophysical Union, Washington, DC, pp. 269-291. <https://doi.org/10.1029/CE058p0269>
- Crocker, D.E., Costa, D.P., Le Boeuf, B.J., Webb, P.M., & Houser, D.S. (2006). Impact of El Niño on the foraging behavior of female northern elephant seals. *Marine Ecology Progress Series*, 309, 1-10. <https://doi.org/10.3354/meps309001>
- Das, K., Beans, C., Holsbeek, L., Mauger, G., Berrow, S.D., Rogan, E., & Bouquegneau, J.M. (2003). Marine mammals from northeast atlantic: relationship between their trophic status as determined by [delta] 13C and [delta] 15N measurements and their trace

- metal concentrations. *Marine Environmental Research*, 56, 349-365. [https://doi.org/10.1016/S0141-1136\(02\)00308-2](https://doi.org/10.1016/S0141-1136(02)00308-2)
- Davis, R.W., Jaquet, N., Gendron, D., Markaida, U., Bazzino, G., & Gilly, W.F. (2007). Diving behavior of sperm whales in relation to behavior of a major prey species, the jumbo squid, in the Gulf of California, Mexico. *Marine Ecology Progress Series*, 333, 291-302. <https://doi.org/10.3354/meps333291>
- Diaz, R.J., & Rosenberg, R. (2008). Spreading dead zones and consequences for marine ecosystems. *Science*, 321, 926-929. <https://doi.org/10.1126/science.1156401>
- Domenici, P., Steffensen, J.F., & Batty, R.S. (2000). The effect of progressive hypoxia on swimming activity and schooling in Atlantic herring. *Journal of Fish Biology*, 57, 1526-1538. <https://doi.org/10.1111/j.1095-8649.2000.tb02229.x>
- Domenici, P., Ferrari, R.S., Steffensen, J.F., & Batty, R.S. (2002). The effect of progressive hypoxia on school structure and dynamics in Atlantic herring *Clupea harengus*. *Proceedings of the Royal Society of London B, Biological Sciences*, 269, 2103-2111. <https://doi.org/10.1098/rspb.2002.2107>
- Emerson, S., Watanabe, Y.W., Ono, T., & Mecking, S. (2004). Temporal trends in apparent oxygen utilization in the upper pycnocline of the North Pacific: 1980-2000. *Journal of Oceanography*, 60, 139-147. <https://doi.org/10.1023/B:JOCE.0000038323.62130.a0>
- Estes, J.A., & Palmisano, J.F. (1974). Sea otters: their role in structuring nearshore communities. *Science*, 185, 1058-1060. <https://doi.org/10.1126/science.185.4156.1058>
- Gilbert, D., Rabalais, N., Diaz, R., & Zhang, J. (2010). Evidence for greater oxygen decline rates in the coastal ocean than in the open ocean. *Biogeosciences*, 7, 2283-2296. <https://doi.org/10.5194/bg-7-2283-2010>
- Gilly, W.F., Markaida, U., Baxter, C.H., Block, B.A., Boustany, A., Zeidberg, L., ... Salinas, C. (2006). Vertical and horizontal migrations by jumbo squid, *Dosidicus gigas*, revealed by electronic tagging. *Marine Ecology Progress Series*, 324, 1-17. <https://doi.org/10.3354/meps324001>
- Gilly, W.F., Beman, J.M., Litvin, S.Y., & Robison, B.H. (2013). Oceanographic and biological effects of shoaling of the oxygen minimum zone. *Annual Review of Marine Science*, 5, 393-420. <https://doi.org/10.1146/annurev-marine-120710-100849>
- Grantham, B.A., Chan, F., Nielsen, K.J., Fox, D.S., Barth, J.A., Huyer, A., ... Menge, B.A. (2004). Upwelling-driven nearshore hypoxia signals ecosystem and oceanographic changes in the northeast Pacific. *Nature*, 429, 749-754. <https://doi.org/10.1038/nature02605>
- Hazen, E.L., Jorgensen, S., Rykaczewski, R.R., Bograd, S.J., Foley, D.G., Jonsen, I.D., ... Crowder, L.B. (2013). Predicted habitat shifts of Pacific top predators in a changing climate. *Nature Climate Change*, 3, 234. <https://doi.org/10.1038/nclimate1686>
- Ito, T., Nenes, A., Jonson, M.S., Meskhidze, N., & Deutsch, C. (2016). Acceleration of oxygen decline in the tropical Pacific over the past decades by aerosol pollutants. *Nature Geoscience*, 9, 443-447. <https://doi.org/10.1038/ngeo2717>
- Kawakami, T. (1980). A review of sperm whale food. *The Scientific Reports of the Whales Research Institute*, 32, 199-218.
- Keeling, R.F., Körtzinger, A., & Gruber, N. (2010). Ocean deoxygenation in a warming world. *Annual Review of Marine Science*, 2, 199-229. <https://doi.org/10.1146/annurev.marine.010908.163855>
- Klinowska, M. (1991). Dolphins, porpoises and whales of the world: the IUCN Red Data Book. IUCN, Gland, Switzerland.
- Koschinski, S. (2001). Current knowledge on harbour porpoises (*Phocoena phocoena*) in the Baltic Sea. *Ophelia*, 55, 167-197. <https://doi.org/10.1080/00785326.2001.10409483>
- Lavery, T.J., Roundnew, B., Gill, P., Seymour, J., Seuront, L., Johnson, G., ... Smetacek, V. (2010). Iron deflection by sperm whales stimulates carbon export in the Southern Ocean. *Proceedings of the Royal Society of London B, Biological Sciences*, 277, 3527-3531. <https://doi.org/10.1098/rspb.2010.0863>
- Le Boeuf, B.J., Crocker, D.E., Costa, D.P., Blackwell, S.B., Webb, P.M., & Houser, D.S. (2000). Foraging ecology of northern elephant seals. *Ecological Monographs*, 70, 353-382. [https://doi.org/10.1890/0012-9615\(2000\)070\[0353:FEONES\]2.0.CO;2](https://doi.org/10.1890/0012-9615(2000)070[0353:FEONES]2.0.CO;2)
- Miller, P.J.O., Johnson, M.P., Tyack, P.L., & Terray, E.A. (2004). Swimming gaits, passive drag and buoyancy of diving sperm whales *Physeter macrocephalus*. *Journal Experimental Biology*, 207, 1953-1967. <https://doi.org/10.1242/jeb.00993>
- Naito, Y., Costa, D.P., Adachi, T., Robinson, P.W., Fowler, M., & Takahashi, A. (2013). Unravelling the mysteries of a mesopelagic diet: a large apex predator specializes on small prey. *Functional Ecology*, 27, 710-717. <https://doi.org/10.1111/1365-2435.12083>
- Naito, Y., Costa, D.P., Adachi, T., Robinson, P.W., Peterson, S.H., Mitani, Y., & Takahashi, A. (2017). Oxygen minimum zone: An important oceanographic habitat for deep diving northern elephant seals, *Migrunga angustirostris*. *Ecology & Evolution*, 7, 6259-6270. <https://doi.org/10.1002/ece3.3202>
- Natoli, A., Cañadas, A., Vaquero, C., Politi, E., Fernandez-Navarro, P., & Hoelzel, A.R. (2008). Conservation genetics of the short-beaked common dolphin (*Delphinus delphis*) in the Mediterranean Sea and in the eastern North Atlantic Ocean. *Conservation Genetics*, 9, 1479-1487. <https://doi.org/10.1007/s10592-007-9481-1>
- Ohizumi, H., Isoda, T., Kishoro, T., & Kato, H. (2003). Feeding habits of Baird's beaked whale *Berardius bairdii*, in the western North Pacific and Sea of Okhotsk off Japan. *Fisheries Science*, 69, 11-20. <https://doi.org/10.1046/j.1444-2906.2003.00582.x>
- Paulmier, A., & Ruiz-Pino, D. (2008). Oxygen minimum zones (OMZs) in the modern ocean. *Progress in Oceanography*, 80, 113-128. <https://doi.org/10.1016/j.pocean.2008.08.001>
- Pierce, S.D., Barth, J.A., Shearman, R.K., & Erofeev, A.Y. (2012). Declining Oxygen in the Northeast Pacific. *Journal of Physical Oceanography*, 42, 495-501. <https://doi.org/10.1175/JPO-D-11-0170.1>
- Rabalais, N.N., Turner, R.E., & Wiseman, W.J. (2001). Hypoxia in the Gulf of Mexico. *Journal of Environmental Quality*, 30, 320-329. <https://doi.org/10.2134/jeq2001.302320x>
- Rabalais, N.N., Turner, R.E., Diaz, R.J., & Justic, D. (2009). Global change and eutrophication of coastal waters. *ICES Journal of Marine Science*, 66, 1528-1537. <https://doi.org/10.1093/icesjms/fsp047>

- Robinson, P.W., Simmons, S.E., Crocker, D.E., & Costa, D.P. (2010). Measurements of foraging success in a highly pelagic marine predator, the northern elephant seal. *Journal of Animal Ecology*, 79, 1146–1156. <https://doi.org/10.1111/j.1365-2656.2010.01735.x>
- Robinson, P.W., Costa, D.P., Crocker, D.E., Gallo-Reynoso, J.P., Champagne, C.D., Fowler, M.A., ... Yoda, K. (2012). Foraging behavior and success of a mesopelagic predator in the Northeast Pacific Ocean: insights from a data-rich species, the northern elephant seal. *PLoS ONE*, 7, e36728. <https://doi.org/10.1371/journal.pone.0036728>
- Roman, J., & McCarthy, J.J. (2010). The whale pump: Marine mammals enhance primary productivity in a coastal basin. *PLoS ONE*, 5, e13255. <https://doi.org/10.1371/journal.pone.0013255>
- Santos, M.B., Clarke, M.R., & Pierce, G.J. (2001). Assessing the importance of cephalopods in the diets of marine mammals and other top predators: problems and solutions. *Fisheries Research*, 52, 121–139. [https://doi.org/10.1016/S0165-7836\(01\)00236-3](https://doi.org/10.1016/S0165-7836(01)00236-3)
- Seibel, B.A. (2011). Critical oxygen levels and metabolic suppression in oceanic oxygen minimum zones. *Journal Experimental Biology*, 214, 326–336. <https://doi.org/10.1242/jeb.049171>
- Steingass, S. (2017). Dietary Composition of Four Stocks of Pacific Harbor Seal (*Phoca vitulina richardii*) in the Northern California Current Large Marine Ecosystem as Synthesized from Historical Data, 1931–2013. *Northwestern Naturalist*, 98, 8–23. <https://doi.org/10.1898/NWN16-05.1>
- Steingass, S., & Horning, M. (2017). Individual-based energetic model suggests bottom up mechanisms for the impact of coastal hypoxia on Pacific harbor seal (*Phoca vitulina richardii*) foraging behavior. *Journal of Theoretical Biology*, 416, 190–198. <https://doi.org/10.1016/j.jtbi.2017.01.006>
- Stramma, L., Johnson, G.C., Sprintall, J., & Mohrholz, V. (2008). Expanding oxygen-minimum zones in the tropical oceans. *Science*, 320, 655–658. <https://doi.org/10.1126/science.1153847>
- Stramma, L., Johnson, G.C., Firing, E., & Schmidtko, S. (2010). Eastern Pacific oxygen minimum zones: Supply paths and multidecadal changes. *Journal of Geophysical Research*, 115, C09011. <https://doi.org/10.1029/2009JC005976>
- Stramma, L., Oschlies, A., & Schmidtko, S. (2012). Mismatch between observed and modeled trends in dissolved upper-ocean oxygen over the last 50 yr. *Biogeosciences*, 9, 4045–4057. <https://doi.org/10.5194/bg-9-4045-2012>
- Tyack, P.L., Johnson, M., Aguilar de Soto, N.A., Sturlese, A., & Madsen, P. T. (2006). Extreme diving of beaked whales. *Journal Experimental Biology*, 209, 4238–4253. <https://doi.org/10.1242/jeb.02505>
- Van de Vijver, K.I., Holsbeek, L., Das, K., Blust, R., Joiris, C., & De Coen, W. (2007). Occurrence of perfluorooctane sulfonate and other perfluorinated alkylated substances in harbor porpoises from the Black Sea. *Environmental Science & Technology*, 41, 315–320. <https://doi.org/10.1021/es060827e>
- Vaquer-Sunyer, R., & Duarte, C.M. (2008). Thresholds of hypoxia for marine biodiversity. *Proceedings of the National Academy of Sciences of the United States of America*, 105, 15452–15457. <https://doi.org/10.1073/pnas.0803833105>
- Vaquer-Sunyer, R., & Duarte, C.M. (2011). Temperature effects on oxygen thresholds for hypoxia in marine benthic organisms. *Global Change Biology*, 17, 1788–1797. <https://doi.org/10.1111/j.1365-2486.2010.02343.x>
- Verfuß, U.K., Honnef, C.G., Meding, A., Dähne, M., Mundry, R., & Benke, H. (2007). Geographical and seasonal variation of harbour porpoise (*Phocoena phocoena*) presence in the German Baltic Sea revealed by passive acoustic monitoring. *Journal of the Marine Biological Association of the United Kingdom*, 87, 165–176. <https://doi.org/10.1017/S0025315407054938>
- Viaud-Martinez, K.A., Brownell, R.L., Komnenou, A., & Bohonak, A.J. (2008). Genetic isolation and morphological divergence of Black Sea bottlenose dolphins. *Biological Conservation*, 141, 1600–1611. <https://doi.org/10.1016/j.biocon.2008.04.004>
- Watwood, S.L., Miller, P.J.O., Johnson, M., Madsen, P., & Tyack, P.L. (2006). Deep-diving foraging behavior of sperm whales (*Physeter macrocephalus*). *Journal Animal Ecology*, 75, 814–825. <https://doi.org/10.1111/j.1365-2656.2006.01101.x>
- Wells, R.S., & Scott, M.D. (1999). Bottlenose dolphin *Tursiops truncatus* (Montagu, 1821). *Handbook of marine mammals: the second book of dolphins and porpoises*, 6, 137–182.
- Weber, D.S., Stewart, B.S., Garza, J.C., & Lehman, N. (2000). An empirical genetic assessment of the severity of the northern elephant seals population bottleneck. *Current Biology*, 10, 1287–1290. [https://doi.org/10.1016/S0960-9822\(00\)00759-4](https://doi.org/10.1016/S0960-9822(00)00759-4)
- Würsig, B. (2017). Marine mammals of the Gulf of Mexico. *Habitats and Biota of the Gulf of Mexico: Before the Deepwater Horizon Oil Spill*. Springer, pp. 1489–1587. https://doi.org/10.1007/978-1-4939-3456-0_5
- Zaitsev, Y.P. (1992). Recent changes in the trophic structure of the Black Sea. *Fisheries Oceanography*, 1, 180–189. <https://doi.org/10.1111/j.1365-2419.1992.tb00036.x>

