Chief
Scientist's
Office

Agriculture in the Lower Mekong Basin: Experience from the Critical Decade of 1966–1976

Willem van Liere and Jeffrey A. McNeely

**Agriculture in the
Lower Mekong Basin:
Experience from the
Critical Decade of 1966–1976**

The designation of geographical entities in this book, and the presentation of the material, do not imply the expression of any opinion whatsoever on the part of IUCN concerning the legal status of any country, territory, or area, or of its authorities, or concerning the delimitation of its frontiers or boundaries.

The views expressed in this publication do not necessarily reflect those of IUCN.

Published by: IUCN, Gland, Switzerland and Cambridge, UK

Copyright: © 2005 International Union for Conservation of Nature and Natural Resources

Reproduction of this publication for educational or other non-commercial purposes is authorized without prior written permission from the copyright holder provided the source is fully acknowledged.

Reproduction of this publication for resale or other commercial purposes is prohibited without prior written permission of the copyright holder.

Citation: van Liere, Willem and McNeely, Jeffrey. A. 2005. *Agriculture in the Lower Mekong Basin: Experience from the Critical Decade of 1966–1976*. IUCN, Gland, Switzerland and Cambridge, UK. 190 pp.

ISBN: 2-8317-0891-5

Cover design by: Creative Service Section, Allied Printers, Thailand

Photographs courtesy: Bangkok Post

Photograph page 4 courtesy: NASA

Cover photo: Angkor Wat, symbol of a great Mekong civilization

Layout by: Creative Service Section, Allied Printers, Thailand

Produced by: Allied Printers, Thailand

Printed by: Allied Printers, a division of The Post Publishing Public Company Limited, Thailand

Available from: IUCN Publications Services Unit
219c Huntingdon Road, Cambridge CB3 0DL
United Kingdom
Tel: +44 1223 277894
Fax: +44 1223 277175
E-mail: info@books.iucn.org
www.iucn.org/bookstore

A catalogue of IUCN publications is also available.

The text of this book is printed on Woodfree, 80 gsm.

Agriculture in the Lower Mekong Basin: Experience from the Critical Decade of 1966–1976

Willem van Liere and Jeffrey A. McNeely

Table of Contents

7 **Acknowledgements**

9 **Preface**

PART I

Experience from the Critical Decade of 1966–1976

17 **Chapter 1**
Geology and Physiographic Units

23 **Chapter 2**
Climate and Water Resources

33 **Chapter 3**
Basin Soils and Their Major Limitations

43 **Chapter 4**
The Natural Vegetation

53 **Chapter 5**
The Native Fauna

67 **Chapter 6**
Resource Conservation and Water Resources Development

PART II

The Agricultural Development Programme

77 **Chapter 7**
Agricultural Systems

97 **Chapter 8**
The Agricultural Development Model

119 **Chapter 9**
Pioneer Agriculture Projects

165 **Chapter 10**
Summary and Conclusions

185 **Chapter 11**
The Lower Mekong Basin Today

187 **References**

Acknowledgements

This book was written thirty years ago to serve as a scientific basis for the Mekong Committee (Committee for the Coordination of Studies in the Lower Mekong Basin) in establishing its work programme in the field of agricultural development and conservation. The basic data for the book were collected by the staff of the Agriculture Division of the Mekong Secretariat and we wish to thank in particular, among its staff, Takeshi Kawai, irrigation engineer, Francois Dauphin, agronomist and Jacques Ceruse, geologist, for their important contributions.

Special thanks go to Sue Rallo, who had the very difficult task of converting scanned text into accurate modern electronic text, a good illustration of the technological changes that have taken place since the book was originally written. Thanks also to Frederik Schutyser, who did the actual scanning work and helped in many other ways. Wendy Price has been responsible for incorporating all of the editorial corrections and preparing the final text for publication, for which our deep thanks. Special thanks also go to Pojanan Suyaphan who provided invaluable moral support during, and after, the actual writing of the manuscript.

Preface

The decade from 1966 to 1976 was a tumultuous time in the history of the Lower Mekong Basin. Three of the four countries were at war (Viet-Nam, Lao PDR, and Cambodia) and the fourth (Thailand) sent troops and provided bases for the conflict. Remarkably, this was also a time of major planning for developing the agricultural production systems of the Lower Mekong Basin. This book describes the pioneering work done in that decade under the auspices of the Committee for the Coordination of Investigations in the Lower Mekong Basin, otherwise known as the Mekong Committee.

We describe this work based on a draft publication that we prepared in 1977, but we have made some editorial changes and have improved the quality of some of the figures. Many of the maps that were prepared by the Mekong Committee remain the best available on the respective topics. Because of the critical importance of agriculture to the people of South East Asia, we are convinced that publishing this information will be a welcome contribution to the literature and provide an essential historical dimension to current agricultural development activities. We have also added a concluding chapter that briefly summarises what has happened in the thirty years since we did our research.

While the agricultural development programme sponsored by the Mekong Committee was concerned with both short-term and long-term needs, its primary objective was the long-term sustainable development of the basin. Such long term planning was seen to be critically needed for three main reasons: population increase; overexploitation of natural resources; and the inevitability of large-scale application of modern technology in agricultural production. The latter requires a systematic approach, because transforming existing agricultural practices into sustained-yielding farm systems able to meet future needs, taking into account ecological and other constraints, cannot be achieved on a piece-meal basis.

An important aspect of the long-term programme related to agricultural experimentation and pioneer agriculture projects. The specific objective of agricultural experimentation is to develop new methods for enhancing agricultural production and to test them under limited-scale controlled conditions. The pioneer agriculture projects were intended to apply these new methods under actual full-scale conditions in areas from 5,000 to 10,000 hectares, including development of farm management practices such as land preparation, water distribution, use of agricultural chemicals, crop storage, marketing, and financing. Thus the pioneer projects served to test and demonstrate the feasibility of new concepts in agriculture, beginning with smaller areas and progressing to ever larger areas as experience was gained, until eventually the new concepts could be applied broadly in the region.

The Mekong Committee's agricultural programme also assisted in planning and implementing short-term agricultural projects in the riparian countries. The objectives of these short-term activities were different for each country, but each programme aimed at immediate results, over as large a geographical area as possible and with an investment as economically sensible as possible.

In northeast Thailand the short-term programme objective was to ensure continued economic growth. To this end, the Government launched various general improvement programmes, including with irrigation, upland crops improvement, and livestock.

In Cambodia, the Lao PDR and Viet-Nam, countries that experienced a devastating war during most of the time described (the wars ended in 1975), the first priority was to regain self-sufficiency in food in the shortest period of time. To this end, agricultural production programmes were carried out with large-scale public participation which ensured a speed of execution never before achieved. It seemed likely that in all three countries the physical facilities to attain self-sufficiency would be in place in the fairly near future.

However, short-term programmes are unlikely to remove the constraints that impede sustained agricultural production, both technically and organizationally. For example, it was imperative to develop farm systems with sustained yields for the poorer uplands. It was equally important to develop intensive and diversified farm systems for the lowlands and to apply these at acceptable risk levels. Achieving optimal overall agricultural production in both uplands and lowlands required a broader approach and the entire watershed had to be considered, with watershed management and protection playing an increasingly important role. Short-term programmes are not well adapted to solving such problems.

Even more disconcerting than the agro-technical and agro-environmental problems noted above was recognition that very soon the shortage of water in the lower basin during the latter part of the dry season, when water requirements are at their peak, was bound to become the major limiting factor in agricultural development. As of 1975, in northeast Thailand and Lao PDR some 15 thousand million m³ of water were being stored in existing tributary reservoirs; much of this water was still unused in the dry season and served to increase the low flow of the Mekong river. However, the short-term programmes of Thailand, the Lao PDR and Cambodia all aimed at increased use of water during the dry season. Within a time-span perhaps as short as 10-15 years, it was expected that no extra water would be available for agriculture in the Vietnamese delta during the critical months of low flow (March-May).

More dams on the tributaries may not be able to contribute substantially to further agricultural development of the basin. In north-east Thailand practically all available dam-sites have been used already; in Cambodia the flows of the tributaries to the Great Lake are small; and additional dams on tributaries in the Lao PDR would be primarily single purpose power projects. It is sometimes said that local storage, using small reservoirs in the flat lowlands of northeast Thailand and in the Vietnamese delta, could compensate for the shortage. However, experience has shown that local storage is scarcely effective precisely when it is most needed, at the end of the dry season; moreover it is costly and it occupies much valuable productive land.

The conclusion is inescapable that the future of agriculture in the basin can be ensured only with significant new investments across a broad front, including new cropping systems, water management and infrastructure.

In summary, sustainable development would enable an effective integration of both conservation and development of the natural resources of the lower basin, so that optimal overall sustained production would be achieved. This was seen as a realistic prospect and assuming it could be achieved, the lower Mekong basin would satisfy not only its own food needs, but the basin could produce significant annual surpluses of food, feed and fiber in the early part of the next century. The agricultural programme of the Mekong Committee was designed to pursue these prospects, and this book described the approaches that were taken. The alert reader will note that many of these approaches, seen as innovations in 1967-1977, are today in the mainstream of sustainable agriculture.

Willem van Liere
Bangkok, Thailand
willem@csloxinfo.com

Jeffrey A. McNeely
Gland, Switzerland
JAM@iucn.org

10 September 2005

PART I
Experience from
the Critical Decade of
1966 – 1976

Chapter 1

Geology and Physiographic Units

Introduction

This introductory chapter presents a broad painting of the evolution of the physical land resources of the Lower Mekong Basin and relates the existing land formations to agricultural practices and potentials. First the geological history is described, followed by a classification of the land forms in the lower basin into five physiographic units and an assessment of the relative potentials for agriculture for each of the units. In subsequent chapters of Part I, further details of climate, water resources, soils, vegetation types, and wildlife will be added to present a comprehensive picture of the resource background for agricultural development in the basin.

Geological History

The Lower Mekong Basin rests on an ancient block of continental crust, called Indosinia by geologists, that has remained relatively stable since the Jurassic Period, a period of over 150 million years. Large areas were covered by an inland sea during the upper Mesozoic Era, during which thick red-bed sandstones and evaporites were deposited.

Strong uplift and igneous activity in northern Vietnam during the late Cretaceous and early Tertiary periods, raised the Fan Si Pan range and developed the Red River rift along its northeast flank (this rift has been filled with nearly 1,000 meters of Tertiary and Quaternary sediments). During this period, Indosinia was warped into broad domes and basins along primarily southeasterly trending axes and secondary axes at right angles. Northeast Thailand and adjacent parts of Lao PDR (the Korat Plateau) remained mostly low-lying, so that the Mesozoic succession is preserved virtually complete in the Mun-Chi and Sakhon Nakhon basins, with deposits of salts which are 250 meters thick in places (Takaya, 1974). Over much of Cambodia, on the other hand, the Mesozoic sandstones were progressively domed up during the Tertiary and as a result were largely leveled by erosion, revealing granite outcrops from intrusions of the basement complex in many areas. In northern Thailand, block faulting formed flat, agriculturally important plains between the steep mountain ranges.

Although the basement complex of Indosinia has remained relatively stable over geological time, the shaping of the present-day Southeast Asian sub-continent has been achieved by various tectonic movements involving the Indian and Chinese continental shields, movements which are still continuing (Molnar and Tapponier, 1975). For example, tectonic uplift has caused the Himalayas to rise 3,000 meters since the middle Pleistocene (Hagen, 1960),

and these orogenic movements have extended from the Himalayas through Burma, Sumatra, central Indonesia, the Philippines, Taiwan, and Japan. This has enclosed the Indosinia region on all sides and has caused the tilting of the basin which resulted in the drowning of the eastern and southern parts of Indosinia to form the South China Sea and the Gulf of Thailand.

The Quaternary period (Pleistocene plus Holocene epochs, basically the past two million years) has been most important in forming the present land forms, river courses, and major soil groups. This was a time of tectonic and volcanic events, major climatic shifts, and changes in sea level. About 600,000 years ago, a stage of vulcanism formed most of the basalt areas in the Annamites, northeast Thailand, and Cambodia. About the same time, or somewhat earlier, a period of deformation further uplifted the Phu Phan range.

The Pleistocene epoch was characterized by climatic fluctuations with humid stages between dry periods. The dry periods probably coincided with glaciations at higher latitudes and they were responsible for major erosion in the Lower Mekong Basin (Carbonnel, 1972). During these periods, much of the forest cover may have disappeared on the Korat plateau and in the Mekong plain, resulting in large erosional alluvial fans with sands, gravels and boulders. On the Korat plateau the older alluvial surfaces have progressively higher elevations and greater slopes; by contrast, in the Mekong plain, each progressively older plain tends to pass under the deposits forming the next younger plain. Elevation of the older parts and the subsiding of the younger parts probably continued to occur as recently as the late Pleistocene.

Very little is known in detail about the Quaternary of the Mekong river upstream of the delta. In contrast with other rivers originating from the Himalayas, such as the Irrawaddy, Salween, and Yang Tse, no Pleistocene deposits of any importance are known along the Mekong's course. Some of the Quaternary deposits found well inland are probably of marine origin as is implied by the fossil coastline with large funnel shaped tidal valleys around the Great Lake, which would correspond with a higher Pleistocene sea level. However, as late as 20,000 years ago, the sea level was again about 100 meters lower than at present.

The Holocene epoch, starting about 10,000 years ago, began rapidly rising sea level. In Vietnam, lagoonal deposits were formed behind a series of sandy shorelines and these account for the acid surface soils of the Plain of Reeds and the Trans-Bassac. Some 1 to 2 meters of recent fresh-water alluvium have been deposited in the Great Lake, the base of which has been dated at about 5,700 years of age (Carbonnel, 1972).

Physiographic Units

As the result of the complex geological history sketched above, the Lower Mekong Basin now consists of five rather distinct physiographic units: the Northern Highlands; the Annamite Chain; the Southern Uplands; the Korat Plateau; and the Mekong Plain. These units each have distinct geology, climate, vegetation, human-use patterns, and potential for development.

Northern Highlands

The Northern Highlands comprise northern Lao PDR as well as Loei province north of Phu Luang in northeast Thailand and the mountains of Chiengrai province in northern Thailand. This is a strongly folded mountainous area where the processes of erosion have carved a highly complex and dissected relief. The topography is steep and rugged, trending mostly northeast-southwest, with mountain ridges of 1,500 to 2,800 meters usually more than 600 meters above the adjoining valley floors, and with most slopes exceeding 30 percent. Most of the rocks are Mesozoic, with linear fold belts of Triassic-Lower Jurassic age. The region has a few relatively large upland plains, such as the Plain of Jars on the Xieng Khouang Plateau. The river valleys usually have small Quaternary alluvial terraces, but the plains of northern Thailand in Chiengrai province are quite extensive, with some 220,000 ha of wet rice fields.

The climax vegetation above 1,000 meters is usually Hill Evergreen Forest, but this forest has been much disturbed by the destructive swidden agriculture practiced by recent maize and opium-growing hill tribe immigrants, resulting in large areas that have become dominated by a few non-native species of unproductive *Imperata* grasslands and broad-leafed weeds (*Chromolaena*). The lower areas are covered with dry evergreen and deciduous forests, including teak forests, with the valley floors invariably given over to crops, especially rice. Human population is sparse, averaging 5-14/km², though populations in valley bottoms where wet rice is grown are considerably higher (as in Chiengrai where the population density is 58/km²).

In the uplands, swidden agriculture is the dominant land use and most inhabitants are hill-tribes (dominated by Khamu, Meo, and Thai Daeng). The potential of the Northern Highlands for hydroelectric power development is substantial, but the potential for irrigation and other agricultural development programs is markedly less than in other parts of the basin (with the exception of the Chiengrai plains).

Annamite Chain

The Annamite Chain extends for 800 kilometers within the basin, from the Langbian plateau in the Sre Pok drainage at about 12°N to the southern edge of the Northern Highlands (19°20'N); the range is 50 to 300 kilometers wide and 500 to 2,500 meters high, dividing the Mekong drainage on the west from the South China Sea drainage on the east. The northwest-southeast trending chain is mostly steep and mountainous in the north and central parts, but forms dissected hills and rolling to hilly plateaus in the south. This discontinuity is a result of an ancient geological history; the Kontum Massif in the south is an eastern outlier of the Pre-Cambrian metamorphic rocks of the Indosinian basement complex. This core of ancient rock has been repeatedly folded, overlain with sedimentary rocks, refolded, and then exposed to an episode of vulcanism about 600-700,000 years ago which resulted in large areas of basalt (including the Bolovens Plateau). Of special interest in the Annamite Chain is the Khammouane Plateau, a hilly karstic limestone area east of Thakhek which is a maze of steep ridges and pinnacles with numerous sinkholes, caves and disappearing streams. This forms the single most extensive area of limestone in the basin.

The rainfall in the Annamites is particularly heavy along the south and west flanks, which receive the full force of the southwest monsoon; some of the inner valleys, however, are considerably drier, resulting in deciduous forests. At one time, the parts of the Annamites which receive over 2,000 mm of rain per year were completely covered in dense rainforest, but many areas have been cleared for swidden agriculture. When this was done at low population levels, it probably represented a sustainable use of the land, but continued burning with increased population density has depleted the soil in many regions (particularly in the south), resulting in wide expanses of unproductive grasslands. Most forests between Dalat and Ban Me Thuot have been cleared, as have the forests along the Sre Pok, extending into Cambodia. The dominant land use is still swidden agriculture, with less than 1 percent of the land used for wet rice. The population is sparse (under 4/km² in the north, 5-40/km² in the south), but is highly diverse in hill-tribe cultures.

Though the Mekong tributaries have more gentle profiles than the streams draining eastwards into the South China Sea, they are broken by many falls and rapids. The northern half of the Annamites is better suited to hydroelectric power development than the less accented relief of the southern half, which provides only limited opportunities for irrigation development in the tributary valleys. Also the basalt areas have some potential for upland crops.

Southern Uplands

The Southern Uplands consist of the Elephant and Cardamom mountains, separating the Cambodian part of the Mekong Plain from the Gulf of Thailand and continuing into Thailand as far as Khao Soi Dao in Chantaburi. The eastern end of the uplands is mostly continuous mountains while the western end is intersected by rolling and dissected plains (now planted with fruit orchards and field crops). The uplands are 500 to 1,700 meters high, formed of the same Mesozoic formations as the Dangrek and Petchabun ranges, with some crystalline massifs. Slope gradients generally are moderate in the north and rather steep and eroded in the south, with only a few steep escarpments.

The Southern Uplands receive a very high rainfall, as much 5,000 mm/year in some places; this results in very dense tropical rainforest and very low human population (under 4/km² in most parts). Hill-tribes are a negligible portion of the population, and even swidden agriculture is very limited (though more common on the drier northern side, which is in a rain shadow). The region has little scope for agricultural development.

Korat Plateau

The Korat Plateau comprises northeast Thailand and adjacent parts of Lao PDR from Pa Mong south to Pakse. The plateau is a large (500 x 500 km) saucer-shaped intermountain basin tilted towards the southeast, the floor of which is 100-200 meters in elevation, with the surrounding mountains reaching 1,400 meters. Hills considered part of the Korat Plateau all consist of Mesozoic sandstones and include the Dangrek range from the Thai-Lao

PDR-Cambodia border west to Khao Yai National Park; the eastern part of the Petchabun Range from Khao Yai north to Phu Luang; and the Phu Phan range in Sakhon Nakhon, an isolated massif in the middle of the plateau. Much of the plateau is underlain by thick deposits of salt formed in the Cretaceous period.

The hills above 1,000 meters are covered in Hill Evergreen and Coniferous forests, with the lower areas in Dry Evergreen, Deciduous Monsoon, or Dry Dipterocarp forests, depending on rainfall and human activity; much of the area is rather dry due to the rainshadow effect of the surrounding mountains. Deforestation has contributed to the recurrent floods and drought which plague the plateau and much of the area is covered with unproductive scrub or grassland vegetation.

The Nam Ngum and Nam Lik rivers rise in the Northern Highlands and form a broad alluvial plain north of Vientiane, at the northern edge of the plateau. The east bank of the Mekong has several major tributaries which rise in the Annamite range, all of which have alluvial valleys in the Korat Plateau, including the Nam Theun, Se Bang Fai, Se Bang Hieng, and Se Done. More than half of the Plateau is drained by the Mun and Chi rivers of northeast Thailand, which experienced some of the earliest development of rice agriculture in the basin and, judging from archaeological sites, must have supported fairly dense prehistoric and early historic human populations. Later populations were considerably smaller, but with recent advances in agricultural technology the population has again increased and much of the plateau supports populations of 80-150/km². About 17 percent of the land is used for growing wet rice, totaling some 3,500,000 ha (over 70 percent of the cultivated area). Other important crops are kenaf (550,000 ha, 11 percent of cultivated area), maize (275,000 ha, 5.5 percent), and cassava (160,000 ha, 3.3 percent) (1973-4 figures). The Korat Plateau offers substantial scope for further agricultural development by means of flood control, drainage, and irrigation on the more productive soils.

Mekong Plain

The Mekong Plain is a vast low-lying area traversed by the Mekong river; however, only a relatively small portion of the Mekong Plain consists of fluvial deposits of the young Mekong. The plain comprises most of lowland Cambodia and the Mekong delta of Vietnam, a small part of southern Lao PDR and a small part of eastern Thailand in Chantaburi and Prachinburi provinces. It is bordered by the Dangrek escarpment on the north, the Elephant and Cardomom mountains on the south, and the southern Annamite Chain on the east. The plain is about 800 kilometers from north to south and 600 kilometers from east to west; it is mostly less than 100 meters in elevation, but a few higher outcroppings are scattered throughout the plain and much of northern Cambodia is characterized by rolling and dissected plains between 100 and 200 meters elevation. The Mekong Plain is the result of erosion and sedimentation; the sediments vary in depth from at least 500 meters near the mouth of the Mekong to only about 30 meters at Phnom Penh, with bedrock outcroppings in isolated hills above the plain in several places.

Leaving the south-eastern edge of the Korat Plateau, the Mekong river plunges over the Khone Falls and reaches the lowlands of the Mekong Plain. In the east, the Se Kong, Se San, and Sre Pok tributaries descend from the Annamite Chain and join on reaching the Mekong Plain at Stung Treng. At Sambor the Mekong crosses its last series of rapids and enters its lower course, characterized by high but narrow natural levees and a broad floodplain. Further downstream, the Mekong receives the waters of a major tributary, the Tonle Sap, at Phnom Penh. The Tonle Sap, fed by the streams which drain into the Great Lake in western Cambodia, reverses its flow in the rainy season, when part of the floodwaters of the Mekong flow up the Tonle Sap into the natural storage basin provided by the Great Lake. Immediately below the Mekong-Tonle Sap confluence at Phnom Penh, the Mekong river divides to form the Mekong and Bassac rivers. Here, between Takeo and Prey Veng, the Lower Mekong Basin is at its narrowest before spreading into its broad alluvial delta. Bordering the sea, at the nine mouths (“Nine Dragons”) of the Bassac and Mekong, the combined action of river deposition and the sea have produced a coastal belt of slightly higher elevation. Deposition in the delta continues to extend the Ca Mau peninsula southwards and westwards at a rate of 150 meters per year in some places.

Although much of the Mekong Plain is subject to annual flooding and the southern tip of the Mekong Delta receives high rainfall throughout most of the year, most of the plain is somewhat dry due to rainshadow effects. Mangrove forests are found along the seaward edge of the delta and fresh-water swamp forest grows in some places around the Great Lake. A large expanse of acid sulphate soils, which is inundated for much of the year, forms the Plain of Reeds, but most of the non-agricultural areas are covered by deciduous forest.

The Mekong Plain is currently the most densely populated part of the basin, with densities over 450/km² in the rice-growing regions of the delta (rice being grown on 50 percent of the land area and 92 percent of the cultivated area). The richest rice-growing areas of Cambodia are also densely populated, especially south of Tonle Sap and on the Battambang plain. The northern and eastern savannas, however, are very sparsely populated, with Preah Vihear, Stung Treng, and Mondulkiri having less than 4 people/km².

The lowlands, particularly the areas of Holocene alluvium, have historically been the most densely populated and productive agricultural part of the Lower Mekong Basin. These areas have substantial water resources development potential for agriculture.

Conclusion

The Northern Highlands, Annamite Chain, and Southern Uplands are generally low in potential for agricultural development and are best used for forestry, swidden agriculture, watershed protection, and protected areas. The Korat Plateau and the Mekong Plain, however, offer very substantial productive possibilities for modernized irrigated agriculture. The potentials for these units will be examined in detail in Part II.

Chapter 2

Climate and Water Resources

Introduction

This chapter discusses two of the most fundamental agricultural resources: climate and water. Though human activities such as forest clearing and burning of fossil fuels may have local effects on the climate, basically the weather is a resource over which people have little direct control. Rather, the climate is a major constraint circumscribing productive human activities. The first half of this chapter discusses how the climate affects crop production in the Lower Mekong Basin.

The second half of this chapter is devoted to the Mekong river, including its flow, sediment load, water quality, salinity intrusion in the delta, and flood patterns. All of these parameters are closely related to agriculture. While little can be done to increase rainfall or decrease droughts, much can be done by people to control the Mekong River's available water in support of sustainable development. This was the major thrust of the Mekong development program.

Climate

The major climate systems operating in the region form part of the global atmospheric circulation; the regional climate is part of a series of mechanisms which maintain the energy balance of the globe through poleward movement of heat energy. The Lower Mekong Basin is situated in the center of the monsoonal region of Asia, characterized by a reversal of wind direction between summer and winter due to differential heating of extensive land and water surfaces. Starting in May, the land surface of China becomes warmer than the surface of the surrounding seas so that a low pressure area develops over the land and a high pressure area develops over the sea; thus the southwest monsoon carries moist air from the Indian Ocean to the land, where it is condensed to fall as rain (75-95 percent of the annual rainfall occurring during this period). Conversely, during the winter, the surface of the sea tends to be warmer than the land so the high pressure area is over the land and the dry continental air masses of the northeast monsoon flow towards the sea, bringing dry conditions to the basin.

Köppen's Climatic Zones

Although the Lower Mekong Basin is often said to lie entirely within the tropical zone and the whole basin is affected by the monsoon, the basin in fact has three major climatic zones

based on rainfall, drought, temperature, and solar radiation (following the Köppen system).

- A. The Tropical Monsoon zone has heavy rainfall most of the year. The short semi-humid period (generally December-February) has less than 60 millimeters mean monthly rainfall, but the ground remains sufficiently moist to support rainforest. Mean temperature in the coolest month is above 18°C, and the temperature has a low annual range which is usually exceeded by the diurnal range. The Tropical Monsoon zone includes most of the Southern Uplands, the southern tip of the delta, and much of the Annamite Chain.
- B. The Tropical Wet and Dry zone has a distinct dry season from December to May, with several months having less than 60 millimeters of mean monthly rainfall. The mean temperature of the coolest month is above 18°C, and diurnal and annual temperature ranges are moderate. This is the dominant climatic zone in the basin, comprising virtually all of the areas of the Mekong Plain and Korat Plateau which are proposed for agricultural development programs.
- C. The Subtropical Wet and Dry zone has a distinct dry season during December to May, with the minimum monthly rainfall only one-tenth that of the wettest month of the wet season. The mean temperature of the coolest month ranges from 18°C to 13°C, and the annual mean temperature is relatively low (though the diurnal range is usually greater than in the Tropical Wet and Dry zone). This climatic zone is found primarily in upland areas of the Northern Highlands and in the highest parts of the Annamite Chain.

Temperature, Rainfall, and Solar Radiation

Illustration 1 shows selected climatic data for each of the five physiographic units in the basin. The marked annual variation in rainfall is notable, with the variation sometimes exceeding the mean. Based on 60 years of rainfall records in three provinces in northeastern Thailand, annual rainfall is widely variable, the 5-year moving average is somewhat less variable, and the 60-year period clearly showed a fluctuating trend (Kawai, 1974). Topography, wind currents, and random thunderstorms also lead to marked local differences, so the climatic data presented should be taken only as indicative of the general climatic picture.

Since plants create new matter through the process of photosynthesis, the amount of solar radiation is a vital parameter for agriculture. Geographical, seasonal, and diurnal variations of global solar radiation have been surveyed for Thailand (Exell and Saricali, 1976), including detailed maps of the geographical distribution of solar radiation prepared from data on cloudiness at 44 stations, duration of sunshine at 18 stations, and linear regressions relating radiation to sunshine at Chiangmai and Bangkok. The highest mean values are above 470 cal cm⁻² day and are widespread during the spring dry season; the lowest mean values are below 350 cal cm⁻² day in restricted localities with heavy rainfall in autumn.

Regional temperature, rainfall, and solar radiation data generally are not good indicators of micro-climate. Forested areas, for example, generally experience less temperature

ILLUSTRATION 1: CLIMATIC DATA FOR SELECTED LOCALITIES IN THE LOWER MEKONG BASIN									
	Location	Precipitation			May-Oct.	Temperature (°C)			Elevation (metres)
		Maximum	Minimum	Mean		Maximum	Minimum	Mean	
I.	Northern Highlands:								
	Luang Prabang (L)	1879.0	493.1	1244.3	84.8	48.8	0.8	24.2	287.0
	Chiengrai (T)	2302.9	1120.2	1628.1	89.7	41.3	2.0	24.8	382.0
	Xieng Khouang (L)	2501.9	1232.3	1715.9	75.1	33.9	3.0	19.4	1149.0
II.	Annamite Chain								
	Boloven (L)	na	na	3742.5	87.7	na	na	na	1200.0
	Pleiku (VN)	3154.0	1937.0	2450.9	92.7	34.7	5.6	21.8	800.0
	Ban Me Thuoi (VN)	2234.0	1480.0	1733.1	87.1	39.4	7.4	23.5	536.0
III.	Southern Uplands								
	Kampot (C)	3120.0	1434.0	1905.0	79.1	37.2	4.7	27.0	5.0
	Val d'Emera ?(C)	na	na	5124.1	76.2	na	na	na	950.0
	Chantaburi (?)	4158.2	1739.6	2837.8	88.5	40.8	8.9	27.2	6.0
IV.	Korat Plateau								
	Udonthani (T)	1998.5	1367.0	229.4	87.5	43.9	2.5	26.8	181.0
	Korat (T)	2415.6	730.7	1180.6	81.8	43.4	4.9	27.1	189.0
	Savvanakhet (L)	1181.9	1182.5	1395.5	87.8	42.4	5.6	26.0	140.0
V.	Mekong Plain								
	Battambang (C)	1970.0	919.0	1376.4	81.0	41.0	10.4	27.0	22.0
	Siam Reap (C)	2056.0	466.0	1431.7	76.2	20.3	9.5	26.7	21.0
	Pakse (L)	2582.4	749.4	1929.5	93.9	39.7	7.0	26.4	96.0

Note: T = Thailand; L = Lao PDR; C = Cambodia; VN = Viet-Nam

variation, both daily and seasonally; humidity is higher in forests than in cultivated areas; and rainfall as measured on the ground is less in forested areas than in comparable cleared areas because of the interception of rain by the various canopy layers of the forest (though rainfall is actually higher in forested areas due to differences in surface albedo). The areas remaining in forest are generally in the uplands, where elevation also has a marked effect on the climate. The temperature is about 1°C cooler for each 200-300 meters in elevation and rainfall is higher in uplands because of local convection currents and wind effects. The distribution of hill evergreen forests in the otherwise dry Korat Plateau is a good indication of higher and more evenly distributed rainfall in the upland areas.

Generally, the highest rainfall areas are in the Southern Highlands, the central and southern parts of the Annamite Chain, and the western coastal areas of the delta (2,000-4,000 mm/yr). Large areas in the Korat Plateau and the regions surrounding the Great Lake are shielded by the coastal ranges and receive less rainfall, averaging between 1,000 and 1,400 mm/yr.

Crops and Rainfall Pattern

Since yields of most crops grown under traditional conditions are heavily influenced by climate, especially the annual rainfall pattern, analysis of relationships between climate and agricultural production is exceedingly important. Kawai (1974) has analyzed rainfall distribution and frequencies to determine the relationship between paddy yields and rainfall and to identify regional irrigation priorities.

It should be mentioned here that total annual rainfall, as measured at rainfall stations, is sufficient to grow most crops anywhere in the lowlands and uplands of the basin. However, the distribution of rainfall during the growing season tends to be erratic and many years bring serious damage from floods or droughts. Since tropical soils do not have a particularly high water-holding capacity, especially for paddy rice, the farmers have traditionally leveled the paddy fields and surrounded them by low bunds (embankments) in order to increase effective rainfall by keeping the water on the field.

For the purpose of analyzing in greater detail the variability of rainfall on paddy production in northeast Thailand, a model simulation was designed by the Mekong Secretariat (Kawai, 1974), using the data from the main provincial rainfall stations over a period of about 20 years (varying somewhat with the station) for the 15 provinces in the northeast, and taking into account constants such as maximum flood depth (in the fields between the bunds, determined by the height of the bund), water holding capacity of the soils, and evapo-transpiration. The main result of this analysis is the determination of the frequency of drought days (total and maximum continuous), a drought day being defined as a day during which water depth and available soil moisture are zero. This shows that in the rainshadow of the Petchabun Range, for example, the average total of drought days during the growing season is as high as 100 (even though total rainfall was 1,200 millimeters or higher) while the maximum number of continuous drought days during the transplanting and the flowering periods is as high as 40 and 25 respectively.

The effect of these drought periods on paddy production can be analyzed in a time series, since paddy growing is relatively insensitive to price fluctuations, weather being the major variable. If adverse weather conditions are met in the early part of the growing season, the planted area is reduced; under adverse conditions during the flowering season, the harvested area is reduced. Drought during either of these periods will thus result in reduced rice production, and available statistics are sufficiently accurate to allow analysis of these phenomena.

The reduction rates were plotted against paddy yield over a period of twenty years, showing a clear trend, but with a considerable scatter of points. This is not surprising, since available statistics are arranged not according to physiographic units but to administrative units, while other variables such as varieties, local differences in the cropping calendar, flood damage, pests, and disease have not been taken into account.

The conclusion is that climate is a serious limiting factor in rice growing in many parts of the Lower Mekong Basin and particularly in northeast Thailand. As shown above, rainfed paddies tend to be erratic in performance; for sustained yields, they need supplemental water amounting to 280 millimeters in an average year and 800 millimeters in a drought year (occurring once in every ten years). The total demand in northeast Thailand for supplemental water on irrigable paddies in drought years is about 14,000 million m³, almost equal to the total watershed runoff in such years. The maximum water supply potential is about 15,000 million m³, assuming it is feasible to store and distribute this amount. Of the 15,000 million m³, some 4,500 million m³ was being stored in 1970, enough for use on about 80,000 ha in the rainy season and 250,000 ha in the dry season. For diversified or double cropping, the total demand could be as high as 41,000 million m³, including the

14,000 million m³ for rainy season paddy. The deficit of 27,000 million m³ will have to come from another source, and the only feasible source is the Mekong river.

Water Resources: The Mekong River and Its Tributaries

The Mekong river is the longest river in southeast Asia, crossing six countries in its 4,200 km course from its sources in the eastern part of the great plateau of Tibet to its mouths in the South China Sea. The Mekong drains an area of about 744,000 km², of which about 606,000 km² or about 80 percent is in the lower basin, beginning where the river leaves China and becomes the border between Myanmar and Lao PDR. The lower basin thus includes almost the whole of the Lao PDR (202,400 km²); the Korat Plateau, the northern tip, and a small part of the southeast of Thailand (184,230 km²); nine-tenths of Cambodia (154,730 km²); and the western flank and southern tip of Vietnam (65,170 km²).

Entering the lower basin, the Mekong has an average annual flow of about 90,000 million m³ (less than 20 percent of the annual flow at the mouth). By the time the river discharges into the South China Sea, over 2,500 kilometers downstream, the average annual runoff has increased to nearly 550,000 million m³. Estimates of the annual peak flow and annual runoff at some key locations on the mainstream are given in illustration 2.

The sources of the surface flow within the basin are quite disparate, with about 70 percent of the total flow coming from the Northern Highlands and Annamite Chain, comprising about 50 percent of the area of the lower basin. By contrast the Korat Plateau and the Mekong Plain, comprising 25 percent of the land area, contribute only 10 percent of the total flow, a reflection of the lower rainfall in these areas.

The seasonal variation of Mekong flow is pronounced, with monthly average flows ranging from 0.12 to 2.89 times the long-term mean flow. This variation generally corresponds to the monsoon rainfall pattern except in the upper basin where snow-melt makes it more uniform. The river begins to rise following the onset of the rains in May and attains its maximum level in August or September for upstream stations, and in September or October for downstream stations. It then falls rapidly until December, recedes slowly during the dry period, and reaches its lowest level in late April.

The wide variability of annual rainfall is reflected by peak annual flows. Records covering 60 years at Vientiane indicate that the annual peak may be as high as 25,000 m³/s, in a flood year, and as low as 11,300 m³/s in a dry year; in a “normal year”, an annual peak flow on the order of 17,100 m³/s is to be expected, which could be confined within the banks of the river. Along the stretch between Vientiane and Nong Khai the river occupies extensive floodways, 10 km wide in Thailand and 5 km wide in Lao PDR. The total maximum overflow in these floodways for the 1966 flood was estimated at 3,700 m³/s against 21,300 m³/s moving in the main channel.

The magnitude of the annual peak flow increases progressively moving downstream to

ILLUSTRATION 2: MEKONG ANNUAL PEAK FLOWS ^a					
Station	Distance from outlet km	Drainage area above station km ²	Annual peak flow m ³ /sec	Annual runoff millions m ³	Annual suspended sediment load million tons per year
Chian Sen	2,330	186,000	10,700	89,870	67.21
Luang Prabang	2,010	264,000	16,600	132,510	-
Vientiane	1,584	301,000	16,300 ^b	144,540	109.14
Thakhek	1,216	366,000	26,100	241,530	-
Mukdahan	1,125	387,000	29,200	263,760	-
Pakse	869	540,000	39,000	301,850	132.20
Stung Treng	668	624,000	52,700	450,310	-
Kratie	545	636,000	53,000 ^b	452,890	-
Phnom Penh	332	653,000	43,700 ^c	428,530 ^c	97.55 ^c
Phnom Penh	332	653,000	50,000 ^d	512,000 ^d	-
Outlet Mekong + Bassac	0	744,000	44,000 ^d	550,000 ^d	-

^a National sources data consolidated by ECAFE Division of Water Resources and Mekong Secretariat (Lower Mekong, Hydrologic Year book)

^b Channel effects

^c Channel flow only

^d Channel flow plus estimated overbank spill (1961)

about Kratie where it reaches a peak as large as 53,00 m³/s. Below Kratie, the river overtops its banks when rising above 7 meters MSL, so peak flows of the mainstream are reduced. Extensive overbank flooding takes place on both banks, but the by-pass flow on the western bank escapes towards the Great Lake. The main channel also adds its flow to the waters of the Great Lake through the Tonle Sap during the rising flood, and the action is reversed during the later flood recession. The Great Lake thus acts as a natural flood retention basin without which the delta would be flooded to a greater depth, and more frequently.

At Phnom Penh, part of the Mekong water begins to flow into the Tonle Sap sometime between April and June and continues until its recession in October. The total inflow to the lake from the Mekong river is about 60,000 million m³. The lake outflow during the rest of the year is about 84,000 million m³, indicating that an additional 24,000 million m³ of water comes from the lake catchment. Just below Phnom Penh, the Mekong divides into two branches, the westerly being the Bassac and the easterly being the continuation of the Mekong itself.

Between Chau Doc and Long Xuyen in the Vietnamese delta, the Bassac and the Mekong are connected by a channel (Vam Nao) through which about 40 percent of the Mekong flow turns into the Bassac during the annual crest stage, when the flow at Long Xuyen rises above 12,500 m³. The result is that the smaller flow of the Bassac (normally 25 percent that of the Mekong) becomes equalized with that of the Mekong below the connecting point. Some overland flow also escapes easterly across the Plain of Reeds and drains into the Vaicos.

While the peak floods are quite spectacular and potentially destructive, the low flow is

usually the limiting factor in determining the scale of development of an agriculture project and its compatibility with existing or future developments. Based on available hydrologic data, the lowest flow and the average annual low flow of both mainstream and selected tributaries are indicated in illustration 3. The lowest flows on record at Chieng Saen, Vientiane, and Kratie on the main river are 543 m³/s, 701 m³/s and 1,250 m³/s respectively, while the average annual low flow at the same stations are 639 m³/s, 1,004 m³/s and 1,764 m³/s.

The great variation in flow between seasons and between years indicates that substantial storage is required in the basin to provide more uniform flows for irrigation, dry-season water supply and power generation as well as for flood control and navigation improvement.

Sediment

The sediment load of the Mekong river is somewhat lower than for other major Asian rivers, such as the Irrawaddy, Yellow, and Yangtze. The suspended sediment load, analyzed over a period of 12 years by an 89-station sampling network, increases from 67 million tons at Chieng Saen to 109 at Vientiane and 132 at Pakse. After Kratie, as the river slows down, the sediment load decreases to 97 million tons at Phnom Penh. Peak concentrations are highest in the upper reaches of the river, averaging about 1,500 parts per million (ppm) near the Myanmar border, but decrease to about 1,000 ppm at Pa Mong and about 500 ppm in approaching the delta. Peak sediment loads occur in August in the upper reaches, but two or three months later in the downstream stretches. Sediment loads are at a minimum, less than 50 ppm, in February.

ILLUSTRATION 3: LOW FLOWS IN MEKONG RIVER AND TRIBUTARIES ^a						
River	Gauging station	Drainage area km ²	Recorded lowest daily flow	Average annual low flow		Low flow ^b m ³ /s
				m ³ /s	litres/s/km ²	
Mainstream	Chian Saen	189,000	543	639	3.38	1,539
Mainstream	Vientiane	299,000	701	1,004	3.36	2,301
Mainstream	Kratié	646,000	1,250	1,764	2.73	7,045
Nam Ngum	Ban Pak Kanhoung	14,300	29	56	3.92	301
Se Bang Fai	Se Bang Fai	8,560	8.2	16.5	1.93	280
Se Kong	Ban Khmuon	29,600	77	192	6.49	710
Nam Chi	Yasothon	43,100	1.6	4.3	0.10	129
Nam Mun	Ubon	104,000	3.2	9.1	0.69	324
Stung Sen	Kompong Thom	14,000	1.8	4.3	0.31	99
Dak Bla (Upper Se San)	Kantum	3,056	3.8	-	-	-
Ea Krong (upper Srén Pok)	Drayling	8,880	18.2	-	-	-

^a National sources data consolidated by ECAFE Division of Water Resources and Mekong Secretariat. (Lower Mekong, Hydrologic Year)

^b Low flow is arbitrarily defined as 50 percent of the long-term average flow, or less.
Source: Indicative Basin Plan, Mekong Secretariat, 1970.

It is often said that these sediments, deposited in the delta by the annual floods, perform a vital service in adding nutrients to the Delta soils. In order to establish the relationship between sediment deposition and soil fertility, the Mekong Committee sponsored an investigation into the mineralogical and chemical composition of the river sediment as well as adjacent soils in the delta (Uehara *et al.*, 1974). Small but significant differences in mineral and chemical composition were measured between river sediment and delta soil, with the sediment samples higher in minerals, such as mica, hematite, kaolinite, and feldspar, but still having an excessively high percentage of quartz. The fresh sediments contained, as expected, a somewhat higher percentage of extractable elements, such as calcium, magnesium, potassium and phosphorus.

Based on these data, the quantity of phosphorus, potassium, magnesium, and calcium added to the soil each year by sediment indicates that the amounts of nutrients added to the soil were quite small, not surprising given the low fertility of the parent materials through which the river flows. The readily soluble nutrient annually added to a one hectare area was estimated to be approximately 1 kg of phosphorus, 3.2 kg of potassium, 4 kg of magnesium, and 50 kg of calcium. Even if these computed values were doubled, the sediment deposit could not significantly increase the fertility of delta soils. This is not to deny the importance of the fluviatile soils for agriculture, but their importance is based more on their physical properties rather than annually-deposited soil nutrients.

Water Quality for Agriculture

Systematic sampling and analysis of the quality of water in the Mekong river was initiated in 1972, using some 29 stations. It was found that the water quality changes very little going downriver from Myanmar to the South China Sea. The water is low in total dissolved solids, hardness, sodium, and chloride, with the concentration of dissolved minerals being reduced by about one-third during the wet season as compared with the dry season. The dissolved mineral quality is thus considered excellent for practically all agricultural, community, and industrial purposes

Salinity Intrusion

Intrusion of saline water from the South China Sea and the Gulf of Thailand affects much of the downstream part of the delta, especially in the dry season when the Mekong discharge and local rainfall are reduced. Inundation by saline water occurs only on a limited land area during high tides. Of the 3.7 million ha in the delta, about 0.5 million ha are sufficiently affected by salt water intrusion to place limitations on agricultural development.

The salinity of the water in the various branches of the river was measured at the banks of the sea arms during the period 1935 to 1942. A more comprehensive evaluation of delta salinity was made under the auspices of the Mekong Committee from 1972 to 1974 by the Netherlands Delta Development Team (NDDT) engaged in preparing a basis for a master plan for agricultural development in the delta. The analysis used available information

of river discharges, especially downstream from Phnom Penh, on bed configurations (especially minimum depths of the bars at the mouths of channels, which are important factors in estuary salinity intrusion), and on water movement by tidal action. Salinity distribution curves were approximated for all the branches of the river, based on the monthly averages of the results of measurements at the banks during the years 1935-1942 and on conductivity measurements taken during 1973. The analysis according to this statistical method yielded penetration lengths into the branches varying from 20 to 65 kilometers.

Flood Patterns

With the aid of the satellite imagery mapping program operated by the Mekong Secretariat, a general overview of the flood and drought problem in the lowlands of the Lower Mekong Basin was obtained. It is apparent that towards the end of the rainy season in September-October, some 10 million ha of lowlands are covered with water at various depths of flooding. These flooded areas include the entire Mekong delta, the lowlands around the Great Lake, the Korat Plateau along the major rivers and streams, large parts of the Vientiane and Se Bang Fai plains, and parts of northern Thailand.

Four variables – water velocity, rate of rise, depth and duration of the flood – act independently relating to flood damage in different localities. Throughout two thousand years of farming experience, rice varieties and cultivation techniques have evolved which are capable of considerable adaptation to these variable conditions.

While much of this flooding has benefits for rice growing, it is generally unregulated, both in timing and in quantity, so the use of the floodwaters is seldom optimal; rice production in the flood zones remains hazardous and higher levels of production can only be reached with flood mitigation. Complete flood control on a river the size of the Mekong is not possible, and is probably not desirable, since flooding with neutral river water is beneficial to many delta soils. However, considerable flood relief is locally possible through the construction of storage dams on the tributaries, as has been shown by the Lam Pao and Nam Pong dams for the valley of the Chi river in northeast Thailand. But the construction of very small reservoirs, such as the tanks in northeast Thailand, of which some 150 have been built, has little impact on flood mitigation (Kreuze, 1975).

Many of the same lands affected by floods also suffer from serious lack of water, even for domestic use, toward the end of the dry season. In the coastal belt, much of the water is saline, but further inland it is highly acidic. Around the Great Lake some waters contain humic acids which become highly concentrated in the dry season. In many of the areas flooded in northeast Thailand during the wet season, virtually no water is available during the dry season. Thus the flood and drought syndrome occurs at various locations in the basin in every year, although the severity and location of damage is variable.

Conclusion

In summary, the basic problem for agricultural development in the lowlands can be simply stated in one single concept: water management, including management of the tributaries to the extent they can contribute and management of the mainstream Mekong for realizing the lower basin's full development potential.

Chapter 3

Basin Soils and their Major Limitations

Introduction

Once flood and drought conditions are improved through water control, soil characteristics become the major constraint for agriculture, so the Mekong program placed great emphasis on basin soil studies. A General Soil Map and, based on the soil map, a General Land Capability Map were prepared to a scale of 1:1,000,000 (van der Kevie, 1972). The soil map incorporated all surveys made to date in the riparian countries and classified the soils in the international FAO/UNESCO nomenclature. With the aid of satellite imagery from LANDSAT II, this map was improved and modified into a Pedo-Morphological Map of the Lower Mekong Basin (Ceruse, 1977).

While these small-scale maps tend to mask local variations, they do give an indicative picture for preliminary agricultural planning. They are especially useful for screening of alternative agricultural systems, both rainfed or irrigated, enabling planners to delete alternatives which do not correspond to the suitability of the land and thus will invariably result in failure or low return on investment.

Processes of Soil Formation

The soils of the Lower Mekong Basin are formed as a function of parent material, climate, hydrology, and age. Vegetation, while important in the reduction of organics and to control erosion, plays a much smaller role as a soil-forming factor in the tropics, since humus formation is not a permanent characteristic (with a few exceptions).

Regarding composition of the parent materials, alluvials comprise about 10 percent, soils on sandstone comprise about 80 percent, and soils on limestone and basalt comprise about 10 percent of the total. These three groups of parent materials give rise to soils with quite different characteristics.

The high rainfall typical of the basin causes a progressive leaching of all soil types, resulting in nutrient-deficient podzolic soils of various degrees of acidity. Another climatic feature, although it involves a much slower process than the leaching of soluble elements, is the separation of silica from the sesquioxides in the clay minerals. This process, enhanced by annual wetting and drying in the monsoonal climate whereby the sesquioxides precipitate at a certain depth (often conditioned by the fluctuating groundwater table), is called “laterization”. The alluvial layer of sesquioxides hardens to laterite when dry, forming a hardpan which is impossible to cultivate by normal agricultural techniques.

A somewhat similar process takes place in wet paddy fields caused by annual extreme reduction (when wet) and oxidation (when dry) whereby the clay minerals tend to break down after leaching. This process, which results in topsoils with very low cation exchange capacities (CEC) is called “ferrolysis” (Brinkman, 1970). The cation exchange capacity of a soil is an empirical value expressing the amount of cations, in milli-equivalents, that 100 gram of dry soil can absorb.

The CEC value is very important, not only as a measure of the inherent fertility of the soil, but of the efficiency of fertilizer applications.

The hydrology of the plains and lowlands results in hydromorphic soils with pronounced mottling (“gley”). However, this hydromorphy (with the exception of some very poorly drained areas in the delta) rarely results in a humiferous topsoil, so these hydromorphic soils are called “low-humic gley soils”.

Age is also important, since all these soil forming processes become more pronounced and evolve over time. Many of the surfaces of the Korat Plateau and the Mekong Plain are very old; the lateritic hardpans of the Sakhon Nakhon basin in northeast Thailand, the Savannakhet plain in Laos, and the plains of Cambodia, probably reflect ancient groundwater levels. In the Mekong Plain of Cambodia, particularly in the eastern part, soils with a clay hardpan are common, forming the so-called “planosols” or “plinthitic” soils.

Soil Types

The General Soil Map classifies the basin soils into 29 types. However, for the purpose of the present discussion, these 29 types have been grouped into eight categories, representing their dominant characteristics. Four of these are lowland soils (hydromorphic) and four are upland soils (well drained).

Lowland Soils

The lowland soil groups occupy the major portion of the Korat Plateau and the Mekong Plain, the areas with the most potential for agricultural development. These soils are primarily hydromorphic with gley characteristics at shallow depths.

a. The Coastal Complex

The coastal complex is an expanse of land along the sea coast of the delta consisting of a series of ancient, sandy beach lines interspersed by clay depressions. In general, the soils of the coastal complex are clayed with a low organic matter content and high CEC. They are usually slightly acid, though some are saline (with a pH of about 7) or alkaline (with a pH above 8.5).

Although these soils are generally fertile, salt water intrusion is the dominant feature as far as soil and land use are concerned. The extensive saline swamps are at present not suitable for intensive cropping, being mainly used for shrimp farming, salt production and some coconut growing.

The soils of the high tidal flats experience only minor salinity problems. For example, tree crops, typically having deep root systems, generally cannot be grown in these locations, with coconuts, having fibrous roots, being a notable exception. These soils are well suited for paddy cultivation because of their good surface runoff drainage and heavy texture. However, several factors related to poor soil structure make them less suitable for crops other than rice. Rice cultivation necessitates puddling of the soil, and chemical factors in these marine soils allow the clay particles to disperse easily, causing a surface crust to develop. Further, high temperatures during the dry season cause destruction of most of the organic matter. Thus these soils are poorly aerated and have a low permeability, and drainage must be improved if crops other than rice are to be grown.

A small area of the coastal complex is occupied by light-textured soils on old beaches. These sand ridges are low in organic matter and CEC and are moderately acid. In general, their fertility is low, but these soils are well suited for dry-foot crops (that is, crops other than paddy rice) because of their good aeration and drainage conditions. Their low water-holding capacity makes these soils very sensitive to drought. At present, the higher sand ridges serve as settlement areas with gardens on which various crops are grown, including fruit trees, though for deep-rooting crops salinity damage may occur due to capillary rise.

The low tidal flats generally have soils which become saline during the dry season, so accumulated salts must be washed away (some 100 to 150 mm of rainfall may be required) before rice can be planted. High bunds are also required to prevent salt water intrusion during high tides. The low tidal flats in the southwestern part of the coasts complex are wet throughout the year, with a rather abundant weed growth during the dry season. Salinity problems here are less severe than elsewhere in the coastal complex due to a very high rainfall and a relatively short dry season.

In the western part of the coastal zone, peat layers have developed from 40 to over 200 cm thick. Little information is available about this area and consequently the agricultural potential is largely unknown.

b. Delta Lands

These soils are clayey with varying organic content, depending on past and present drainage conditions. The clays have been deposited under special conditions, including regular inundation by brackish water, presence of mobile iron, a slow sedimentation rate, a dense vegetation with a fibrous root system producing large amounts of organic matter, and prevailing lack of free oxygen (NDDT, 1974). Under these conditions, sulfates in the brackish water are reduced to sulfides by bacterial activity. The sulfides accumulate in the clay sediment over a depth which probably exceeds several meters in most areas. When

the soil dries, the chemical processes are reversed and sulfates are released which makes the soil strongly acid.

Active oxidation of sulfides occurs roughly between the highest and lowest groundwater levels, forming various acidic iron sulfate compounds, notably lemon-yellow mottles of the mineral “jarosite”. These acid compounds will react with the clay minerals in the soil, thereby releasing aluminum which is toxic to the crops. At present, the very acid soils with a jarosite layer within 50 centimeters of the surface are not usually cultivated, but soils with a jarosite layer at depths between 50 and 120 centimeters are typically under rice cultivation.

Paddy is the most suitable crop for sulfide soils because the topsoil loses its acidity after about 2 weeks of water saturation, due to chemical reduction processes. The main hazard occurs when droughts cause drying of the soil and sulfates are formed as a result of oxidation; the pH may drop to below 4 or even 3 at these times.

The hazard of soil acidification has been the key factor in determining the agricultural potential of these delta soils. Under irrigation, acidification can be minimized or even prevented by using neutral irrigation water and by maintaining a high groundwater table during the dry season (for example, by growing a second rice crop which is not harvested before mid-March). According to the Delta Team Report (NDDT, 1974), reclamation should begin near the main rivers and proceed gradually away from the rivers to guarantee irrigation water of good quality and to prevent acidification of floodwater. Careful liming will also be necessary to reduce residual acidity.

c. Flood Plains

The soils of the riverine plains are mostly clayey except for small areas occupied by river levees. These plains are formed from flood-deposited sediments and because of their recent origin, their soils are mostly undifferentiated, with only the A-horizon distinctly developed (through accumulation of some organic matter). The clayey soils of this type may be somewhat more suitable for dry-foot crops than the marine clays because of their better physical properties. The hazards of salinization of alkalinization do not exist (except for some minor areas).

In general, the levee soils have a low organic content, medium-high CEC, and a slightly acid to neutral pH. While they are currently only moderately fertile, they have high potential with good water and soil management. They are well-suited for dry foot crops including fruit trees, though flood conditions are a limiting factor.

The soils of the backswamps are low in organic matter, have a high CEC, and are slightly too moderately acid. Most are moderately fertile, but have high potentials with improved management. They are well suited for paddy cultivation unless flood conditions interfere.

Around the Great Lake, floods normally rise more smoothly, allowing the cultivation of transplanted or floating rice, depending on the depth of the flooding, although in some areas flash floods may occur. Damage from flash floods can be greatly reduced by regulation

of tributaries. In general, the flood plains will have a much higher potential for agriculture once flooding is brought under control.

d. Groundwater Complex

These soils are characterized by a fluctuating groundwater table which rises to or near the surface in the wet season. The typical groundwater soil is a silt loam with low clay content in the surface horizon, the clay content increasing with depth. When used for paddy cultivation, a dense ploughpan usually develops. The organic matter content is low to very low and the pH and CEC are low. Under conditions of alternating reduction and oxidation of iron, a large part of the clay minerals have been destroyed in these paddy soils. This accounts for the low clay content in the topsoil and the very low CEC of the clay fraction in the topsoil (Brinkman, 1970).

Aluminum is released with the decomposition of clay minerals and becomes exchangeable on the absorption complex. Being a trivalent cation, aluminum is, under non-saline conditions, usually absorbed on the clay-humus complex, leaving a very low aluminum concentration in the soil moisture solution. However, with heavy fertilizer applications aluminum may be exchanged by cations from the fertilizer sufficiently to become toxic to young plants. Under continuous reduced conditions aluminum will precipitate, so that when heavy fertilizer dressings are required, they should be applied about two weeks before transplanting.

In the case of acid soils, which occur in northeast Thailand, free aluminum will replace other cations on the clay complex as soon as the soil moisture solution is diluted. Rice should be transplanted after two weeks of inundation or lime should be applied, though over-liming should be avoided because of the low buffering capacity of the soil (Brinkman and Dieleman, 1977).

In northeast Thailand and the Vientiane Plain, the groundwater areas occupied by this complex contain some salts, especially sodium chloride, which may cause salinity problems under certain hydrological circumstances, as in areas which are not annually flooded during the wet season. The salts originate from residual salinity of certain geological formations, accumulating during the dry season through capillary rise and evaporation of groundwater (Takaya, 1974). These saline areas can be found on slightly higher spots surrounded by paddy soils and on transitional soils between low and higher levels, which occur in northeast Thailand, particularly in the Mun-Chi Basin. Reclamation of saline soils should include leaching with improvement of the drainage conditions.

Soils of the groundwater complex originate mostly from pre-weathered acidic rock, with the soils of the Battambang Plain (which originate from basic rock) as the main exception. Normally a texture-B horizon has developed here, but processes like those described for the other paddy soils of the groundwater complex do not take place. These are fairly productive soils, well suited for paddy and possibly for upland crops when drainage is improved and the crop is protected against flash floods.

Upland Soils

The upland soils have a groundwater table deep below the surface, even in the wet season, although some transitional areas resemble the soils of the groundwater complex. In northeast Thailand the lands of these transitional areas are in bunded rice fields, but they are very drought-susceptible and of low productivity.

a. Podzolic Soils

These are well drained, in low to very low clay and very low organic content when cultivated. They also have a low CEC and low to very low water holding capacity. They are marginally suited for annual upland crops, with the exception of kenaf and cassava, which can be grown for several years until soil nutrients are exhausted. However, with modern management and fertilizers many of these podzolic soils would be capable of sustained agriculture. Typical for the lower phase of the podzolic soils is the abundance of termite mounds.

b. Red and Black Soils (Derived from Plateau Basalts and Limestones)

The red and black soils form a special group among the upland soils. While the group has wide variation, these soils are by far the best for upland rainfed agriculture, both for annual and perennial crops. In the higher rainfall areas at lower latitudes in Cambodia and Vietnam, rubber is the dominant crop. At higher latitudes, such as in northeast Thailand, the main cash crops are corn (maize) and cotton. In northern Thailand and northern Laos, opium is grown on these soils. In all cases, the red and black soils are capable of sustaining a variety of diversified farm systems under rainfed conditions.

c. Lateritic Soils

Lateritics, which occur on the Korat Plateau and in the Mekong plain, can be considered to be fossil groundwater soils with an oxidized and hardened laterite at shallow depth. They are generally not suited for field crops, but grasslands could be quite productive.

d. Mountain Soils

These are undifferentiated, since pedological surveys in the Lower Mekong Basin have usually been confined to the plains. Satellite imagery does not provide sufficient information to identify mountain soils, even on small-scale maps. However, the mountain soils are generally freshly weathered from the underlying rocks and therefore their characteristics reflect the geology. Geological maps can thus give an impression of the soil types, suggesting that the vast majority of the mountain soils are shallow, were derived from sandstones and acid igneous rocks, and have a limited potential for agriculture, even for shifting cultivation. Their carrying capacity for growing livestock is also low; but a few of these soils are favorable for growing a wide variety of high value crops, favored by special micro-climatic conditions, including fruits, vegetables, herbs and spices, and insecticidal plants.

Conclusion

The soils of the Lower Mekong Basin have many important limitations for agriculture, typically including low organic content, high acidity, saline intrusion, and periodic damage from floods and drought. Mitigation of these constraints is one of the key objectives of the agricultural development efforts of the Mekong program.

Chapter 4

The Natural Vegetation

Introduction

Previous chapters have discussed geology, climates water regime, and soils, which leads to the floristic expression of these parameters: the natural vegetation. This chapter first describes the natural types of evergreen and deciduous forests, leading into a discussion of forests as a resource and showing that forests have played a key role in the socio-economic history of the Lower Mekong Basin. The basin forests are now seriously threatened by over-utilization, especially in Thailand.

Definition of Natural Vegetation

In the minds of many, “natural vegetation” means “virgin forest” where human influence is slight or lacking. However, such vegetation probably no longer exists in significant amounts anywhere in the basin. Due to the long-term activities of swidden agriculturalists (who have been working the basin forests for hundreds of generations), most forests in southeast Asia have been cleared at some time in history, and most have been cleared many times. No country in southeast Asia appears exempt from extensive modification of its forest by man and fire, from low alluvial plains with densely populated wet-rice areas to high altitudes in mountainous tracts (Wharton, 1968). “Even in the mountains of Laos there are few areas of forest that have not been partially cleared for shifting cultivation at some former time” (Talbot, 1973). Spencer (1966) concludes “Most of the mature forests of the Orient today are not virgin in the proper sense, but merely old forests that have reached a fairly stable equilibrium of ecological succession after some early clearing by human means”.

Perhaps “natural vegetation” can be best defined as any non-agricultural vegetation, with “climax vegetation” being the idealized mature forest (or grassland in some seasonally inundated areas) that would occur if human influence were eliminated.

Forest Types

Natural vegetation can be divided into two major groups, forests and grasslands. The basin forests can be further divided into two main groups, evergreen and deciduous, though it should be remembered that these are not exclusive; an evergreen forest may have many species typical of deciduous forests and a deciduous forest may have some evergreen

trees, especially in the understory. Each of the major groups is further divided into types, with evergreen types including Tropical Rainforest, Hill Evergreen Forest, Coniferous Forest, Fresh-Water Swamp Forest, and Mangrove Forest. The main deciduous types are Deciduous Monsoon Forest and Dry Dipterocarp Forest. In addition, early successional stages of fire climax forests may contain trees as a minor factor.

Tropical Rainforest

The tropical rainforest of southeast Asia is the most complex terrestrial ecosystem known to science, with an amazing diversity of plants and animals. Rainforest occurs from sea level up to 1,000 meters in areas which have an annual rainfall of at least 2,000 mm, fairly evenly distributed throughout the year. A drier sub-type is found in areas with a seasonally dry climate but with high rainfall during the rainy periods. Rainforests are formed on a wide variety of soils, but are especially common on red-yellow podzolics (“Orthic Acrisols”). These heavily leached, acidic tropical soils are notoriously infertile, since most nutrients are contained in living vegetation and decomposers. A luxuriant tropical forest can grow on almost sterile soil and attempts to grow crops on such soil inevitably exhaust the residual fertility very quickly.

The foliage of tropical rainforest typically forms four major layers: (i) a ground flora of sparse low-growing herbs, shrubs, and ferns, with grasses occurring mostly along streams or in temporary forest clearings; (ii) a relatively open layer just above the ground, formed of shrubs and young trees; (iii) an almost continuous middle canopy of tree crowns at 20–25 meters above the forest floor; and (iv) an upper canopy of relatively isolated emergent trees (many of which may be deciduous) rising to 50–60 meters. Vines, strangling figs, and epiphytes are prominent. Typical tree species include *Streblus zeylanica*, *Hydnocarpus illicifolius*, *Shorea vulgaris*, *Dipterocarpus costatus*, *Diospyros* spp., and *Hopea recopei*. In mature condition, these forests have a maximum total biomass of about 350 tons/ha, with an annual net growth rate of about 2.5 percent.

In Thailand, tropical evergreen forests are best developed in the far southeast, but a drier sub-type is found along the Dangrek range to Khao Yai and in patches from Khao Yai north to Loei. In Lao PDR, these forests occur in the wet areas of the Annamites, with one of the last great reserves in continental southeast Asia found in the Se Kong Valley bordering the Bolovens Plateau. Patches of evergreen forest are found in places along the Mekong. In Cambodia, rainforest grows in the Southern Uplands in the south and along the Annamite Chain in the east, comprising about 30 percent of the forested area of the country. In Vietnam, evergreen forest is found in parts of the Mekong Plain, stretching into the Annamites; these forests have suffered greatly from defoliation and, in the hilly areas, from swidden agriculture.

Hill Evergreen Forest

Hill Evergreen Forest occurs above 1,000 meters (1,200 meters in the southern Annamites),

in areas where annual rainfall exceeds 2,000 millimeters and is fairly evenly distributed throughout the year. The canopy is typically lower than in tropical rainforests, often forming just 2 layers; mosses, ferns, and epiphytes are very common in these moist cool forests. The total biomass is much less than in tropical evergreen forests, amounting to no more than 230 tons/ha. Typical trees include *Schima*, *Engelhardia spicata*, *Michelia champaca*, *Quercus*, and *Castanopsis*.

Hill Evergreen Forest occurs in Thailand on the upper elevations of Khao Soi Dao, in the upper elevations of the Petchabun range, and in the upper parts of Chiengrai. It is widespread in the higher elevations of northern Laos (though much of it has been replaced by grasslands following soil exhaustion by maize and opium growing hill tribes). Cambodia has small sections of Hill Evergreen Forest in the Southern Uplands and the Annamite chain. In Vietnam this forest type is found only in the Annamite chain, where it has been greatly disturbed by swidden agriculture.

Coniferous Forest

Coniferous Forest is typically found in a few restricted areas on sandy soils at elevations of 600-1400 meters, though a few outcrops are found in the plains and along sea coasts. The conifers are often found in open stands with few other tree species (as at Phu Kradeung in Thailand's Petchabun Range), but in other places they are found in denser mixed stands (as at Nam Nao, also in the Petchabun Range). The undergrowth is usually grass, indicating the effect of fire. Southeast Asian coniferous forests consist mostly of *Pinus merkusii* and *P. khesya*. When found with other trees these are usually *Quercus*, *Castanopsis*, and *Irvingia malayana*. Conifers may mix with Dry Dipterocarp forests on dry lateritic soils.

In Thailand, Coniferous Forest is also found in Khon Kaen, and in small areas of Sisaket, Surin, and Ubon, as well as in a few areas of Chiengrai. In Laos, Coniferous Forest is reported to cover perhaps 300,000 ha (Turbang, 1973). Much of the Plain of Jars was covered in pine, although most pine forests have been destroyed by the recurrent battles that have raged across the plain. In Cambodia, Coniferous Forest covers only 18,000 ha, mostly on the Kirirom plateau with pockets in the Cardamom mountains and on the plains of Pursat and Kompong Thom. In Vietnam, pines cover about 120,000 ha in the upper Sre Pok watershed, but they have been seriously damaged by war; a small area of *Pinus kempfii* (*dalatensis*) occurs at about 2,000 meters in the Chen Yang Si mountains (Singh, 1972).

Mangrove Forest

Mangrove Forest occurs in the basin only in the Mekong Delta, covering 2,800 km². Mangroves occur in the zone between the neap tides and the spring tides, where the soil is formed of silt from the land and sand from the sea. The flora consists of relatively few species (seldom more than about 30-40), dominated by *Avicennia*, *Bruguiera*, *Rhizophora*, *Nipa*, and *Melaleuca* (in sequence from the sea to the land, reflecting salt-water tolerance).

Mangroves are considered to be very important ecological resources, as breeding grounds for crabs, shrimp, and many species of fish, but perhaps equally important is their role in claiming land from the seaward edge of the delta. The mangrove trees both retard erosion from the ocean (of vital significance in a typhoon area), and build up soil around their unique air-breathing root systems. As the forest advances toward the sea, the landward margin becomes progressively higher and drier and the forest tends to merge with the adjacent dry-land forest. This land eventually becomes suitable for rice cultivation and other agricultural uses. Unfortunately, the mangroves have been particularly susceptible to herbicides, with about half their area destroyed. Five years after defoliation, these mangroves are not yet regrowing, dramatically demonstrating that an intensive reafforestation scheme is required.

Fresh-Water Swamp Forest

Fresh-Water Swamp Forest is found mostly in Cambodia, along the edges of the Great Lake and along certain banks of the Mekong, totaling about 680,000 ha (in 1960). Dominated by *Homalium brevidans* and *Hydnocarpus anthelmintica*, this forest type is considered vital to the continued productivity of the Great Lake. The recent clearing of much of this forest has led to increasing siltation, thus threatening the very existence of the lake (which is less than 1 meter deep during the dry season). During the rainy season, when the lake quadruples in size, the swamp forest serves as the spawning ground for many species of fish, so the loss of this forest will also result in a reduction of the fish populations (Bardach, 1960).

Deciduous Monsoon Forest

Deciduous Monsoon Forest receives 1,250 to 2,000 mm of rain annually) and has well pronounced wet and dry seasons. It is generally confined to elevations below 1,000 meters. It is formed on well-drained lateritic soils, including Ferric Acrisols, Gleyic acrisols, and Ferralic Cambisols. In comparison with the evergreen forests, much more light reaches the forest floor and the undergrowth is much denser. The annual net growth is about 2 percent, with the total above ground biomass about 70-80 tons/ha. During the rainy season, movement through these forests is difficult and visibility is low, but during the dry season much of the undergrowth is burned off to be replaced by new grasses springing from the protected underground rhizomes. Most of the trees lose their leaves during the dry season, including the dominant species such as *Dipterocarpus alatus*, *Shorea obtusa*, and *Hopea odorata*. Of outstanding economic significance in some mixed deciduous forests is the presence of teak (*Tectona grandis*).

In Thailand, Deciduous Monsoon Forest occurs around Aranyaprathet, in part of the Dangrek range, along the Lao border in Ubon, throughout the hilly regions of Sakhon Nakhon, along the Petchabun range, and in much of Chiangrai. In the recent past, it was much more widespread throughout northeastern Thailand, but this forest type has been greatly reduced in favor of growing cash-crops such as cassava, sugar cane, and maize. In

Laos, this forest type is found in the south around Pakse and in much of the area from Vientiane to the Burmese border below 1,000 meters (these are good teak- producing forests). In Cambodia, it is widespread in the north, covering about 40 percent of the forested land of the country (including Dry Dipterocarp Forest). In Vietnam, it occurs in the southern-most part of the Annamite chain, where the hills are low and rolling.

Dry Dipterocarp Forest

Dry Dipterocarp Forest (also called “Savanna Forest”) replaces Deciduous Monsoon as the annual rainfall drops below 1,250 mm and the dry season lengthens to 6 months (arid months total 8.5 in some places in northeastern Thailand). Soils are typically dry and shallow, lacking humus. These forests are more open than the Deciduous Monsoon Forests with more grass and bamboo undergrowth; most are burned annually. With continued burning, large areas of Dry Dipterocarp Forest have become grasslands with only scattered trees and groves along the watercourses. Typical tree species include *Shorea obtusa*, *Dipterocarpus obtusifolius*, *D. tuberculatus*, and *Pentacme siamensis*. Annual net growth is about 1.5 percent, with biomass no more than 65 tons/ha and often considerably less.

In Thailand, this type of forest is found in much of the northeast, especially in Chaiyaphum and Nakhon Ratchasima. In Laos, Cambodia, and Vietnam, it occurs in the driest areas among the Deciduous Monsoon Forests. “Bamboo Forest” is often considered a separate forest type in the deciduous forests, but bamboo is a successional stage which is usually caused by fire. Allowed to mature, bamboo forest will usually develop into Deciduous Monsoon or Dry Dipterocarp Forest, though in wetter areas it may eventually develop into Tropical Evergreen Forest.

Savannas and Grasslands

Whitmore (1975) has shown that it is most probable that all tropical grasslands, with the obvious exception of those found on some more or less permanently swampy sites, are entirely derived from woodlands or forests as a result of long-continued burning by fires started by man or lightning and that there is no such thing as a tropical grassland climate. This thesis is strongly supported by Whyte (1972), who suggests that no less than 80 percent of the grasses now recorded for Malaya come from human introductions.

While the drier parts of monsoonal Asia may have been richer in native grass species, the clearing of forests in the Mekong basin has resulted in considerable immigration of grass species from outside the basin. The replacement of forest by Imperata grasslands is one obvious manifestation of this phenomenon. The relative paucity of natural grasslands (except in the seasonally submerged areas) suggests that the edible grasses, which provide the main food for grazing animals, will prosper only where appropriate growing conditions are maintained by people (often through the use of fire).

Forests as a Resource

The great diversity of forest types and of tree species within each forest type has played a vital role in human economies in the basin for thousands of years. They have provided food, fuel, construction materials, medicines, resins, gums, pesticides, and supporting wildlife. Modern urban populations remain dependent on the forests for construction timber, fuel, and paper, with wildlife and forest plants and products somewhat reduced in importance.

Consumption of wood in the basin is difficult to quantify, since many of the statistics seem unreliable. For example, FAO (1975) reports that per capita consumption of firewood and charcoal in Thailand is 0.37m³ per year, while Backer and Openshaw (1972), who carried out considerable fieldwork in Thailand, concluded that per capita consumption of fuel wood and charcoal was actually 1.23m³ per year (totaling 49 million m³ in 1970). Given the level of reliability of the available statistics, the values cited should be taken as indicators of the dimensions of the problem rather than precise figures. In general, it is believed that per capita consumption in the lower basin of roundwood, sawlogs, sawnwood, fuelwood, and charcoal averages from 1 to 2m³/year.

ILLUSTRATION 4: POTENTIAL FOREST PRODUCTION AND TOTAL REQUIREMENTS OF WOOD AND WOOD PRODUCTS (from FAO, 1976)					
Station	Operable forest area	Average growing station km ² (m ³ /ha)	Annual growth increment (m ³ /ha)	Total potential production (106m ³)	Total requirements (106m ³)
Thailand	10.0	41	0.82	8.2	40-80
Cambodia	9.0	77	1.54	13.8	7-14
Lao PDR	6.0	80	1.60	9.6	3-6
S. Vietnam *	2.9	80	1.60	4.6	19-38

* South Vietnam only (data not available for North Vietnam)

Illustration 4 presents information on the operable forest area, average growing stock, annual growth increment (calculated at 2 percent of the average growing stock), and total potential production for the riparian countries. The last column lists total requirements which can then be compared with total potential production.

It appears that potential production is in balance with requirements in Cambodia and production is well in excess of requirements in Lao PDR; but requirements are 5-10 times production in Thailand and 4 to 8 times production in Vietnam. Moreover, the projections for the future indicate an even more serious problem, since population is increasing and the area of operable forest is being reduced by 5 to 10 percent per year. Clearly, the forest resources of the basin area being over-utilized, and this is resulting in an essentially permanent depletion of the resource.

While wood is the most important forest product, many other products are of considerable significance, including bamboo (as food, construction material, pulp, and a wide variety of other local uses), rattan, mushrooms, many commercial species of orchids and other ornamentals, tannin for tanning hides, lacquer (*Rhus spp.*), resins (from *Hopea*, *Shorea*, *Dipterocarpus*, and other dipterocarps), lac (a resin excreted by a scale insect, *Laccifer lacca*, used to make shellac), and a wide range of antibiotics and other medicines.

Many plants which are currently cultivated in the basin have wild ancestors in the forests. When crossbred with domestic varieties, the new breeds may be more productive than currently cultivated breeds since they are more likely to have genetic resistance to the various indigenous insects and other pathogens which plague agricultural areas. Cultivated plants in the basin that have wild relatives in natural forests include durian, mango, banana, jackfruit, mangosteen, lingsat, rambutan, coconut, sugarcane, pomelo, pomegranate, taro, tamarind, bel-fruit, "sataw" (*Parkia* sp.), breadfruit, carambola, chempedak, hog plum, lesser yam, pigeon pea, cotton (*Gossypium* spp.), tea, cucumber, ginger, anise, cassia, cinnamon, betel-nut, black pepper, cardamom, turmeric, kenaf, and many others. These plants provide some of the basic raw materials with which plant breeders and geneticists can create new varieties suitable for local conditions.

Tropical plants are in constant competition with leaf-eating and boring insects, defending themselves with a wide range of chemical defense mechanisms, many of which are of direct importance to man. Rotenone, which kills ticks, fleas, lice, beetles, corn borers, aphids, mosquitoes, cockroaches, and house-flies without hazard to the human or domestic animal consumer and without residue tolerance problems, comes from several plants found in the basin, including *Derris elliptica* and *D. malaccensis*. Pyrethrum (from *Chrysanthemum*) is another important pesticide which has been widely used and grown in the basin, possibly for thousands of years. Although insect pests are likely to be increasingly controlled in the future by biological means (for example, use of hormonal substances applied to upset the life cycle of the target species) and by diversification of crops to avoid the vulnerability of monoculture, when a serious insect outbreak does occur, an insecticide is needed. Because most synthetic chemical insecticides have many deleterious side effects and their general application is not recommended, the botanical pesticides like rotenone and pyrethrum are in many situations much more suitable for agricultural use. Many additional botanical products found in basin forests are useful as synergists, emulsifiers, adhesives, and stabilizers in insecticides, and future research in the basin will undoubtedly reveal other useful botanical pesticides as well.

Besides the direct consumption of forest products, the forests have indirect resource values as well. Forests are particularly important as protection for watersheds, and this aspect of forests in the Mekong basin has been treated in some detail by Singh (1972). He pointed out that adequate forest cover can increase productivity of the land, mitigate floods and drought, increase rainfall, conserve soil, and protect reservoirs and irrigation systems from siltation (although few quantitative data are available to date on these points). With the rapid clearing of forests in recent times, flood hazards have increased; many tributary streambeds have been degraded and considerable erosion affects the geologically recent deposits on plains below foothills (Battambang, Vientiane plains).

Several other indirect values deserve mentioning. A period of forest fallow can return fertility to the soil under shifting cultivation, an essential ingredient of swidden farming (see Chapter 8). Forests also provide the best habitat for many forms of wildlife, ultimately supplying protein to people living in and near the forests. Finally, forests have considerable scenic value for tourism, forming the basis of a potentially outstanding system of protected areas (see Chapter 6).

In the long run, forests are capable of playing an even more important role in maintaining the basin civilizations. As the non-renewable resources of coal and oil decline, organic production will grow in production. Wood will remain valuable as a versatile raw material, and more species will have economic value. Forests can also be expected to become a major source of raw materials for the new biochemical technology which can be expected to develop to replace oil and coal as sources of complex molecules for food and manufacturing (Whitmore, 1975). Already, edible protein can be extracted from leaves, and lignin is being used to manufacture plastics, ion-exchange resins, vanillin, tanning agents, stabilizers for asphalt emulsions, as dispersants for oil-well drilling, and for ceramics processing. Cellulose can be utilized for rayon and plastics and as a raw material for hydrolysis to sugar. Complex organic molecules such as steroids are conveniently obtained from plant sources. While the technology exists to synthesize complex organic molecules from simple inorganic ones, this involves the expenditure of enormous quantities of energy which often still comes from non-renewable sources. It thus seems wise to utilize plants to perform the initial reduction of carbon dioxide by using solar energy and to restrict human intervention to manipulating plant products.

Conclusion

The basin forests are essential natural resource systems, with many direct applications to agricultural problems. More important, the forests are a renewable resource, which, with proper management, can give sustained yields for the indefinite future. Over-exploitation and lack of management have reduced the productivity of the basin forests, and growing stock per unit area and mean annual increment have both dropped to relatively very low levels. The forests of Southeast Asia are being exploited at the fastest rate in history, and at a rate far beyond the natural ability of the forests to replenish themselves. Much forested land is being turned into low-productivity grasslands which are permanent in timescales relevant to land-use planning. Productivity of the forests can now be raised only through intensive management of remaining natural forests and establishment of forest plantations to meet the future requirements for forest products in the basin.

Chapter 5

The Native Fauna

Introduction

Previous chapters have shown that the Lower Mekong Basin has a wide variety of geological formations, moisture regimes, soil types, and vegetation. These factors contribute to a very rich fauna, both on the land and in the water. Studies supported by the Mekong Secretariat (McNeely, 1975; Mekong Basin Fisheries Study, 1976) have reported at least 212 species of mammals, 696 species of birds, 213 species of reptiles and amphibians, and 800 species of fish (including some saltwater species which enter the basin seasonally). More complete information can be found in the reports mentioned above; here we will briefly describe the fauna and discuss its relevance to agriculture and food production.

Land Animals

The terrestrial mammals include 21 species of insectivores, 66 species of bats, 15 species of primates, 51 species of rodents, 34 species of carnivores, 16 species of grazing animals, one hare, one flying lemur, and 2 species of pangolins. Ten of these are endemic (occurring only within the Mekong basin) (Illustration 5) and 14 are considered rare and endangered by IUCN (Illustration 6).

The basin birds include 71 families and 305 genera, with over 500 species resident and another 160 as winter migrants. About 30 species are resident at low densities but are joined by migrants during the winter. Among the more prominent groups are hawks (35 species), pheasants (17 species), ducks (14 species), owls (18 species), woodpeckers (23 species), babblers (67 species), flycatchers (34 species), and warblers (54 species). Species endemic to the Mekong basin include the Chestnut-headed Partridge, Imperial Pheasant, Germain's Peacock-Pheasant, Red-vented Barbet, Bar-bellied Pitta, Yellow-billed Nuthatch, Short-tailed Scimitar Babbler, Sooty Babbler, Gray-faced Tit-Babbler, Collared Laughing Thrush, White-checked Laughing Thrush, and Gray-crowned Crocias.

Generally, faunal diversity closely follows floral diversity, so the most diverse wildlife is found in the evergreen forests and the least diversity is found in agricultural areas. Since wildlife is most common in the non-agricultural areas, it is broadly complementary to watershed protection. However, selective forestry and even some limited clear-cutting is seldom detrimental to wildlife as a whole and may even be beneficial, since the new secondary growth is often more productive for the larger grazing animals. Generally, the southern Uplands, Northern Highlands, and Annamite Chain have the most wildlife. The

ILLUSTRATION 5: MAMMALS ENDEMIC TO THE LOWER MEKONG BASIN	
Species	Distribution
Northern Smooth-Tailed Treeshrew (<i>Dendrogale murina</i>)	SE Thailand, Cambodia, southern Vietnam
Marshall's Horseshoe Bat (<i>Rhinolophus marshalli</i>)	SE Thailand
Bourret's Horseshoe Bat (<i>Rhinolophus paradoxolophus</i>)	NE Thailand, northern Vietnam
Lesser Slow Loris (<i>Nycticebus pygmaeus</i>)	Annamite Chain
Douc Langur (<i>Pygathrix nemaeus</i>)	Annamite Chain
Francois Langur (<i>Presbytis francoisi</i>)	N. Laos, N. Vietnam (Annamites)
Pileated Gibbon (<i>Hylobates pileatus</i>)	West of Mekong, S. of Pakse, SE Thailand
Lao marmoset Rat (<i>Hapalomys delacouri</i>)	Annamite Chain
Owston's Civet (<i>Chrotogale owstoni</i>)	Laos, N. Vietnam (Annamites)
Kouprey (<i>Bos sauveli</i>)	Southern Laos, N. Cambodia, small part of NE Thailand

ILLUSTRATION 6: IUCN ENDANGERED SPECIES OF MAMMALS IN THE MEKONG BASIN	
Species	Distribution
Douc Langur (<i>Pygathrix nemaeus</i>)	Laos-Vietnam
Pileated Gibbon (<i>Hylobates pileatus</i>)	Kampuchea, SE Thailand
Asiatic Wild Dog (<i>Cuon alpinus</i>)	Basinwide
Clouded Leopard (<i>Neofelis nebulosa</i>)	Basinwide
Tiger (<i>Panthera tigris</i>)	Basinwide
Leopard (<i>Panthera pardus</i>)	Basinwide
Asian Elephant (<i>Elephas maximus</i>)	Basinwide
Javan Rhinoceros (<i>Rhinoceros sondaicus</i>)	Basinwide
Sumatran Rhinoceros (<i>Dicerorhinus sumatrensis</i>)	Basinwide
Brow-antlered Deer (<i>Cervus eldi</i>)	Basinwide
Wild Water Buffalo (<i>Bubalus bubalis</i>)	Cambodia, S. Laos
Gaur (<i>bos gaurus</i>)	Basinwide
Banteng (<i>Bos javanicus</i>)	Basinwide except N. Laos
Kouprey (<i>Bos sauveli</i>)	S. Laos, N. Cambodia, NE Thailand

Note: "Basinwide" species are found in small numbers in widely scattered localities in all riparian countries; however, the total population of the two species of rhinoceros totals no more than 10-15, and some of the listed species may no longer occur in the basin.

Annamites are particularly important as a centre of endemism; of 11 endemic birds, 9 occur only in the Annamites, and of 10 endemic mammals, 5 are found only in the Annamites. In the Mekong Plain and the Korat Plateau, by contrast, the wildlife is mostly confined to the uplands and thinly populated areas such as northern and eastern Cambodia.

Wildlife as a Resource

Wildlife contributes to the well-being of basin residents in many ways, including direct consumption as food, sale of live animals, skins, horns, and ivory, production of phosphate fertilizer (bat guano), controlling insect and rodent populations, and as a genetic reservoir. Virtually all of the 4 to 5 million indigenous people living in the basin are upland farmers who obtain much of their protein from wild game. In the Korat Plateau and much of northern Cambodia and southern Lao PDR, farming is possible only during a limited period of the rainy season. During the dry season, the villagers become hunters and gatherers like their ancient ancestors, collecting forest fruits, vegetables, and mushrooms, and hunting birds, reptiles, amphibians, small mammals, and (in the remote uplands) large mammals.

The existing and proposed Mekong river water resource development projects serve to greatly increase the surface area of water, offering great potential for improving habitat conditions for aquatic birds, including ducks (14 species), grebes, pelicans, cormorants (4 species), herons and egrets (16 species), finfoots, rails, gulls and terns (12 species), jacanas, wading birds (about 37 species), and many others. Provided management measures could be enforced, these birds could provide the basis for greatly increased production of protein in the tributary and mainstream reservoirs. The even-toed ungulates, including pigs, deer, and wild cattle, include most of the game species which have potential for controlled hunting, harvesting, and captive breeding. In optimal habitats, the biomass of such game animals may average as much as 4,000 kg/km², with as much as 10,000 kg/km² in certain localized habitats (especially alluvial grasslands). In competition with domestic animals, these densities will invariably be lower; however, it should be emphasized that the wild ungulates often feed on different plants or different vegetative stages than do domestic animals and they are seldom dietary competitors to the extent where one excludes the other (except where stocking density of domestic forms is extremely high). Thus a combination of wild and domestic ungulates can, in many situations, be more productive than either one alone.

Deer especially have a high potential for captive breeding or semi-domestication. They utilize habitat in a manner which is often complementary to that of cattle, and they produce high quality meat (worth 30-60 percent more on the local market than that of domestic cattle). Some species are able to live in close proximity with man; they are hardy and resistant to many cattle diseases; and the male annually produces antler velvet of a value equal to 2-3 times its own market value as meat (Tan, 1968). Tan estimates that in Vietnam, one buck can earn for the farmer an income double or triple that of a bovid, with the same expense.

The four species of wild cattle in the basin are closely related to the domestic forms and

may compete with them for fodder where large numbers of domestic animals are grazed in the forests. In such situations, disease transmission is a real danger, especially from the domestic form to the wild form. Conversely, the wild bovids have much to contribute to future cattle breeds if they can maintain their genetic distinctiveness. Cross-breeds between wild tropical cattle and domestic breeds from Europe, the United States, or Australia might be more productive in tropical conditions.

The gaur, banteng, and kouprey (the last large mammal to be discovered by science) are in the same genus *Bos* with domestic cattle. All three species have a long and intimate relationship with fire, which burns old grass and brings up tender new shoots, so slash-and-burn agriculture actually has a beneficial effect for these species, provided the density of human population is not so high as to make hunting a limiting factor. The kouprey may have contributed to the ancestry of the Indian zebu, the most successful tropical cattle breed. It has been suggested that kouprey may be resistant to rinderpest and other cattle diseases, indicating the possible importance of this species for breeding.

The wild water buffalo is the same species as the domesticated form, though some domestic breeds have a different number of chromosomes than the wild form (again showing their potential importance for breeding). The buffalo is probably the most important domestic animal in the basin, providing power for pulling plows in rice fields, threshing rice, pulling carts and logs, and providing fertilizer. Domestication of the buffalo allowed the development of rice agriculture in the lowlands, and the maintenance of a viable wild stock should contribute to the continuing success of the breed.

From many points of view, the elephant is the most important large wild mammal in the basin. Tamed as a working animal, the elephant is vital in developing valuable teak and other forests in rough hilly areas (1970 forest production in Thailand was approximately US\$100 million). As a ceremonial animal, the elephant plays a central role throughout the basin; as a wild animal the elephant is an attraction to visitors to Khao Yai National Park and potentially in other areas, such as the Petchabun range and the Elephant mountains.

As transportation for tourists, the elephant has proved its usefulness in such areas as Kaziranga in Assam and Chitwan in Nepal and has similar potential in the Lower Mekong Basin. As items of foreign trade, 245 elephants were exported from Thailand from 1967 to 1971, earning about \$250,000. As tools in wildlife research, trained elephants have been shown to be the most practical means of making a census of Asian wildlife. As beasts of burden, elephants have long been vital to transport systems and logging throughout the basin.

At present, forestry elephants generally work with trucks, with the elephant bringing cut logs downhill to the roadhead where the truck carries them to the sawmill. In former times, the rivers were used as transport routes and in many places, the reservoirs created by dams existing and envisioned can in the future similarly be used as transport routes. Many basin forests can best be worked by elephants, since high-quality trees are not concentrated enough to make roads economically feasible, especially given the reality of rising petroleum costs and the increased siltation likely to be caused by construction of logging roads and by modern clear-cutting techniques.

Wildlife and Pests

The role of wildlife in controlling pest outbreaks is seldom appreciated and precise data are generally scarce. A brood of young woodpeckers, for example, may consume 8,000-12,000 insect pupae per day. When tree trunks are attacked by wood-boring insects, they are regularly worked over by woodpeckers, completely debarking diseased tree trunks and helping to control further spread of disease. Frogs and toads generally breed at times when insects are at their peak populations and since most amphibians feed on insects, they exert a strong influence on insect populations (as well as supplying protein to local people).

The basin has about 57 species of insect-eating bats, some of which have populations numbering in the millions. Indications are that many bats consume over half their own weight in insects each night, with different species of bats preying on different groups of insects, so that a rich fauna of bats ensures that a wide spectrum of insect species is preyed upon. Widespread application of chemical pesticides is likely to adversely affect the numbers of insect-eating birds, amphibians, and bats and therefore the stability of insect populations, leading to periodic pestilence. The replacement of the insect eating fauna by chemical pesticides thus does not seem to be a viable long-term alternative. Instead, the value of these forms to agricultural development should be realized and efforts made to conserve them. Chemical pesticides should be used at the minimum rate required for each specific situation.

Another pest problem is rats, which are estimated to consume 10-20 percent of all grain growing in Thailand. Losses of up to 100 percent of the rainy season rice crop have been reported from the central plains of Thailand, and damage to the dry season crop is intensified in all areas (Shuyler and Ratanaworabhan, 1970). Recent reports from northern Thailand suggest that in some upland areas (where natural predators have been exterminated), up to 60 percent of the crop is destroyed by rats. The damage to crops is done by a relatively few species which are commensal with people and replace the less destructive forest rats when forests are cleared. The commensal rats benefit from human effects on the environment, breeding rapidly and sheltering under houses, barns, and other shelters or digging burrows in ricefield bunds or termite mounds.

Many of these rodents are hosts for disease vectors, including mites (scrub typhus), fleas (bubonic plague), and ticks (several diseases). In addition, their droppings in stored grain often carry such diseases as leptospirosis and salmonellosis. Efforts to control rodents through poisoning or trapping have failed; for example, at the beginning of the 1969 rainy season, a week-long poisoning campaign in one province in Thailand killed 680,000 rats (6 rats/ha) but this did not significantly reduce damage to the next crop (Shuyler and Ratanaworabhan, 1970). The best approach to rodent control is in improved village sanitation and in storing rice and other foods in rodent-resistant containers, and in maintaining natural predators of rodents such as owls, hawks, snakes, mongooses, and civets.

Wildlife Breeding and Genetic Resources

As mentioned earlier, domesticated buffalo and cattle owe a genetic debt to wild ancestors currently living in the basin. Several other species are important genetic resources as well, furnishing the raw material for developing new breeds which may have characteristics leading to greater productivity, such as disease resistance and ability to prosper on available natural vegetation. Among the native species of importance to agriculture and food production are the red jungle fowl (*Gallus gallus*), the ancestor of the domestic chicken; the various ducks (*Anatidae*), at least some of which ancestral to domestic ducks; the wild pig (*Sus scrofa*), the ancestor of the domesticated pig (many in remote areas are cross-breeds); jungle cat (*Felis chaus*), which may have played a role in the development of the domestic cat (an important pest controller); and the jackal (*Canis aureus*), a possible contributor to the ancestry of the domestic dog. Further, the domesticated elephants are all recently descended from the wild form and probably should not be considered "domestic" since they are identical to the wild form; many tame elephants are bred with wild elephants in the forest.

Live animals from the basin are in high demand from zoos and laboratories throughout the world, with the animal trade earning about \$4-5 million in Thailand in 1970. Douc langurs are worth \$2,500 each, Sumatran rhinoceros about \$200,000 per pair (if they could be obtained), elephants \$1,500 each, gibbons \$1,000 each, and clouded leopards \$2,500 each. These figures are given as indicators of the potential magnitude of the animal trade, which could yield a sustainable economic return for the foreseeable future if properly managed. Unfortunately, the live animal resource has been grossly abused, to the extent that there are now appropriate bans on the trade of many species (including all of those listed above). The future of the animal trade is likely to depend increasingly on captive breeding, with the few remaining forests serving as national parks or genetic reserves to be used sparingly or only for non-consumptive uses.

Virtually all species of wildlife in the basin can be bred in captivity (exceptions are flying lemurs and pangolins), but those likely to be particularly productive include hog deer, sambar, Eld's deer, banteng, gaur, kouprey, elephants, gibbons, monkeys, pheasants, quail (the quail trade in Hong Kong earns \$5.7 million per year), wild ducks, and crocodiles (the crocodile trade in Thailand earns \$400,000 per year).

The captive bred wildlife can be used not only for export or local use, but also for releasing in natural habitats when protective measures can be enforced. Many forested areas which are prime wildlife habitat are being cleared for extractive agriculture, though the soil is too poor to support a crop for more than a few years. Such land will soon revert to grassland and forest, but the wildlife has usually already been exterminated. Natural repopulation of these areas might take decades, but a boost from populations bred in captivity can significantly increase the rate of repopulation, thus making the land again productive. Captive breeding of species of wildlife in demand for zoos and laboratories could also help reduce pressure on the national parks and other protected areas while also providing jobs for villagers and foreign exchange to the economy. However, it is vital that captive breeding should be seen not as a replacement for habitat conservation but rather as a necessary

supplement in a period of rapid exploitation. The major thrust of any conservation program must be in protection of fairly extensive areas of natural habitats (see Chapter 6).

The Aquatic Animals

Importance of Basin Fisheries

Protein production through aquatic animals is highly efficient in the tropics, since the cold-blooded animals grow quickly in the nutrient rich warm waters of streams, rivers, swamps, and ponds, rapidly and efficiently transforming their food supply (phyto and zoo-plankton, algae, molluscs, aquatic plants, detritus) into high-quality protein available for human consumption. It is thus no surprise that the freshwater animals, including fish, shrimps, crabs, shellfish, and frogs, are currently much more important than the wild land animals to basin economies. Virtually all farmers living in the lowlands and near streams engage in subsistence fishing as part of their daily activities and depend on fish for their primary source of protein. The extent of this subsistence fishing is estimated to be at least 100,000 tons/year.

The aquatic fauna of the basin is very rich, including over 800 species of fish, 5 species of dolphins (only one of which has been known to reach above Khone Falls), 2 species of crocodiles, 49 species of frogs and toads, 12 species of turtles, and numerous species of crabs and other crustaceans. The fish and crustaceans are by far the most important economically and nutritionally and have the greatest potential for development, especially through aquaculture.

In view of the importance of the fishery resource, a comprehensive evaluation of the fisheries in the lower basin was carried out, under Mekong Committee sponsorship, over the period 1974 to 1977. This study, termed the Mekong Basinwide Fishery Survey (MBFS), had the objective of determining the production, consumption, natural history, development potential, and impact of water resources development on the basin fisheries. Though the fieldwork was curtailed, a great deal of original data was collected, in many cases the first such data ever available.

Much remains to be learned; for example, complete life history data is still not available for any species, production and consumption figures are still gross estimates, and so on. Therefore final considerations of fishery impacts and future directions of changes in the fisheries as a result of water resources development, although relatively complete, should be considered tentative. Nonetheless, the data obtained were sufficient for qualitative generalizations about the potential impacts on fisheries of the basin development. These indicate that substantial management, particularly in the area of aquaculture, will be needed to overcome a projected loss of total fish stock and catch in the basin.

The MBFS found that most fish species in the Mekong, as in most tropical rivers, are well adapted to a widely fluctuating environment, with a wide range of tolerance for temperature,

pH, dissolved oxygen, and other environmental parameters. Adaptations to survive frequent and abrupt environmental changes have evolved in some of these species, such as accessory breathing organs found in some catfish, climbing perch, and murels that enable the fish to utilize atmospheric oxygen, so the fish can survive periods of deoxygenation of water as usually occurs in the peripheral zones of inundated areas during flood seasons. Another interesting adaptation is the ability to move over damp land, in some species, which enhances survival when habitats dry out.

Importance of the Annual Inundation Cycle

The life cycle of many fish species in the Mekong is adapted to the annual inundation cycle. With the onset of floods and the submergence of ricefields, ditches, swamps, oxbows, and other low-lying areas over a considerable distance on either side of the riverbank, large numbers of the fish population migrate from the rivers to the inundated areas, mostly for spawning. While the breeding adults are quite hardy, large numbers of eggs and hatchlings die due to desiccation if the flood recedes quickly; if, on the other hand, the floodwaters persist for a relatively long period, large scale mortality of fry and fingerlings occurs due to deoxygenation and drastic changes in pH resulting from decay of organic matter in flood waters. This emphasizes the importance of the very high rates of reproduction invariably found in basin fish.

An estimated 85 to 95 percent of the total freshwater fish population in the basin follows this inundation spawning pattern. However, even those that do not follow the inundation pattern could be adversely affected by the environmental changes following changes in the inundation pattern.

Trend of Basin Fisheries Production

Illustration 7 summarizes current production and consumption of fish in the basin and projects the demand for the year 2000. This shows that fish supply must double if per capita consumption remains the same and must quadruple if recommended minimum dietary requirements are to be met. In this respect it is important to estimate the probable trend in the production of the natural freshwater fisheries of the basin, to evaluate the extent to which this supply can continue to meet the local needs. However, the basic data for the fisheries of the basin are sparse and often misleading, so the trend of fisheries production in the area is somewhat speculative.

One of the most important fishery resources of the basin is the Great Lake of Cambodia, sometimes called the world's greatest freshwater fishery. As Henri Mouhot, the French explorer and naturalist, said as early as 1860, "The Great Lake is a source of wealth to the whole nation; the fish in it are so incredibly abundant that the play of the oars is frequently impeded by them". However, production has fallen from nearly 80,000 tons in 1940 to only 25,000 tons in 1965, and the decline is continuing (Bardach, 1966). This decline in Great Lake fisheries, whatever the reasons, has effects throughout the Mekong Plain. The

MBFS has shown that the aquatic production of the Tonle Sap-Great Lake Mekong floodwater complex above Phnom Penh is a critical factor in sustaining the entire lowland delta fishery.

Much of the exploitation in the Great Lake and its tributaries results from the netting of fish during the gradual recession of the fish-laden floodwaters. Fish not captured during their migrations to and from the vast inundation areas support the fisheries of the mainstream channels. During the dry season, almost all of the fish that have survived the extreme but essentially unmanaged fishing pressures of the declining flood period and commencing low water stage have moved into the major river channels. With the beginning of the wet season these fishes migrate again both upstream and laterally to their spawning grounds.

ILLUSTRATION 7: PRODUCTION, CONSUMPTION, AND FUTURE DEMAND OF FISH (METRIC TONS)			
Country	1970		2000
	Production	Consumption	Projected Demand ²
Cambodia	145,000	180,500	580,000
Lao PDR	30,000	30,000	100,000
NE Thailand	80,000	143,700	790,000
V.N. Delta	<u>230,000</u>	<u>137,000</u>	<u>480,000</u>
Totals	485,000¹	491,200	1,950,000

Source: Netherlands Economic Institute; MBFS.

¹ Includes approximately 40,000 tons from aquaculture.

² These estimates assume an increase in per capita consumption of fish. If per capita consumption remains the same as in 1970 (that is, if the increase in animal protein needs are met by livestock and poultry or if protein deficiencies still remain), then the demand for fish in year 2000 will be about 1 million tons/year.

During these migrations, the fish are also subjected to intense fishing pressures. Those that succeed in reaching the principal spawning grounds in the flooded areas produce their young there. The adults soon make their way back to the main rivers, and enroute are subjected once again to heavy fishing pressure. Meanwhile, the young grow vigorously over the inundated land, but as the water begins to recede they also follow it toward the main river channels, with only a few remaining in the residual lakes, ponds, and swamps. Enroute to the dry season riverine habitats, the young are significantly reduced in numbers by a fishery perhaps unparalleled in the world for its extreme intensity and for the ingenious variety of gear involved. Much of the population of the young is caught before reaching the haven of the dry season habitat. The survivors support much of the productivity of the freshwater deltaic areas, but also must provide the future spawning population which will move back into the inundation zones during the soon-to-come rising water period.

Because of the difficulties involved in trying to study the fishery of the Great Lake and its associated waterways during wartime, it has not been possible to establish the reasons for the decline in the lake fishery. However, population growth in the lowland areas has increased demands which in turn has stimulated the generation of the world's greatest concentration of aquaculture in parts of the delta downstream from Phnom Penh. The MBFS found that in a small farmer-fisherman hamlet halfway between Can Tho and Long Xuyen, each of the 20 households owned a fishpond which annually yielded about 37 tons of fish per hectare of surface water area; they had started the ponds seven years previously, when the fish catch in the river had begun to decrease. The total production of cage culture in Vietnam could amount to 40,000 tons annually, or 8 percent of the total fish production of the Lower Mekong Basin.

The increases in population have definitely resulted in adverse effects on several of the larger fish species of the Mekong. For example, the famous “pla buk” or giant catfish (*Pangasionadon gigas*), the largest fish in the Mekong, was once quite common in the lower stretches of the river, with several thousand a year being caught below Khone Falls (Pavie, 1904). They are now essentially extinct below Pak Sae (Davidson, 1974) and are rare and declining elsewhere.

While the information noted above may indicate a trend toward over-exploitation of the overall fisheries in the lowland areas of the Mekong plain, definitive conclusions are hardly possible because of lack of basic data in this zone. Also, in northeast Thailand the available data indicate increases in overall fish catch but no overall over-exploitation or depletion. Increasing use of irrigated agriculture will mean increasing use of agricultural chemicals including toxic chemicals, which would greatly depreciate the paddy fisheries which have traditionally contributed to the farmer's diet. For these various reasons, when estimating future fish production from natural freshwater sources, it is probably conservative to assume that future production rates will not exceed present levels. Hence the additional fish food needed to support new populations will have to come from other sources, including new reservoir fisheries, aquaculture, or marine fisheries.

Role of Marine Fisheries

Exploitation of marine fisheries has greatly increased in Thailand since 1960, due to market stimulation from Bangkok and other urban areas and to availability of new technology, and national policies will no doubt press for continuing expansion of this industry. However, most of this product is consumed by the urban sector, and is generally too expensive for meeting the needs of the rural population. Marine fisheries are also an important industry in Vietnam and seem destined for considerable expansion. Again, however, it seems likely that the protein needs of the rural population will be met mostly by freshwater fish sources.

Conclusion

The terrestrial wildlife of the Lower Mekong Basin is a natural resource of historical and current importance and offers considerable potential for the future. However, it is an inescapable fact that this wildlife resource is greatly suffering from over-exploitation, and most of the basin can be accurately described as defaunated, closely following the rate of deforestation. If nothing is done, the wildlife will continue its long slide downhill, to the ultimate detriment of the human population. By considering wildlife as an agricultural resource which must be managed as an integral part of the agricultural development plan, however, wildlife could continue to contribute to human well-being for the foreseeable future.

The situation regarding exploitation of the natural freshwater fisheries of the lower basin is much more difficult to quantify. The increasing demand for freshwater fish will likely be met by a combination of aquaculture and reservoir fisheries (including enhancement of reservoir fish yields to capitalize on all available feeding niches).

The orientation of the agricultural development model advocated in this report is not limited to the traditional concept of optimizing farm production alone, but to develop a truly comprehensive model which will yield at any future point of time sustained productivity of the overall complex. Perhaps the term “environmental agriculture” best indicates the scope and objectives of this type of planning. Thus the model will help to conserve wildlife and to supply an increased amount of fish. The integration of such a program into the overall basin agricultural development model presents a unique opportunity to attain this goal, and when combined with wildlife and forest conservation measures, it can provide the means of the conservation of these natural resources for human benefit.

Chapter 6

Resource Conservation and Water Resources Development

Introduction

Chapters 4 and 5 showed that the forests, wildlife, and fisheries of the lower Mekong basin are being exploited at a rate far exceeding the sustainable yield level, primarily due to improved access to previously remote areas, increasing human population, the replacing of a subsistence economy by a market economy, improvement of the technological means of forest clearing, fishing, and hunting, and inadequate enforcement of regulations in the rural areas. The situation in Thailand, at least, is rapidly approaching a terminal state. It is anticipated that within another decade Thailand's forests will be depleted to the extent that the country will become dependent on timber imports from the surrounding countries. The depreciation of forests will also mean a radical reduction in wildlife and many species adapted to mature evergreen forest may be lost, along with most of the larger game species. The situation in parts of Lao PDR, Cambodia, and the Vietnamese highlands is much better, and these countries have the potential for developing their forest resources on a rational sustained-yield basis.

Role of Protected Areas in Sustained-Yield Agriculture

Chapter 7 will show that traditional and extractive methods of agriculture are reaching a critical stage and will not be able to contribute very much to further agricultural development. Expansion of total land under crops in the basin has reached its productive limits, and modernization of basin agriculture will depend on more intensive use of the suitable agricultural land. With agricultural development stabilized on the more productive land, the marginal areas now being cleared of forest for transitory, extractive forms of agriculture can revert to its most productive long-term uses: forestry, watershed protection, recreation, and wildlife conservation.

The best way of controlling over-exploitation, and assuring that at least representative areas of forest remain to regenerate the abandoned areas created by extractive agriculture, is through a system of protected areas. Recognizing that the only conservation technique which even approaches success in the short term is total protection from modern land development, the Mekong Committee has sponsored a study which has designed a system

of protected areas for the basin, totalling about 7 per cent of the land area. These reserves are seen as integral parts of the land-use plan for the basin. The proposed and existing national parks and other protected areas are briefly summarized in illustration 8, including location, vegetation, and purposes (further details are found in McNeely 1975).

In the long term, the protected areas will be most effective if they actually form “core areas” within a larger land and resource management unit. Within the core area, no extractive activities are allowed, but in the surrounding “buffer zones” the forests and wildlife would be managed for the direct benefit of the local people, through controlled access for gathering of forest products, limited subsistence hunting in some specified zones, selective logging, sustained-yield swidden agriculture, and many other economically productive activities.

The unplanned extractive exploitation of the basin forests is disturbing watersheds (leading to recurrent cycles of floods and droughts), changing weather patterns, reducing fertility of the land, depleting forest reserves, reducing wildlife populations, and generally reducing the value of the land. In contrast, forested upland areas zoned for protected areas can reduce floods and siltation rates, increase usable water yields, improve dry season water supplies, and decrease water waste. All of these will help contribute to the longevity and productivity of reservoirs and irrigation systems in the lowlands, thus contributing to increased agricultural productivity.

Another important function of protected areas for agriculture is to maintain diversity in the basin. The various hill tribes often cultivate a very wide variety of crops and collect an even wider variety of plant products from the forests and abandoned swiddens. In contrast, modern agriculture tends to concentrate on relatively few crops which give a high economic return. As basin conditions change in the future, the upland forested reserves can provide the source material (of both plants and techniques) for solving as yet unforeseen agricultural problems. On a more immediate level, the reserves also contain the wild relatives of domesticated plants and animals discussed in Chapters 4 and 5. The wild forms evolve in competition with local insects, diseases, and other pathogens and may provide good breeding stock for creating new insect or disease resistant domestic or cultivated forms. Also, research is showing that the best way to control insects and other pests is often through using a natural enemy of the pest; such biological controls may often be found in the national reserves.

The protected areas thus can play an important role in the overall land use plan for the basin, of which water resources development is the central feature. It has been argued that major dams and reservoirs are very destructive of wildlife and forests, but it is clear that if current trends continue *without water resources development*, most wildlife and forests is doomed to disappear (as is already the case in many areas). *Water resources development*, on the other hand, offers greatly improved possibilities for conservation: more intensive agriculture in the lowland irrigated areas can reduce pressure on the agriculturally marginal uplands; the power produced by dams can help reduce the reliance on wood as a cooking fuel; by locating protected areas in and around reservoir watersheds, and by planning and financing them as integral parts of the reservoir projects, opportunities for extending control to the protected areas are optimal; and the overall economic

ILLUSTRATION 8: PROPOSED AND EXISTING PROTECTED AREAS OF THE LOWER MEKONG BASIN

Name	Location	Vegetation	Values to be protected
1 Elephant Mountains	Kamphot, Kompong Speu, Koh Kong, C.	Pines, wet evergreen, deciduous	Pileated Gibbon; Northern Smooth-tailed Treeshrews; Chestnut-headed Partridge
2 Cardamon Mountains	Koh Kong, Pursat, C	Wet evergreen, deciduous	Elephants, rhinos, gibbons, treeshrews
3 Khao Soi Dao - Khao	Chantaburi, TH	Wet evergreen, hill evergreen	Elephants, rhinos, Marshall's Horseshoe Bat
4 Delta Mangrove	Bac Lieu, An Xuyen, VN	Mangroves	Salt-water crocodiles
5 Great Lake	Pursat, Battambang, C	Swamp forest	Natural fisheries; waterbirds (perhaps flamingoes)
6 Angkor Wat	Siem Reap, C	Deciduous	Tiger, Eld's deer, Banteng
7 Köulen	Preah Vihear, C	Deciduous, man-made savanna	Kouprey (only place?); Giant Ibis, Siamese Crocodile
8 Kratie	Kratie, C	Deciduous, man-made savanna	Wild cattle
9 Lompat	Ratanak Kiri, C	Deciduous, open savanna	Wildlife in vicinity of Sre Pok, So San, Stung Treng dams
10 Langbian Plateau	Tuyen Duc, Darlac, VN	Wet evergreen	Langbian Rat, Collared Laughing Thrush, Gray-crowned Crocias
11 Senmonorom	Monduliri, C	Wet evergreen	Pygmy Slow Loris, Red-vented Barbet, Germain's Peacock Pheasant
12 Hodrai-Sou	Ratanak Kiri, C; Kontum, VN; Attopeu, C	Wet evergreen	Pygmy Slow Loris, Langbian Rat, Lao Marmoset Rat
13 Bolovens Plateau	Attopeu, Saravana, L	Wet evergreen, deciduous	Langbian Rat, Douc Langur, Lao Marmoset Rat, Red-vented Barbet
14 Tehepone	Savannakhet, Khammouane, C	Wet evergreen, deciduous	Imperial Pheasant, Yellow-billed Nuthatch
15 Khammouane Plateau	Khammouane, C	Wet evergreen, deciduous	Imperial Pheasant, Sooty Babbler, Douc Langur, others
16 Xieng Khouang	Borikhan, Xieng Khouang, C	Wet evergreen, deciduous	Owston's Civet, François' Monkey, Douc Langur, Yellow-billed Nuthatch
17 Luang Prabang	Sayaburi, C; Nan, TH	Deciduous, hill evergreen	Hunting and gathering tribe, the Yumbri
18 Nam Ou	Luang Prabang, C	Mixed	Wildlife of north Laos
19 Phong Saly	Phong Saly, C	Mixed	Southern China fauna
20 Upper Mekong	Houa Khong, C	Mixed	Burma/Lao/Southern China fauna
21 Khao Yai	Nakhon Nayok, Nakhon Ratchasima, Prachinburi, TH	Evergreen, grassland, deciduous	Elephants, tiger, gaur, Pileated Gibbon
22 Phu Kadoung	Loei, TH	Pine, grassland, deciduous, evergreen	Sandstone massif with numerous waterfalls; tourism, plentiful bird life
23 Nam Nao	Khon Kaen, Loei, Petchabun, TH	Diverse	Elephant, gaur, tiger, gibbons, plentiful birds
24 Phu Kheo	Chaiyaphum, TH	Grassland, evergreen, deciduous	Elephant, Sumatran Rhino, Bourret's Horseshoe Bat
25 Phu Luang	Loei, TH	Evergreen, deciduous	Elephants, gaur, tiger
26 Dangrek Range	Buriram, Surin, Sisakhet, TH; Oddar, Preah Vihear, C	Evergreen, deciduous, grassland	Many archaeological sites; elephants, Pileated Gibbon, Eld's Deer, Kouprey
27 Middle Mekong	Ubon Ratchathani, TH	Flood plain	Flood plain fauna
28 Phu Wua	Nong Khai, TH	Evergreen, deciduous	Hog Deer
29 Phu Phan	Kalasin, Sakhon Nakhon, TH	Evergreen, deciduous	Isolated massif with isolated fauna

development which will follow water resources development should stimulate sufficient government control and financing for enforcement of land-use regulations.

Conservation in Upland Areas Outside Protected Areas

Well over 60 per cent of the land area in the basin is not amenable to modern agricultural development, but much of this land will still be used for agriculture. It is difficult to predict in which direction upland farming systems will develop. Whether the trend would be towards forest farming, integrated livestock farming, or rotational systems of field crops will depend on future commodity prices, transportation systems, and government policies. These variables cannot be charted at this stage, but it is expected that with full implementation of the planned irrigation projects, beneficial changes will ensue for all agriculture in the basin. These changes will include improved markets, transportation and communication, availability of electrical energy, storage and crop protection methods, agricultural extension, and improved varieties of crops. With these very basic changes, both upland and lowland farmers can be expected to greatly improve their efficiency; for example, "mini projects" are planned for some of the best upland areas. But even with all of these changes, it is expected that some areas outside the protected areas will still be most suitable for low population density traditional agriculture and other areas will be most suitable for forestry.

The system of protected areas, wherein natural high-diversity forests would be conserved through protection, should be accompanied by a maximal effort in reforestation. Demand for wood will continue to increase at a rate which may exceed the rate of population increase, and current consumption trends indicate that future requirements of forest products will not be able to come from the natural forests. The best alternative during the reforestation process seems to be forest plantations both for firewood and for construction timber (including teak). Several species of very fast-growing softwoods (such as *Acacia catechu*) would be suitable for charcoal, particularly if they are formed into briquets; fast growing species of *Casuarina* can be used for poles; pine for construction and fuel can be grown as part of a forest farming system as described in Chapter 7; bamboo and kenaf can be used for paper pulp thus reducing pressure on the forests; and mixed species plantations of hardwoods can be grown for the more exacting furniture and construction requirements. Above all, the areas allocated to conservation must be protected from further depredations and the remaining production forests must be used at a sustained yield level.

Similarly, wildlife conservation in the protected areas should be accompanied by livestock development programmes in the agricultural areas. Improved production of pigs, cattle, buffalo, chickens, and ducks could play a significant role in improving the level of protein consumption in the basin and reducing the reliance of villagers on wild protein.

As mentioned earlier, captive breeding of species of wildlife in demand for zoos and laboratories could also be quite productive (though never as a substitute for habitat conservation). Areas being reforested with mixed species of trees also provide a good opportunity for reintroducing wildlife which has been bred in captivity. Such a boost from wildlife managers would be a vital step in helping exhausted, deforested land recover to previous levels of production.

Aquaculture

Chapter 5 showed that the natural freshwater fisheries of the basin, which have traditionally been the primary source of animal protein for farmers, will need to be supplemented by aquaculture and reservoir fisheries in order to meet the fish food needs of continuing population growth. While reservoir fisheries can meet part of this demand, major dependence must also be placed on aquaculture utilizing the year-round water supply made possible by water resource development.

While individual farm family ponds have been commonly used in irrigation areas, drawing water from the irrigation system, the main new opportunities envisioned in the Mekong programme are in commercial scale fish farms served by irrigation systems, using cage culturing of fish in lakes, rivers, and other waterways.

Commercial Aquaculture in Irrigation Service Areas

Land considered unsuitable for agricultural purposes because of its topographic position has been assessed for aquaculture development. For example, the Mekong Committee developed preliminary plans and designs for aquaculture farms utilizing a range of production rates depending upon investment levels. Usually these farms would be complexes of about 50 ha each, comprising a network of 2 ha farms suited to family size operation plus a small area devoted to fish stocking and hatchery purposes.

Included in the programme are plans for developing more efficient methods of fish feeding, such as manufacture of pelletized food in fish feed mills, and of storage, processing, and sale of the harvested fish. The current programme of the Mekong Committee includes provisions for constructing and operating demonstration fish farms of this type both in the Lao PDR and in Thailand. Such fish farming must pay careful attention to maintaining good water quality, i.e., to exclude harmful agricultural chemicals from the system. One of the objectives of the demonstration farms will be to incorporate these safeguards in the design of pond operations.

Delta Rehabilitation Potentials

Because of the possibility that development of the Mekong might adversely affect shrimp and other shellfish production in the Mekong delta, and also in order to enhance the delta fishery in general, the Mekong programme has considered three promising potentials for optimizing the benefits of delta development between cropping and aquaculture, taking into account the productive potentials of specific areas and the relative economics of alternative ways of using them. These include:

- a. Prawn and shrimp farming. This project encompasses training of riparian technicians in farming techniques, biological surveys of prawn resources and cultivable water areas in

the delta, feasibility studies for establishing a pilot prawn culture farm, and adaptive research and pilot operations in spawning shrimp and prawns under captive culture conditions.

- b. Catfish propagation. This project has the objective of developing practicable means for artificial propagation of the giant catfish *P. pangasius*, not only to enhance commercial farming of this species but to compensate for any losses in natural fry sources which might result from mainstream project development. The methodology could be subsequently adapted to conserve other valuable fish species which might be endangered by water resource development.
- c. Brackish water fish seed prospecting. To offset possible losses to estuarine and inshore fisheries due to dam construction farm fish production could be enhanced in the brackish zones of the Mekong delta by ensuring a reliable supply of seed for valuable brackish water species of both fish and shrimp. The project includes field explorations for locating seed sources, determining seasonal distribution and favourable exploitation times, fabricating gear for seed collection and segregation, and training.

Conclusion

The natural resources of the lower Mekong basin, particularly forest habitats and wildlife, are being depleted at an alarming rate, and unless this trend is changed most of the remaining forests and wildlife could be lost within the next several decades. However, there is considerable hope for the future *provided* that conservation is seen as part of overall land use planning which will yield optimal benefits to the regional economy. Conservation of the upland watersheds, including the allocation of sizeable areas for protected areas, must be included in all water resources development plans from the earliest stages. Conservation in the lowlands is also best based on a sound land use plan, but because most of the lowland areas are now being used for agriculture, conservation essentially means optimal agricultural development, as discussed at length in Part II.

With regard to freshwater aquatic fauna or fisheries, the natural fisheries must be supplemented by aquaculture as well as by the new fisheries of reservoirs, in order to meet the increasing protein needs.

PART II

The Agricultural
Development Programme

Chapter 7

Agricultural Systems

Introduction

Part I has discussed the wide variety of soils, climates, water regimes, and natural flora and fauna found in the Lower Mekong Basin, illustrating the point that the basin is very diverse. As is to be expected from such ecological diversity, many different agricultural systems are found in various parts of the basin, each adapted to local conditions. These same local conditions must be considered as essential ingredients in any agricultural development plan, so it is important to have an understanding of the existing agricultural systems.

For the purposes of this discussion, the basin agricultural systems can be divided into three broad categories: traditional agriculture; extractive agriculture; and modern agriculture (though the categories have considerable overlap and considerable variability is found within each category).

Traditional Agriculture

Traditional agriculture includes several systems that have been used in the basin for at least 2,000 years, thus demonstrating their essential soundness under traditional conditions. We divide these systems into two main categories: swidden agriculture (mostly in the uplands); and banded field agriculture (mostly in the lowlands).

For both categories, the most important crop is rice; both employ systems of permanent subsistence farming, suitable for closed agricultural economies with low population pressure. Upland farmers require at least 30 ha per family to maintain subsistence levels, whereas banded field farmers will generally be self-sufficient with 1.5 to 3 ha depending on the local situation. Thus the total area in use for each of the types may be approximately equal even though the population of upland farmers is only 10 percent of the total farmer population.

Swidden Agriculture

(Thai, Lao, Khmer: “rai”; Vietnamese: “mur”)

Swidden agriculture (“swidden”= burned field) is also called “slash-and-burn”, or “shifting

cultivation”. It is a rotational system based on cutting living vegetation in early winter, letting the vegetation dry during the dry months, burning it late in the dry season, and planting a crop of rice, corn, bananas, and a wide range of other plants in the fertile ash. After one or two crops, the field is left to fallow for at least 10 years, after which the cycle is repeated. Agricultural tools are minimal; the plow is not used and even the harvest is often reaped by hand, without the use of a blade. While no draft animals are required, most villagers keep dogs, chickens, and pigs; nevertheless most of their animal protein comes from fish and wild game (at least under traditional conditions).

An essential part of swidden agriculture is the great diversity of crops which are grown, to some extent mimicking the diversity of natural forests. The Lua (Lawa) of northern Thailand, for example, grow about 120 crops, including 75 food crops, 21 medicinal crops, 20 plants for ceremonial or decorative purposes, and 7 for weaving or dyes; the fallow swiddens continue to be productive for grazing or collecting, with well over 300 species utilized (Kunstadter, 1970). The most important crop is upland rice, and it is not unusual that 20 varieties of seed rice are kept in a village, each with different characteristics and planted according to soil type, fertility, and humidity of the fields.

Yields are often very high in upland swiddens. Depending on soil conditions and climate, rice yields may vary from 1 to 4 tons per ha, with an average of about 1.6 tons. In addition, the yield per unit of effort is typically higher than in lowland systems, giving the upland farmer more time for hunting, social interactions, and ceremonial activities. However, if the fields are used for more than one year the yields of many crops drop very quickly (see illustration 9).

ILLUSTRATION 9: NAN PROVINCE, NORTHERN THAILAND: YIELDS FOR SWIDDEN AGRICULTURE UNDER RAINFED CONDITIONS ON THE SAME LAND OVER A PERIOD OF THREE YEARS (KG/RAI)					
Cotton	Mung	Bean	Rice	Corn	Peanuts
Year 1	1.75	153	406	428	100
Year 2	127	178	302	101	185
Year 3	96	69	222	55	108

Source: Chapman, 1970

Although swidden agriculture has come under wide abuse as being destructive of forests and watersheds, it is highly adaptive to a wide range of conditions and when properly performed it may be one of the least harmful ways of cultivating areas where poor soils, steep gradients, and heavy rainfall make conventional farming methods unproductive or impossible. As practiced by stable groups, swidden agriculture is not particularly destructive of forest, land, or wildlife. Permanent villages are established, moving only if forced to do so by extremes of economic hardship, political disturbance, or population pressure, not as a logical consequence of their agricultural techniques (Hinton, 1970).

Sedentary swidden agriculturalists have a strong interest in maintaining the fertility of the village territory and practice several long-term conservation measures, including:

- preservation of stands of timber in and around the swidden to serve as a seed reservoir for new secondary forest;
- sophisticated control of fire (including fire breaks, fire fighters, and coordinated burning);
- early cutting of forest to retain soil moisture, reducing transpiration losses so that swidden soil is often more moist than adjacent forest soil;
- careful rotation of swiddens, using each one for only one year (a second burn or continued use typically leads to the invasion of *Imperata* grass, deflecting the succession from forest to grassland);
- a bush-fallowing period of at least 10 years to allow the flow of nutrients to reverse the trend toward leaching and be recycled into secondary growth plants, which then release their nutrients through burning;
- careful control of weeds (which may be the dominant reason for abandoning a swidden); and
- minimal disturbance of topsoil, thus minimizing erosion.

Under such ideal swiddening systems, wildlife flourishes, with elephants, wild cattle, deer, and wild pigs all feeding in the abandoned swiddens. When the swiddeners have tenure over their land, the game is carefully cropped, with over-hunting prevented by religious customary sanctions, as well as by primitive weapons. In addition, the ability to maintain a stable and permanent relationship with forested land enables some swidden farmers to invest time and effort in other permanent assets like fruit trees, fenced gardens, terraces, and irrigation canals. In the most suitable areas, the swiddens are supplemented by irrigated rice fields, thus allowing a considerably higher population density than under swidden conditions alone. These rice fields require buffalo for plowing and harrowing.

In areas above 1000m, the dominant crop is maize, supplemented by opium (to purchase rice). Such groups as the Hmong also keep large numbers of cattle which are slaughtered only on ceremonial occasions. The Hmong also see their cattle as insurance against extremely adverse conditions, to be sold in an emergency.

However, in recent times, swidden cultivation has revealed some serious limitations. Since per capita land requirements are relatively high and upland agricultural systems seem unable to respond to labor surpluses with corresponding increases in productivity, the pressure of increasing population has forced more intensive use of the available land.

The fallow periods are thus shortened, leading inevitably to a decrease in fertility and, ultimately, to *Imperata* grasslands which cannot be cultivated by traditional means. Thus the over-utilization of the potentially renewable upland agricultural resources has led to an essentially permanent depletion of those resources and reduced carrying capacity of the land.

The increase in hill tribe population is also resulting in unsuitable areas being cultivated. Permanent swidden farming can only be practiced on soils derived from igneous and basic sedimentary rocks, including shales, limestones, and basalts, or on well-structured red soils. All other soils tend to regenerate very slowly, requiring an excessively long fallow period. Many of the higher elevation areas previously avoided by swidden cultivators

because of their inherent limitations, are now being used by a very destructive type of swidden agriculture, which might be called “extractive swidden agriculture”. This system is based on using the same swidden for growing opium for 5 or 6 consecutive years (with other crops such as maize grown during the “off-season”), until all residual soil fertility is lost, soil organisms are removed, seed trees are burned, weeds become rampant, and topsoil is eroded; upon abandonment, the land reverts to *Imperata* grassland. The villages are temporary and of fluid composition, discouraging capital investment or long-term land conservation. Such areas are seldom self-sufficient, with opium being used to purchase rice from the lowlands or lower lying hill tribes.

Another variety of unstable swidden agriculture is practiced by an estimated 15 to 20 percent of lowland farmers. Lowland swiddening is typically very destructive because land titles are seldom involved in the clearing of reserved forests, so the farmers have little motivation to take long-term conservation measures. Short periods of cultivation are followed by periods of fallow which are too short to restore soil fertility, leading to permanent abandonment of the land. Fire control measures are usually neglected and since the lowlands are usually covered by fire-prone deciduous forest, the dry season often experiences massive destruction of forested land. The extent of dry season fire in the plains of northern Cambodia is especially notable.

While the Mekong Project is primarily concerned with the lowland areas, the uplands are seen as an integral part of the basin agroecosystem, and as such must be considered as an important part of the overall plan. The social structure of most hill tribes is organized around the holding and control of swidden lands (Kunstadter, 1970), but since each tribe tends to have its own set of customary laws (seldom recognized by other tribes or by the dominant lowland government), land disputes are very common. Only if the tribe unequivocally controls its swidden land will long-term conservation measures be effective, not only for the land and vegetation but ultimately for the tribe itself.

Even so, this will not be enough. Employment opportunities in forestry should be created, along with reforestation in the *Imperata*-covered uplands. Research on making the better upland soils more permanently productive should be conducted, and new cash crops should be introduced along with new systems of transport and marketing. Within its limits, swidden farming can continue to make a meaningful input into the total agricultural productivity of the basin, if it is developed as part of the overall system.

Traditional Bunded Field Farming

(Thai Lao: “naa dam”; Khmer: “sre”; Vietnamese: “ruong”)

More sophisticated from the point of view of traditional technology is bunded field farming, which has been practiced in the basin for at least 2000 years. Much of northeastern Thailand, southeastern Cambodia, and the Mekong Delta are covered with an intricate patchwork of small, traditional bunded fields. For such farming, the land has to be cleared (often by using slash-and-burn techniques), leveled, and bunded. Experience has taught that it often takes about one generation for a rice field to become optimally productive;

apart from perfect leveling, an impervious layer must form at shallow depths in paddies, to keep the rainwater ponded on the field with minimal infiltration losses. Contrary to swidden farming, no fallow periods are necessary other than the annual short fallow period during the dry season.

Plowing turns under the mulch from the previous year, the dry season fallow produces organic matter on which buffalo and cattle are grazed (adding manure to the soil), and the submerged conditions of the field during the growing period ensure enough nitrogen to be fixed biologically under the anaerobic conditions to maintain a relatively constant minimum yield. Some nitrate nitrogen is formed during the fallow period as well (Anderson, *et al.*, 1973).

While soil fertility plays a dominant role in swidden cultivation, the hydrological conditions of each field may limit the yield of the banded fields, since the drying-out process may make the soil more acidic and disturb the nitrogen-fixing process. As previously noted, the range of hydrological conditions in the Mekong basin is very wide, and here again, traditional technology has been adapted to these conditions with remarkable sophistication. Part of this adaptability is in the wide range of traditional rice varieties which are in current use, varying from varieties with an 80-day growing period (for drought-susceptible fields) to varieties with a growing period as long as 270 days (for lands that are moist over a long portion of the year).

Another part of this adaptability is traditional water manipulation. In many areas of the basin, banded-field farmers of the lowlands have since time immemorial made small canals which would help to spread wet season floodwater more equitably over their fields. In some areas, the rising waters of the flood periods were raised by temporary weirs and the water diverted for gravity-fed irrigation. Large ponds for domestic use during the dry season are widespread throughout the basin, especially in conjunction with temples. In a few areas, very large canals were built during the Funan and Angkor periods, and some of these are still in use.

Banded field farming typically involves monoculture; other crops are seldom grown on the same land. Moreover, it is very labor-intensive and requires agricultural tools and draft animals. While optimal upland swiddens may give a slightly higher yield in any given year, their long fallow period means that the sustained yield of banded fields is typically at least 10 times higher than in upland fields. Furthermore, the banded field system is more adaptable to population pressure. The yields of the banded fields actually increase when fields become smaller with population increase, because on smaller farms better care is taken of the crop, weeds are more carefully controlled, and more work is done to ensure satisfactory hydrological conditions.

The banded fields are characterized by finite patterns which have evolved throughout the ages and in some cases are specific for a certain tradition. Though prehistoric and early historic field patterns are unknown, three well distinguished systems are now common in the basin:

- a. The *medieval Khmer* system consists of small square fields with high and broad bunds, overgrown with shrubs and trees for utilitarian purposes. The Khmer system, common around the Great Lake in southern Cambodia and in parts of the Mun basin of northeast Thailand, indicates that the Khmers probably did not use the plow, but puddled the fields by trampling; this system probably developed in the low terrace lands. The Khmer field patterns show a remarkable regularity from the air, somewhat resembling the centuriations that are found all over the Roman Empire from England to North Africa and the Middle East.
- b. The *Vietnamese system* is characterized by long, narrow fields, sub-divided when necessary; the bunds are narrow and not overgrown. This pattern, which can be efficiently plowed due to the relatively few turns required, suggests that the plow was used from the beginning and that the system originated in the flat lowlands.
- c. The *Thai system*, which is not bound by tradition, is entirely adapted to the topographic and hydrologic opportunities of the terrain. On level land the fields are large and square; on sloping land they are long and narrow, following the contours of the slope. Sometimes bunds are high, sometimes low, as the need dictates.

The traditional banded field systems represent a feat of land reclamation of tremendous magnitude in the Lower Mekong Basin. Some 50 million individual banded fields exist in the basin, all constructed by traditional means without the benefits of agronomists, foreign agricultural experts, benefit-cost analyses, and loans from the international assistance agencies.

The rich store of down-to-earth technology for traditional agriculture efficiently adapted to each micro-ecological situation should not be under-estimated. It is remarkable how this invaluable knowledge has remained intact throughout the ages, in spite of historical vicissitudes. Civilizations may come and go, but the efficiency of the individual subsistence farmer seems to remain untouched. However, traditional farming systems evolved for traditional population levels, so their potential for further expansion is limited. Swidden agriculture is already over-extended and banded field agriculture is in many areas at the limit of maximum productivity. In northeast Thailand, half of the banded fields are marginal rice lands which give unpredictable yields and fail in many years; with traditional means the production ceiling in northeast Thailand would be limited to about 6 million tons per year.

Expansion of Traditional Agriculture in Mekong Delta, 1860-1975

Because both swidden and banded field agriculture were adapted to local environmental conditions and low population pressure, the surpluses from banded field farming historically were channeled into the construction of monumental religious structures or into the support of priesthoods and military establishments.

Beginning about 1860, traditional banded field farming entered a phase of rapid expansion. Stimulated by foreign markets and cash demand for rice surpluses, the area under rice

cultivation in the Mekong Delta increased from 170,000 ha in 1872 to 1,300,000 ha in 1908, an increase of about 7.5 times or a compound rate of 4.7 percent per year. The revolutionary opening up of new land subsequently slowed down, with the increase from 1908 to 1961 being only 270,000 ha, an increase of 21 percent in 53 years (0.4 percent/year) (illustration 11). Most of the increased production was exported, with the exports of rice from Ho Chi Minh City increasing from 58,000 metric tons in 1860 to 900,000 tons in 1910, an increase of 14.6 times in 50 years, representing a compound rate of 5.6 percent/year (illustration 10).

ILLUSTRATION 10: RICE EXPORTS IN THOUSANDS OF METRIC TONS, BANGKOK AND HO CHI MINH CITY, 1860-1910 (ADAPTED FROM OWEN, 1971)		
Year	Bangkok	Ho Chi Minh City
1860	96	58
1870	156	230
1875	237	341
1880	209	286
1885	225	455
1890	490	529
1895	498	567
1900	423	625
1905	861	714
1910	970	905

In the late 1860s, most of the Mekong delta was an unhealthy environment, with human settlement confined to the immediate vicinity of the major rivers and their tributaries. Although the Vietnamese had begun opening up the delta by constructing canals as early as the beginning of the 19th century, this development gained considerable impetus late in the century. The policy was to give large land grants to Vietnamese, Chinese, and French investors, who then recruited tenants to build canals, clear the land, and cultivate the fields. The government built only a few of the largest canals (sometimes following the same pattern as canals built during Funan times, 15 centuries earlier). The new canals provided some limited drainage, but their most important function was to provide access to markets through boat transport; actual irrigation was minimal.

These developments, using essentially traditional techniques but stimulated by foreign markets and some limited foreign technology, led to a great increase in population. In 1880, an estimated 1.7 million people were living in “Cochin China”, but by 1930, this figure had increased to 4.5 million, an annual rate of increase of 2 percent (Netherlands Delta Team, 1974). The population has subsequently increased to about 20 million.

A similar development took place in the lower Chao Phya delta, stimulated initially by the Bowring Treaty with Great Britain in 1856. The large and continuous foreign demand for rice, not a spontaneous development of her national economy, stimulated Thailand to develop rice cultivation in the lower delta of the Chao Phya (Tomosugi, 1965), and the export of rice from Bangkok rose from 96,000 tons in 1860 to 970,000 tons in 1910, an

increase of 4.7 percent per year (illustration 10). The objective of increasing food supply was attained not by raising the productivity of the land, but by expanding the area under cultivation. The development of canals and drainage projects was the key factor in this expansion; the Rangsit Canal, for example, was begun in 1908 and rapidly converted a thinly-populated swamp north of Bangkok into a thriving agricultural area where tens of thousands of rice farmers came to settle.

In both the Chao Phya and Mekong deltas, rapid expansion of agricultural land was in some ways detrimental to sound agricultural development. The way rice was farmed underwent virtually no significant changes; yields rather than increasing seem to have fallen slightly between 1880-1890 and 1930-1950, apparently due to the application of traditional technology to land somewhat less suitable for agriculture (Owen, 1971), though the time-lag factor in obtaining optimal yield from new rice fields may also have been a factor. The increases in crop yields which characterized agricultural development in other parts of Asia (Korea, Japan, China, Taiwan) were rendered unnecessary by the ready availability of large expanses of new land in the river deltas of southeast Asia.

The expansion of agricultural land and traditional agriculture has now reached its limits in the lowlands and (as noted earlier in the discussions on swidden and extractive agriculture) has exceeded its limits in the uplands. The only alternative for increasing agricultural productivity is to increase the yields on the available land, and this can best be accomplished by the techniques of Modern Sustained Yield Agriculture.

ILLUSTRATION 11 AREA (IN THOUSANDS OF HECTARES) PLANTED IN RICE, MEKONG DELTA, 1872-1908				
Year	Trans Bassac	Upper Delta	Lower Delta	Total
1872	20	51	99	170
1878/9	47	60	129	236
1888	204	135	275	614
1898	382	160	363	905
1908	571	242	455	1268
Trans Bassac: Bac Lieu, Can Tho, Rach Gia, Soc Trang, Ha Tien Upper Delta: Chau Doc, Long Xuyen, Sa Dec, Tan An Lower Delta: Ben Tre, My Tho, Tra Vinh, Vinh Long				

Source Owen, 1971

Extractive Agriculture

The basin countries possess a vast source of wealth in their natural forest vegetation; but while the forests are very rich, the forest soils, particularly in the upland areas, tend to be rather poor, with most available nutrients contained in the living and decomposing

vegetation. Thus these areas are fertile only when maintained in forest, or in swidden agriculture (subject to the constraints discussed above).

When the upland forest soils are cleared, crops can be grown for only a few years before inevitable depletion of soil fertility follows. When such soils are over-utilized (“mined”), the land reverts to a low-productivity grassland. This is now the situation in much of northern Laos, northeast and northern Thailand, and parts of the Annamites; it was historically the case in northern Cambodia, where much of the area was cleared of forest during the Khmer civilization, resulting ultimately in permanently depleted soils which can still be clearly seen by satellite imagery.

“Extractive agriculture” as defined here, refers to exploitation systems that deplete the soil or deplete the vegetation cover to the point where regeneration becomes impossible or impracticable and the land must be abandoned. The system is therefore not self-sustaining and is in fact self-destructive. However, extractive agriculture has the advantage that it gives an immediate high return at relatively low short-term cost, but at the expense of depletion of basic natural resources, particularly soils, forests, wildlife and watersheds. Therefore, if it must be used at all, it can only be applied for a period of limited duration, resources permitting, to raise the economy of a country or a region above the subsistence level. Care must be taken to ensure that at the appropriate time extractive agriculture can be transformed into systems providing more sustainable yields, and this will require that some of the profits obtained from extractive agriculture be reinvested in the more productive systems.

Extractive exploitation of the natural forests has been going on for centuries, with earlier efforts primarily carried out by the fringes of the lowland civilizations, where attempts were made to utilize soils which were basically unsuited for agriculture. More recently, extractive forestry has been practiced by recent hill tribe immigrants as a side effect of growing opium. Ultimately, wide areas have become grasslands, forcing continuing migration of these hill tribes. A terminal situation has now been reached in much of northern Thailand and Lao PDR.

Only since the 1950s has clearing of the lower-lying evergreen and deciduous forest land for commercial cropping expanded very rapidly, following the development and expansion of international markets, especially for feed crops. Of the basin countries, Thailand has experienced the most extensive clearing of forests for extractive commercial agriculture. During the period 1950 to 1975, the upland crops cotton, sugar, kenaf, cassava, and maize added over US\$ 4,000 million to the foreign exchange earnings of Thailand, an appreciable increase in wealth for the country.

However, increasing amounts of marginal land unsuited for permanent production are being forced into temporary cultivation, so the long-term contribution of these upland commercial crops to the economy of the basin is likely to be considerably lower than their present contribution (at least under the current method of extractive agriculture). The effects of clearing forest may have long-term costs which are considerably greater than the short-term benefits. The cleared land, being low in soil nutrients even when covered in

forest, apparently does not regenerate into forest very quickly, being essentially permanent wasteland which can contribute little to economic resource development. Perhaps the best use of this wasteland is for an intensive reforestation program.

Modern Agricultural Systems

Modern sustained yield agroecosystems include ranching, integrated livestock, forest farming, commercial plantations of specialized crops, and irrigated farm systems. While these types may be combined in some situations, we will discuss them separately.

Ranching

Ranching is an appropriate extensive land use system in dry temperate grassland areas lacking population pressure, but natural forest vegetation and population pressures on the land in the Mekong basin make extensive cattle ranching a generally inappropriate approach to land use in most of the basin, while small farms exclusively managed for beef production fail to generate sufficient cash income to be attractive. Grain and feed concentrates are regarded as ill-suited for the basin, due to the lack of premium beef price incentives, human competition for grain sources, and the more efficient conversion of these feed sources by poultry and hogs. Crump and Sanoh (1972) have shown that dairy farming is not likely to be a major agricultural activity in the basin, though it may be profitable for individual farmers in a few special situations. A changeover from small farm beef production to dairying might be feasible at a much later stage of livestock development. Dairy product imports, totaling over US\$ 30 million annually in Thailand, consist mainly of butter, condensed milk, and dry milk powder and are not likely to be replaced by local dairy products for the foreseeable future.

Generally, the potential for extensive ranching in the Mekong basin is fairly low at present, but a few areas might be suitable for ranching in the longer term. These include the plains of northern and eastern Cambodia, which are sparsely-populated grassy savannas which have a wide range of wild herbivores and might be best developed for "mixed ranching" using both domesticated and wild herbivores. At present, Thailand exports large quantities of maize and cassava, primarily as animal feeds. In the long term, it might be worth investigating the possibility of using these feeds to produce the meat within the basin, and exporting meat rather than animal feeds.

Integrated Livestock

Of greater potential for the basin is integrated livestock, where the domestic herbivores are managed as part of an integrated crop and livestock enterprise. Under this system, the animals primarily subsist on crop waste residues, weed growth, and associated natural grazing on areas unsuited to cropping, and where little pasture improvement is attempted.

The animals provide draft power for crop cultivation and farm transport and, as a by-product, accelerate cycling of nutrients to maintain fertility on crop lands. Sale of surplus animals and consumption or sale of animal products can be seen as a banking system and a hedge against inflation. It is apparent, however, that this system is not capable of producing a surplus of beef or buffalo sufficient to meet projected requirements, particularly if economic development is expected to result in higher protein consumption by the farmers.

It has often been suggested that the growing of fodder crops should be encouraged since no part of the basin is considered unsuited for planted pastures, and improved species are available which are well adapted to the various soil, moisture, and temperature regimes in the region. Experimental farms have revealed some promising forage crops, both grasses and legumes, which could support 1.5 head of cattle per ha for rainfed pastures. However, such pasture use is not, in itself, a sufficient economic asset to warrant the considerable investments needed.

A more promising approach would be based on the rotation of grass/legume pastures with rainfed cash crops (maize, mung beans, peanuts, cotton, cassava, and kenaf). Studies have shown that for this system, the individual farm size would have to be about 8 to 10 ha. About 6 million ha of upland areas are estimated to be suitable for development under this system in northeast Thailand, thus providing for a minimum of 600,000 farms. While most farmers will also grow some rice in nearby bottomlands, if such areas are available, the typical farm will have about 3/8 of the land in cash crops and 5/8 in grass/legumes pasture. In marked contrast to swidden agriculture, where the fields are rotated, this system is based on the sophisticated rotation of crops. But since this system is new to the basin, much assistance, both technical and financial, will be required if it is to become established.

The proposed rotation pattern would maintain soil fertility for sustained agriculture in suitable upland areas, farmed with the management abilities and available labor supply of contemporary northeastern farmers. It would allow a relatively constant production of cash crops at a somewhat higher yield than at present, while producing about 10 times the present amount of cattle per unit of land. The number of cattle could be even higher when by-products from cassava and mung beans are available as supplementary cattle feeds. It is projected that annual cattle production would be about 160 kg/ha/yr. Still, it should be remembered that this is a hypothetical system, which has not yet been actually demonstrated to be economically feasible.

Forest Farming

The situation discussed in Chapter 4 suggests that steps should be taken promptly to improve forest production in the basin, especially in Thailand. One way to accomplish this is through the forest farming system, where a crop of trees is grown along with a cash crop such as maize, cassava, or kenaf. As the cash crop depletes the soil and the yield drops, the crop is no longer grown, but the trees which were planted with the crop will continue to grow to a harvestable age, restoring nutrients to the soil so that the cash crop can be

grown again after the harvesting of the tree crop. Forest farming can, in effect, transform extractive agriculture into a long-term sustained yield system. Trees would thus be viewed as capital on a farm, in much the same way as livestock, though capital management would be an even longer-term proposition.

Karlberg (1976) has described such a system with kenaf grown with pine and suitable indigenous hardwoods; the system would be suitable for various land reform areas where the soils have already been depleted. The costs for growing trees in areas already planted to kenaf would be very low, since the heaviest cost for establishing forest plantations is for clearing the land. Annual yield of wood products should be at least 15 m³/ha/year, about the same as the annual fiber production for kenaf; however, the raw material costs for trees would be lower since planting and harvesting are easier. For kenaf and cassava, harvesting and income would come every year, while the tree crop can begin to be thinned only after 6 to 8 years. The trees would utilize the ground when the yield of kenaf and cassava is dropping, and by having an even distribution of age classes, the trees would give a cash crop every year, thus yielding a safe sustained annual income.

Karlberg (1976) recommends establishing a Forest Farmers Cooperative as part of the system. This cooperative would act as middleman, buying wood from the farmers and selling it to industry (sawmill or pulp-mill) as well as running a nursery for the pine seedlings (with costs for seedlings deducted when wood is delivered by the farmer).

While Karlberg's forest farming system was designed for northeast Thailand, similar principles can be applied in other upland areas which have been disrupted by extractive swidden cultivation of opium, or in any swidden area where rotation cycles are becoming dangerously shortened. While growing an upland rice crop for one year, the upland farmers will be encouraged also to plant a tree crop in the swidden, which will then be used for producing wood for 15-20 years. In essence, this is planned rather than random regeneration, a system that has been applied widely in Myanmar (where it is called the "taungya" system).

In addition to supplying construction wood, fuelwood, and pulp, forest farms can also produce the raw material for liquid fuels to replace fossil fuels. Aerobic fermentation of wood materials is an economically feasible process to make such fuels as ethanol, methanol, and butanol, in amounts equaling 40 percent of the dry weight of the raw material.

Finally, forest farming would also provide extensive employment opportunities for much of the currently underemployed labor force in both the lowlands and highlands. The major drawback to the forest farming system is that it involves a 15-to 20-year cycle, thus requiring considerable time and investment to become established. It would also require some basic changes in land tenure regulations on forested lands.

Commercial Plantations

Certain garden crops, such as onions, chili, mushrooms, cucumbers, eggplants, string beans,

pineapples, grapes, oranges, rambutan, longan, and mulberry leaves (for silkworm production), can be economically very attractive in the lower basin. In addition, as canning and freezing plants increase in the basin with cheap hydroelectricity, the prospects for increased export of fruits and vegetables are quite promising. In a few localized situations with appropriate soils, large corporate or state farms for crops such as pineapples may be feasible. Pineapple exports from Thailand, for example, totaled nearly US\$14 million in 1974. Still, highly specialized commercial fruit and vegetable growing will occupy only a relatively small area and will not be discussed further in this section.

In the high rainfall zone on deep red soils in Lao PDR, Cambodia, and Vietnam, plantations of rubber, tea, and coffee are feasible, especially when extended and integrated with field crops. In the case of rubber, both Cambodia and Vietnam have much experience in growing and marketing the crop, and post-war conditions are likely to see a great expansion of rubber plantations and considerable increases in production per hectare. The plantations will become even more economically attractive when more rubber processing plants and tire factories are built in these riparian countries.

Coffee grows best on well drained red soils. Tea is native to parts of the Mekong basin, growing best on well drained acidic soils with high rainfall. Both of these crops can be productively grown only in relatively confined areas of the basin and are not likely to become dominant export crops; however, both crops can form the basis of labor-intensive processing plants, furnishing employment for thousands of workers.

Irrigated Farm Systems

Ranching, integrated livestock, forest farming, and commercial plantations of specialized crops have various degrees of feasibility for certain areas within the Mekong basin, but none of these has the potential to be widely applied and to meet the greatly increased requirements in the foreseeable future. The only sustained-yield farm system that is currently available to be applied on a broad scale is the irrigated farm system, including water control and systematic irrigation, along with supporting infrastructure and hydroelectric energy. With full development of the ongoing and proposed irrigation schemes, nearly 4 million ha will be affected, totaling about half of the currently planted area. Apart from some uncultivated floodlands, which will become cultivable with water control, the irrigation schemes will serve existing farmlands.

Irrigation will have many advantages. Cropping diversity will be greatly increased, with a wide variety of crops grown on what was previously monoculture rice-growing land; these will include maize, peanuts, cassava, tobacco, mung beans, cotton and a wide variety of fruits and vegetables. Cropping intensity will also be greatly increased, ranging up to 200 percent (double cropping). Yields will dramatically increase, with rice yields, for example, increasing from 0.7-1.6 tons/ha to 3.0-4.0 tons/ha or even more with high yielding varieties. In addition, yields in non-irrigated areas will also increase modestly, due to improved extension markets, and other services that will be stimulated by the greater economic productivity of irrigated areas. Net farm income and employment opportunities will be

double, while land value will be tripled.

Because of the pervasive importance of irrigation, the next chapter will discuss how it will be applied in the various agricultural zones in the lowlands, how it fits into the agricultural development model, and how it contributes to the long-term development strategies for the entire basin. Before going into these development aspects, however, it will be useful to first consider some of the problems that have been encountered with existing irrigated agricultural systems.

Problems with Irrigated Agricultural Systems

Whenever countries attempt to establish new irrigated agricultural systems, as has been advocated by the Mekong Project, some difficult problems are invariably involved. Before these problems can be solved, they must first be clearly identified. Ten problem areas identified relevant to irrigation in the lower basin are:

1. Inadequate water supply

The adequacy of the water supply is related to several factors including the basic water supply and reservoir or river operation impact (see Chapter 1). Even though the water supply may physically exist, it may be impossible to deliver it to the individual farm unit in the amount or at the time desired. The most common reason for this is the lack of adequate canal or lateral capacity. The limited capacity may be attributable to inaccurate determination of farm water requirements, differences of opinion regarding the irrigation methods to be used, or inadequate maintenance.

The farm water requirement is a function of consumptive use of plants, deep percolation, evaporation, and runoff. These in turn are affected by the number of crops grown per year, the evapotranspiration rate of the particular crops, the texture and slope of the fields (the drainage characteristics of the soil), the length of the runs, and the irrigation method selected.

In the Lower Mekong Basin, which has two basic land types (uplands and lowlands), soil differences can be very important. The flat heavy textured lands, which are commonly underlain by hardpan and characteristically used for rice production, have a different water requirement than the uplands which are usually lighter in texture, have pronounced variation relief, and are open and freely drainable. One of the main problems in irrigating these upland soils is poor management leading to excessive deep percolation.

The economics of design of canals and laterals are often in conflict with the farmer's desire for water delivery. Generally speaking, the canal designer will assume that water will be delivered on a rotation basis which minimizes the canal capacity required and related costs, while the farmer would prefer to have water delivered on a demand basis (whenever

he wants it). This increases the canal capacity requirements with attendant increases in costs, as well as presenting additional management problems. The manager has to increase or decrease the amount of water in the canal depending upon the collective farmers' orders. Otherwise, at certain times more water is diverted into the canal than is needed, and water is thus wasted.

The matter of how the water is delivered to the farmer (demand or rotation) is extremely important to him. This is particularly true of the upland farmer because the upland soils have a low moisture-retaining capacity and crops become stressed very quickly. For this reason, farmers like to have a water supply available so they can irrigate immediately before the crop reaches the wilting point.

As a general rule, farmers the world over are reluctant to buy improved seeds, fertilizer, and pesticides unless they have considerable assurance of the required supply of water. Water is necessary to secure the expected responses from improved practices, particularly the application of commercial fertilizer. Farmers simply will not make the investment in additional or more expensive inputs if they do not have an assured supply of water when they need it.

Therefore, the design of the system and the method of delivery of water have much to do with how farmers use water and the response they get from it. The plans for use of water supply should be worked out in close consultation with those who will actually use the water.

2. Inclusion of non-irrigable lands

Many projects have failed or proved to be less successful than expected because of soil problems, including harmful accumulation of salts and alkali, water-logging because of poor drainage characteristics, excessive erosion or water requirements because of light texture of the soil or excessive slope, and limitations of productivity because of shallowness of the soil.

Other kinds of soil problems develop as a result of the manner in which the project was developed. One of the most common examples is the exposure of subsoil, hardpan, or gravel by excessive or improper leveling. This is a serious planning mistake, and somehow it continues to be made in the development of new irrigation projects; but it can often be avoided by selecting the sprinkler method of irrigation (though this is only economically justified for specialized crops).

3. Drainage

It is now generally recognized that successful irrigation requires the ability to remove water from the land at will as well as to apply it, so in most cases the service canals and laterals constitute only part of the total irrigation facilities required. Perhaps partly due to the

difficulty of anticipating where and when drainage problems will develop, it is often tempting to ignore the potential drainage requirements; hence it is important that drainage facilities be included in the initial plans and estimates. If the drainage water is not polluted by harmful accumulations of salts or alkali, it can be reused, and project costs can often be minimized by making provision for the recycling of such drainage water.

Drainage and periodic flushing are particularly important in those lands that have, for example, salts in the soil profile. In the process of irrigation, these substances are dissolved and brought to the surface of the soil by evaporation, leading to concentrations of the chemicals on the soil surface. This makes it impossible to germinate certain seeds, i.e., many crops will not tolerate even moderate concentration of these chemicals. In some cases the problem can be controlled by drainage and water management; however, badly affected areas should be excluded from the service area. Fortunately, with modern technology, it is now possible to identify such areas in the process of selecting the lands for service (land classification).

Another facet of the drainage problem in tropical areas, where double or triple cropping can be practiced, is the need to remove water so one crop can be harvested and the seed bed prepared for the next in as short a time as possible.

4. Operation and maintenance of irrigation facilities

Many irrigation projects encounter difficulty in delivering adequate water, not because of inadequate supply, design, or canal capacity, but because of inadequate maintenance. The operation and maintenance of irrigation systems is not very glamorous; it does not attract the public attention as does construction of a large dam and reservoir. But maintenance is as important as construction to the success of an irrigation project and must be provided for in the planning stage.

An improperly maintained irrigation system will quickly deteriorate to the point where it will be impossible to make adequate deliveries. Farmers adjust to this situation by reducing the irrigated area or limiting the crops to those that can be produced under natural rainfall, thus defeating the goals of the project. Common problems of maintenance include loss of capacity because of silt accumulation or weed growth, and inability to distribute water because of deterioration of diversion structures and head gates. Loss of water by seepage is another common problem, affecting the project water supply and waterlogging the lands immediately below the canals, thus seriously limiting their usefulness.

One solution to the maintenance problem is to employ a highly qualified staff, including ditch riders, who will deliver water to individual farm units under the method of water delivery being employed; but it generally takes a rather large and prosperous irrigation enterprise to afford this kind of staffing. An alternative that has worked particularly well on small irrigation enterprises is for the water users themselves to take over the operation. When this system is followed, it is generally the practice to elect a manager and a Board of Directors from among the water users. In some cases the Board employs a trained manager.

The manager and Board are then responsible for managing and directing the activities of the enterprise. This system has been known to work very well, particularly in those cases where the water users own the system; the farmers then have a proprietary interest in the irrigation system.

5. Water control

Certain facets of this problem have already been discussed. It has been noted that a farmer should have control of his water supply so that he can put it on and take it off his fields at his will, especially where multiple cropping is expected. In order to facilitate such control it is essential that a service outlet be provided for each individual farm unit. With small and irregularly shaped units, this often becomes difficult. Modern ways of delivery, for example by constructing concrete "canalettes" may be too costly to be practical. It is therefore preferable to plan an open ditch gravity delivery system for the basin, requiring careful and detailed planning. Land consolidation may be an important part of this planning.

One of the problems in making such delivery, particularly in areas that are already farmed, is the acquisition of the right-of-way for the canal or lateral. Because of the lay of the land, such canals or laterals often dissect established farm units. The owners of such lands are not always pleased by the prospects of having a canal cut through their farm and generally oppose such construction. Because of this problem, most countries where irrigated agriculture constitutes an important part of the economy have enacted legislation which requires that any person must make available a right-of-way across his land if needed to serve adjacent lands (this, of course, is done with reasonable compensation). The possibility of isolating land from a source of water is thus eliminated.

6. Full farm development

For areas that are not currently used for agriculture, it is necessary to clear and level the fields for irrigation so the water that is delivered to each farm unit can be effectively used. The leveling (slope of the fields) will depend upon the land type, texture or soils, and the type of irrigation to be practiced (rill, furrow, flood, border, or sprinkler). For instance, with flood irrigation it is desirable to prepare the lands as flat as possible, but with furrow irrigation it is generally desirable to level the lands so the field has some fall. The fall or slope of the field and the length of the run is related to the texture of the soil and the heads of water to be used.

In the Lower Mekong Basin much of the future irrigation is projected in lowlands that are now in use as banded rainfed fields. The problem in this case is to adapt the field pattern to the irrigation/drainage system. In many cases, land consolidation and field re-leveling is necessary. The irrigation system, the land preparation, and the method of irrigation to be used should be properly integrated so they will be compatible. This will be much more of a problem on the uplands than in the flat soils which are generally used for rice production.

7. Farm management practices

After the irrigation system is developed and the farm unit has been prepared for irrigation, the farmer is then ready to engage in irrigated agriculture. At this point he is then faced with a multitude of decisions such as which crops to produce, which varieties to use, the fertilizer to be used and how much, what plant diseases and insects can be expected and how should he control them, how often should he irrigate, how much water should he apply, etc.

If the farmer has had irrigated agriculture experience, he may very well know the answers to many or even all of these kinds of questions; but if he has had little or no experience, then he will likely need much guidance. Thus provision of demonstration farms and farm extension services should be considered an integral part of the irrigation development program.

8. Farm supplies and equipment

The development of a newly irrigated area often creates large demands for farm supplies, equipment, and machinery. In established areas such products can normally be procured, but in new large areas this may not be the case, so it is important that necessary trade centers be established and that the farm supplies that will be needed in the development of the newly irrigated lands will indeed be available.

9. Credit

It is evident from the foregoing that the development and operation of a newly irrigated farm is expensive. If the farmer has to buy the land, clear it, level it, and build his home, he will likely need some sources of long-term credit to finance his operation. In some cases these may be provided as part of the project cost and plan.

Intermediate-term credit may be needed for the purchase of live-stock, farm equipment, and machinery; and short-term credit may be needed to finance the current production expenses. Moreover, all these types of credit should be available at reasonable interest rates; otherwise farmers have little incentive to borrow money and assume obligations to repay. If available at reasonable interest rates, credit can considerably accelerate the return from irrigation projects.

10. Public services

Life in a newly irrigated area may sometimes be a discouraging experience, but with sufficient thought and attention to provision of essential public services some of the hardships can be removed and with modern technology and capital the process of farm unit development

can be greatly accelerated. The basin now has examples of irrigated farms that have been completely developed within two or three years.

The difficulties of adjusting to the new environment can be reduced also by providing some of the public services or amenities that are usually associated with “the good life”. These include such things as roads (closely related to the question of markets discussed earlier), potable water supply, schools, public meeting places, electric services, and medical and public health services. The availability of these kinds of services will provide a higher quality of life for the new farm family and thus accelerate the development of the irrigation enterprise.

The provision of these kinds of services can often be expedited by settling the farm families in cohesive villages; the farmers can then go into the fields each day to perform their farm tasks. The village method of settlement results in economies in the provision of public services and facilitates the development of a cohesiveness in the society and a spirit of cooperation.

From the standpoint of efficiency in the management and operation of the individual farms, the village type of settlement has certain disadvantages. Nevertheless, in a new development the advantages to the community pattern of settlement warrant its serious consideration.

Conclusion

Traditional agriculture, including both swidden cultivation in uplands and paddy rice in the lowlands, has now probably reached or exceeded its productive limits in the Lower Mekong Basin. Further increases in production will depend on a more intensive use of the lowland areas through modern sustained-yield agriculture. While forest farming, ranching, and commercial plantations all have important roles in the agroecosystems of the basin, only irrigated agriculture systems have the potential for the massive increases in food production needed to meet minimum projected food requirements for the lower basin. Such irrigation on a wide scale will also require large supplies of water but with adequate storage these needs can readily be met by the substantial water supply resource of the Mekong river.

Chapter 8

The Agricultural Development Model

Introduction

In 1958, an FAO mission led by G.H. Bacon was assigned to study the available basic data which related to the agricultural, fisheries, and forest production of the Lower Mekong Basin, with special reference to the potential for water resources development. The mission was asked to determine the scope and nature of the further investigations which would be necessary to provide an integrated knowledge of the agricultural resources of the basin, and to indicate how these resources could be best utilized in the light of the anticipated development of the water resources of the Lower Mekong Basin. The Bacon mission made 27 recommendations, including: compiling data on climate, soils, and natural vegetation; studying fertilizer requirements, fisheries, salinity, current land use, development of forest industries, irrigation, and hydrology; establishing a set of experimental stations and pioneer agricultural projects; and training students in agricultural fields in foreign countries (FAO, 1959).

In 1961, the Ford Foundation sponsored a team of White, de Vries, Dunkerly, and Krutilla, which was invited by the Mekong Committee to study economic and social aspects of Lower Mekong development (White, *et al.*, 1962). In the field of agriculture, they recommended that: (a) a team of agricultural experts be organized to assess water management as a means of agricultural development; (b) an area of 3,000 to 5,000 cultivated hectares be established as a comprehensive demonstration project where the full impacts of Lower Mekong management upon rural life could be observed (this is essentially a pioneer agricultural project); (c) studies be conducted on opportunities for fish cultivation in ponds, reservoirs, and natural lakes; (d) prospects be assessed for marketing of rice and other potential agricultural products from irrigation, flood-control, and drainage projects; (e) maps of soils, water balance, and vegetation types be compiled from already existing material; and (f) land capability, land use forest cover, and settlement patterns be inventoried by aerial photo interpretation.

These two reports were not designed to solve problems, serving rather to form the early conceptual background for the agricultural work of the Mekong Committee. This was followed in January 1969 by a request from the Mekong Committee to FAO and UNDP to carry out a review of the experimental and demonstration work on irrigated agriculture initiated by the Committee on the basis of the Bacon and White recommendations. The joint FAO/UNDP mission was also asked to recommend guidelines for long-term planning of agricultural development in the basin.

In July and August 1969, the joint mission visited the four riparian countries under the

leadership of Mr A. Goldschmidt. The mission found that the work of the agricultural stations sponsored by the Mekong Committee was not sufficiently well coordinated and the research projects were not clearly related to each other; nor were the results of the studies and research being made available for application and use. The stations were found to differ in stages of development, in operational arrangements within the governments, and in the adequacy of their budgets. While as a whole they were considered to constitute a good beginning, their work needed to be augmented and related to the other research and experimental work carried out in the basin.

The joint mission recommended that the Mekong Committee should expand to include an Agricultural Division within its Secretariat, with the Director of the division working directly under the Executive Agent. The joint mission also recommended that an Agriculture Sub-Committee of the Mekong Committee be established, consisting of four Directors-General of the Agriculture Ministries of the riparian Governments, to ensure that cooperation would not only cover the stations and related works of the Mekong Committee, but would also include all similar undertakings of the riparian governments for agriculture research, demonstration, extension, training, agricultural planning and institution-building relevant to the Mekong program. The Goldschmidt report also pointed out that any major change-over from existing subsistence agriculture to more intensive use of the land must be justified by substantial economic gains and, above all, by an equitable and wide distribution of benefits among the people of the four countries concerned.

The program which has been subsequently developed by the Agriculture Division follows the basic concepts presented by the review teams. With the recent important breakthroughs in agricultural research, particularly in the development of new high-yielding varieties requiring modern techniques in the use of fertilizers, herbicides, pesticides, and water control, Mekong development becomes even more critical to the future of agriculture in the countries of the basin. It is therefore essential that research and experimentation be directed to applying the new methods to the various physical and human environments within the basin. Changes will have to be introduced step by step, with research and experimentation oriented accordingly so that all proposals for improvements will be realistic in terms of the existing constraints of the farmer: financial, economic, educational, and others. The Mekong Agricultural Research Program must assist in creating the basis for a strategy of development that will ensure the quickest possible realization of the benefits of water control provided by the various projects.

Plans for the development of agriculture in the basin should not be limited to irrigation with full water control but should encompass all patterns of cropping from dry farming to irrigation by gravity with full water control, with numerous intermediate solutions such as irrigation with partial water control, drainage only, pumping ground water, and pumping directly from the mainstream or tributaries. Watershed management with improved shifting cultivation should also be included in the development program.

The experimental and demonstration program should supply the essential data for feasibility studies of major and medium-sized irrigation schemes, and at the same time provide a firm basis for the planning or other agricultural aspects of the Mekong program, including the

establishment of priorities and timing of development. The results of these studies will help the basin governments in planning their country development plans. The experimental programs of the individual stations, their locations, and their relation to other stations in the basin, should not be limited by currently envisaged projects, or indeed to any immediate program. Circumstances and conditions in the basin are certain to change over time, and the experimental stations must be geared to help agricultural practices evolve with the changing conditions. The experimental stations will also keep abreast of pests, diseases, and other dynamic environmental factors.

The combined research, experimentation, demonstration, extension, and institution building must be woven into the work of the Mekong countries and dispersed throughout the basin in order to take account of differing soil and moisture conditions, crops, and cultural patterns. This is the purpose of the pioneer projects, which represent an intermediate step between the demonstration area and full project development. In a sense, these projects are pioneering the larger projects, and should be on a scale that will make possible all necessary economic, social, and institutional inputs, including credit and marketing facilities. The scope will vary, but in principle the size of a pioneer project should not be broader than required to represent an economic unit where all factors and their interactions can be tested on a “real-world” scale.

In conclusion, the agricultural development model adopted by the Mekong Committee is based on three major consecutive steps: agricultural experimentation and demonstration (Chapter 8); pioneer agriculture projects (Chapter 9); and full project development (Chapter 10). Some introductory aspects of each of these steps are discussed below, with further details in the subsequent chapters.

Role of Agricultural Research

Farmers have always been innovators. Traditionally, innovation has been a slow but sure process based on long-term knowledge of local environmental conditions and on a judicious balance between needs and resources. Typically, the process can be speeded up only by new approaches that lead to improved productivity. Thus agriculture has passed from the gathering of wild crops to cultivation, from the swidden field to the rice paddy, from the digging stick to the plow.

The industrial revolution brought new pressures for increased agricultural production, through the factors of exploding human population and accelerated demand for crops for exports, stimulated by newly-accessible markets and the creation of newly perceived needs. The traditional process of slow but sure innovation was unable to respond very well to the needs of modern agriculture. The inability of traditional technology to meet increasing needs has unfortunately tended to set the stage for accelerated clearing of new land (often unsuitable for permanent agriculture) and the reduction of the fallow periods. This “extractive agriculture” inevitably leads to rapid depletion of potentially renewable natural resources.

Thus it is necessary to call on modern agricultural research, based on rigorous scientific

techniques and a worldwide fund of information and assistance. However, whether it is used as a means of accelerating the traditional process of innovation or as a tool for the introduction of exotic technology, agricultural research often meets resistance from the traditional farmers. The effect of such resistance is not always negative – it often eliminates technologies that do not bring decisive and certain improvement to the farmer. The resistance is based on numerous constraints, including financial, economic, and technical problems (growing calendar, crops, water control, etc.), or even the simple matter of taste. Research should not ignore any of these aspects of the farmer's life, but should attempt to integrate innovations into the traditional cultural framework.

Agricultural research, taken in a very general sense, is thus a starting point for all development projects, for efforts to halt the destruction of renewable resources, and for the introduction of new, more productive technology. This section does not record all of the agricultural research done in the basin or even give a summary; the more detailed results are found in the following chapter. The purpose here is rather to clarify the lines of research which have been followed, to present a few examples of significant results, to enquire about the reasons for sub-optimal results or even failures, and to suggest several lines of research for the future.

Introduction of New Technology

The phrase, “introduction of new technology”, is used here to refer to the research work that has been accomplished so far in the basin, especially studies of crop varieties or techniques which have been imported for testing at the various research stations. The “New Technology” is usually compared with a control which hypothetically represents the local conditions.

Regarding varieties, excellent results have been obtained for second crops like maize with the Guatemala variety in Thailand or with the soybean varieties SJ1 and SJ2 (from Japan and Taiwan), also in Thailand. But the situation is different with rice, which has been cultivated for centuries in the region where the twin forces of selection by man and by nature have resulted in varieties perfectly adapted to the local conditions. The Mekong Committee paper, “Note on the Role of High Yielding Varieties of Rice in the Lower Mekong Basin”, shows that only in the Mekong delta have the imported hybrid varieties found the necessary conditions to express their potential: control of water; subsidies which permit the use of fertilizers; and strong demographic pressure.

This last factor has several correlates, including pressure for higher yields (and therefore the utilization of fertilizer) and the necessity for several crops a year (therefore encouraging varieties that have a short growing cycle and are photo insensitive). Outside the delta, the high-yielding varieties are grown only in a few irrigated areas, mainly in the dry season; they seldom compete with local varieties under traditional conditions.

Introduction of new technology also includes research on the use of fertilizer, based on variations on the N/P/K (nitrogen, phosphorus, potassium) complex and their recommended

dosages. Except in the past few years, this field of research has suffered from needless duplication, inadequate research design, and too few trials in the farmers' fields. Outside the Mekong delta, the benefit of fertilizers in doses as recommended by various government agencies, remains marginal on most crops; the few times that the economic feasibility of fertilizers has been calculated, profitability has been shown to be nil or very low. Research regarding organic fertilizers, liming, and the deficiencies of micro-elements has been much less extensive but has suffered from the same weaknesses mentioned above, hence has had minimal impact on the farmers.

The introduction of agricultural machines has had some impact in the basin; such machines include tractors, pumps, and small hand tractors with attachments. Tractors are typically owned by entrepreneurs who rent their machines and services to farmers on a contract basis per unit of land plowed (often to be paid after harvest); tractors are also used for local transport of agricultural produce, shelling corn, and many other farmyard tasks which significantly reduce the labor of the individual farmer. Pumps and tractors are more typically owned by the individual farmer, and they often accompany more intensive use of the land (2 or even 3 crops a year are being obtained in some areas). Improved hand tools have also been developed at some of the research stations, and these will be discussed in more detail in the following chapter. However, it seems that relatively few examples of new equipment adopted by farmers have resulted from research in the basin.

Research into new techniques of forest production has been conducted for many years, concentrating mostly on teak, new species for pulp (bamboo, kenaf), and development of plywood and fiberboard. Some research on reforestation has been conducted, but it is still in its infancy and must greatly expand if it is to meet current and future needs. Productive research can be carried out on fast-growing species for firewood and charcoal, watershed aspects of forest conservation, rotation of trees with crops, and sustained-yield forestry.

Insufficient research attention has been given to the control of pests. Rodent control, for example, has been conducted mostly as it relates to rice, both growing and stored, but little reduction in rates of damage has been obtained. The control of insect pests has centered on the use of chemical pesticides (often stimulated by the large chemical manufacturing firms), but research in other parts of the world is clearly indicating the drawbacks of the abusive use of such chemicals (especially since they often enter human food chains). Research on new chemical pesticides will certainly continue outside the basin, with research within the basin possibly concentrated not on trying to duplicate outside research, but taking advantage of it by finding the most suitable formulas and minimal applications required for specific local conditions. Biological control, i.e., using beneficial attributes of nature against her detrimental aspects, is receiving greater research emphasis in many countries and could be essential to the agricultural development of the basin.

Livestock research is another generally neglected field, possibly because milk and beef have not been important in the traditional diet of most basin residents. While the wild cattle of the basin (gaur, banteng, kouprey) are massive, handsome beasts, domestic cattle tend to be rather scrawny, with little meat or milk (being used mostly as draft animals). Private investment in ranching in Thailand has not, so far, been profitable because of difficulty in

developing an appropriate diet for breeds which are optimal for basin conditions; marketing has also been a serious problem. However, much of the basin area seems appropriate for grazing, so research and development of suitable grasses and legumes should be greatly expanded in the future. The possibility of using native deer and wild cattle should be carefully considered, since these species are already well adapted to local conditions and may require minimal care and investment.

Applying Research to Local Conditions

We include in this category all the programs of work on the selection of local varieties, adapting the results of research to local needs and the improvement of local techniques. Excellent results have been obtained with rice: a broad genetic diversity together with good adaptability to a wide range of environmental conditions has made it possible to select from local populations some broadly suitable and hardy varieties with very satisfactory yield potential. These varieties now occupy the great majority of the paddylands of the basin. One of the most striking examples is the Niaw Sanpatong variety, selected in the 1950s in Chiangmai province (northern Thailand) and now by far the most common glutinous rice variety in north and northeast Thailand and in the Vientiane plain of Laos. It has excellent grain qualities and moderate resistance to drought, flood, pests, and diseases; its flowering in October just at the end of the rainy season assures a minimum yield even when late rains result in late transplanting in July or even in August. The potential yield of this variety is quite high and can almost reach the hybrid yields of 6 t/ha with comparable doses of fertilizer (80-40-40 N/P/K kg/ha), and with appropriate cultivation techniques.

Improvement by selection alone still has obvious limitations, and the introduction of IR varieties has enabled successful hybridization programs to start, once the reasons for previous failures have been understood. If from the beginning researchers in Thailand have endeavored to adapt the IR varieties to local tastes, it is only after comparing the new RD hybrids with the local varieties that the qualities of the latter have been understood and appreciated. As a result, some of the characteristics of the local varieties are now being tentatively incorporated into the new hybrid generation. Even better, current research in the field of drought and flood resistance suggests the possibility of creating short varieties well adapted to variable conditions, in particular those of northeast Thailand. The work on floating rice in Bangkok has shown the existence of genes for elongation between nodes, enabling the plant to adapt itself to the level of the flood (Yantasast *et al.*, 1970).

Some work – still very modest – on mineral fertilization also belongs in this category. The objective is to determine the best usage of small doses of fertilizers, for example 10 to 20 kg/ha of nitrogen. This differs greatly from the still common approach of seeking the maximum yield or the economic optimum (maximum benefit). The research yield is the highest level obtainable with available technology (and this is the level approached by agriculture in the industrialized countries), but this is not realistic in the basin, since the farmers' limited credit, high prices of fertilizers, high interest rates for loans (if such are available), insufficient control of pests, and lack of sufficient water control prevent farmers from approaching the "research yield".

Work now being done in Thailand on maize and rice seems to show in both cases a profitable response to low doses of fertilizers when applied primarily as a top-dressing. Firms in the northeast typically apply ammophos (16-20-0 N/P/K) as a top dressing of 50 to 100 kg/ha on fields which appear to need it most, and this may be the most profitable input of fertilizer suiting the existing constraints. In the case of maize, practically no fertilizer is used; the profitability of fertilizers as measured by previous work is even lower than for rice and the response is unpredictable because of variable rainfall. The rather high risk in using fertilizers can be minimized by applying low doses as a top-dressing only.

Economic Impact of Agricultural Research

The brief survey of agricultural research in the basin outlined above might indicate a trend toward pessimism in that much of the work has not resulted in practical application and the overall agricultural productivity of the lower basin has not been improved much by the research. Accordingly the Mekong Secretariat has undertaken a study on the economic impact of research in Thailand. Two case studies, one on rice and one on maize, are being completed. In both cases, because of difficulties in obtaining precise statistical data and data on the adoption of recommended techniques by farmers, two separate approaches are being carried out in order to cross-check the results. The first approach consists of evaluating, for the main research directions, the yield increase due to the recommended technology and the cropping area on which this technology has been adopted. The second approach is an attempt to explain the evolution of provincial or national yields over a period of years, taking into consideration all the factors which could have had an effect on the yields, including irrigation, decrease in cultivated area per family, and changes of the fertility of the soil under continuous cropping. The net increase in yield after subtracting the effect of these factors is attributed to research.

It is concluded that the effect of research on rice production has thus far been relatively weak. Varietal improvement, including dry season crop and developments made possible by the new varieties, have led to a yield increase of 7.5 percent and fertilizers have increased yields by 12.2 percent. The other factors (cultural techniques, control of pests and disease, etc.) play only a marginal role or are practically impossible to evaluate. The economic impact, on the contrary, is far from being negligible. The Mekong Secretariat has calculated this economic impact by evaluating the internal rate of return for research in Thailand, based on budget allocations for research and experimentation of the Ministry of Agriculture compared to benefits to the nation (Mekong Secretariat, Information Notes Mkg 50/Mkg 51, 1977). For example, the internal rate of return for research on RD hybrids is over 20 percent even though the impact on production has been only 2.5 percent. On the whole, the internal rate of return for research on rice in Thailand carried out from 1955 to 1971 amounts to almost 47 percent, which is considerable. Similar conclusions were reached on maize.

It can be concluded, then, that research has been a profitable activity in spite of all of its insufficiencies. However, this is neither wholly reassuring nor satisfactory. The first steps are usually the easiest and the most profitable ones; such is the case with varietal selection

for example, while breeding involves much greater effort and is rarely followed by rapid success. Research can thus be expected to become increasingly costly, and will be justified only if its impact increases significantly. The ample potential for increased impact is suggested by the yawning gap between yields in the research stations and on the farmer's fields. We therefore cannot consider as satisfactory the present achievements of research which produced only some 15 to 20 percent production increase in 20 years; this is considerably less than the increase in population.

Although extension services have been usually lacking, the rapid diffusion of some innovations, with no or limited government involvement, such as cassava and certain new varieties of maize and of rice, demonstrates circuits of diffusion other than extension. Hence the reason many innovations are not adopted is due primarily to the nature of these innovations.

Research at the Scale of the Basin

The present organization of research has two basic levels. One level is represented by the international research institutes (here mainly IRRI in the Philippines), well-equipped and attracting some of the world's best experts in the relevant fields. The second level is represented by an assembling of research stations at the regional or even provincial level, covering many types of work, but not forming a structured network. Our purpose here is to suggest some organizational principles and to ask which tasks should be assigned to which level.

Role of International Research

The "Green Revolution", which was incited by the international research institutes such as IRRI, has been celebrated in the world press as the answer to world hunger. The predicted world famine, it was said, would certainly be prevented by miracle rice and miracle wheat. But when the green revolution failed to live up to its full potential, it similarly received exaggerated criticisms from many planners and social scientists. As Janzen (1972) has shown, with the introduction of high-yielding hybrid varieties, the local strains and practices which are closely adapted to local farming conditions, are quickly abandoned (especially when low-interest, long-term loans are made available by international development agencies). These high-yielding varieties are strongly dependent on pesticides, fertilizers, and complex breeding programs to keep control of the pest problem in single strain monocultures, leading to increased work and risk in the production process. Further, the new varieties themselves may have some problems. Intense selection for higher yields and other energy and nutrient-consuming traits, may force the plant to reduce its defense outlay in order to balance its internal resource budget. Thus "miracle grains" may be especially susceptible to insects and disease for internal reasons, as well as for reasons of genetic and botanical uniformity. Further research may eventually overcome these problems.

The fall from grace of the “green revolution” is comparable to the height which it attained during its heyday. But IRRI and other international institutions are neither so low nor so high; even taking into account recent modifications to the program, the results still appear quite promising. The concentration of intellectual, financial, and technical means at the international research centers can certainly produce radical and productive innovation, but it must be realized that these innovations cannot have universal impacts and must be carefully adapted to local reality.

Problems of Research Within a Region

In addition to glaring insufficiencies in staff and budget, one of the main characteristics of research at the local level is that it tends to be exceedingly narrow and the results are valid only for the conditions within the station. Interviews with researchers in north and northeast Thailand have revealed a surprising lack of knowledge by the researchers of the actual techniques used by farmers in their region. This myopic perspective also affects communication with basic or applied research carried out elsewhere, even within the same region; even the international literature (which is considerable) is very often ignored. This splendid isolation permits a comfortable work routine which can hardly be expected to be effective. The presence of foreign experts unfamiliar with the region and too often preoccupied with the interests of their parent organization or country cannot improve this situation very much.

As a consequence, an intermediate level has evolved, that of the national institutes or national research programs. These have been given increasing importance, are staffed with competent national and foreign researchers, and work closely with the international institutes. Their field of activity, however, is still limited (2 or 3 crops at the most) and their efficiency has been also limited due to lack of a strong anchor in the rural environment through the local stations. It is clear that the weakest level is at the base (the local stations) and this is the level which should be reinforced as a first priority, as is now the case in Thailand.

Towards a Network of Local Stations

One station for each broad ecological zone, say a dozen for the whole lower basin, seems sufficient, if these stations are well equipped and in close contact with the farmers and with other stations, both within the basin and internationally. The various types of soils within the zone could be covered through “outreach programs”.

Organizing these stations in a network should result in higher productivity at a lower cost. The network can be extremely versatile and could be started at any time in any of the countries. The first step consists simply of organizing periodic meetings of researchers and, if possible, putting at their disposition certain common services, however modest they might be. As an example, consider the impact that one statistician (bio-metrician) might have on northeast Thailand; adequately equipped and assisted, he or she would be

able to analyze all the trials carried out in this region, although this analysis will be fruitful only if communication exists between the statistician and the researcher. The statistician can help the researcher to better formulate his problem, to translate it into an appropriate experimental design, and to obtain a quicker answer to the problem which is being investigated (the report of the FAO project, “Coordination of Plant Production Research in Thailand”, gives many examples of the methodological weaknesses that could be thus avoided).

In many ways, the Lower Mekong Basin seems to be a geographical unit which could form a sound framework for such an integrated research program. Despite some environmental differences, the common traits are sufficiently numerous to treat the basin as a unit. This is well attested by the similar behavior of numerous crop varieties. We have already mentioned the case of Sanpatong rice, which is grown in Laos as well as north and northeastern Thailand. Another example is the PBS variety of maize. Also, the Thai variety of soybean gave the best results among the varieties tested in 1973 at Prek Leap (Cambodia) and Hegari is still the recommended variety of sorghum in all four countries. This list is very incomplete, but without doubt coordinated systematic trials within the basin would confirm the trend. Finally, we should note that the similarity between the major river basins of southern and eastern Asia gives good reason to expect that the results obtained in one basin would find large areas of application in the other basins as well.

What Type of Research?

The role of applied research (the only type considered here) is not to create varieties or techniques capable of producing 10 t/ha of rice, but to allow the farmers to go from 1-2 t/ha to 3-4 t/ha (and perhaps ultimately to 6 t/ha in a few localities). The reference to the farmer, his conditions and techniques of work, and the constraints of the physical environment are then fundamental, so research clearly must relate intimately to the local environment.

The studies of the agricultural environment, including detailed studies of a village or a small natural region, gathering geographical background information (soils, relief, vegetation, hydrology, climate) and sociological data (history of occupation land use, modes of land tenancy, sources of agriculture and non-agricultural revenue, distribution of the surplus, the agricultural calendar, etc.), would form a solid base for the type of research which would be most appropriate. The goal would be to understand the rationality of present agricultural techniques, and to understand their efficiency in a given regional context. Such understanding will also lead to the realization that any action on a single part of the system will eventually affect the whole system. Admittedly the system is rarely in stable equilibrium; hence we must also understand its dynamics and internal pressures so that we can identify the constraints which will have to be overcome to ensure its harmonious evolution. It may be difficult to incorporate such studies in a research program at a station, but they could be carried out by students of faculties of agriculture as research theses for higher degrees.

The necessity for another type of study, which is complementary and apparently easier, might be better understood by the planner; it consists of categorizing and comparing different types of farms. The grouping of farms into categories calls for sound understanding of the ongoing agricultural practices because the first step is to choose relevant criteria which allow placing the farms into homogeneous classes (these criteria might include cropping area, farm tenancy, main crops, main cropping systems, or any combination of these). Comparing mean economic results of the different classes may yield valuable indicators and guidelines, the more so when it is possible for a farmer to progress from a less productive to a more productive farming system. Perhaps even more fruitful is the analysis of results within one category of farm: which are the most productive individual farms, what techniques do they use, what cropping systems do they use and what constraints prevent others from obtaining the same results? It can be readily seen how this exercise can lead to the formulation of research programs.

Testing of the proposals resulting from research by appropriate in-station trials represents only the first step in relating the research to the local environmental conditions. The second step consists of outreach programs on various soil types. This second is often neglected for the third step: a series of replicated trials in the farmers' fields. The importance of a series of trials both as a research and an extension tool is being better understood by local researchers but such trials are still very rare (except the work done by the FAO soil fertility project). But in case of interaction between treatments and soils, the omission of step two will seriously affect the efficiency of step three: interpretation of discordant unreplicated trials is quite hazardous, and their demonstrative effect is obviously lowered. Hence when time is limited it is preferable to shorten or even to cancel the first step and directly test the treatments within various environments.

However, certain techniques cannot be reliably tested through this type of field trial. In the case of rotations, the techniques of conserving fertility, and mixed farming, it might be necessary to envisage another stage, which will be discussed below.

Research on New Farming Systems

Rarely does the farmer grow only one crop and as soon as he grows several crops they will start interacting (if only in arranging the working calendar). Moreover, intensification generally leads to more complex systems through accelerated rotation and overlapping of crops, multiple cropping, and crop association.

Therefore it is not sufficient to work on each individual crop as if it were always the main crop, benefiting from the best of care and being planted at the best time to give the best yield, etc. Research must also be conducted on cropping systems, and in analyzing each crop in relation to the other crops in the system, including looking for varieties that can be grown under suboptimal conditions and working on the methods of economical land preparation which would permit quickly and cheaply planted secondary crops. This perspective should also be applied to evaluating effects on the fertility of the soil (long-term trials of various rotations) and on the economics of the system (which might, for

example, reveal a more efficient use of fertilizers under multiple cropping where fertilizer application on a single crop is not economic).

If in the first stage these studies must be conducted on the station, considerable new difficulties and risks arise in applying the results directly to the farmers of the region. The complexity of such trials forbids the casual use of farmers' fields which, in addition, can hardly be expected to lead to a sound economic analysis. Since our thinking is directed at the farm level it is at this level that the salient trials should be conducted; we must have farms which are the focus of research operated by suitably motivated farmers who will implement the proposed solutions, and which are under permanent survey for observing the agronomic and economic results (inputs, working time, yield, etc.). It is only under such conditions that coherent techniques which are compatible among themselves and acceptable to the farmers can be elaborated.

Another significant aspect of this perspective is the maximum yield of a given crop which could be obtained through the optimization of all production techniques. Such optimization is clearly not possible in the farmers' fields, but when a single factor is being studied (such as planting density), all the others are ignored, i.e., put at their optimum level (fertilization, planting date, disease control, etc.). The notion of a "package of techniques" is now very familiar, but it is generally impossible to convince farmers to accept the whole package. Given the near total absence of factorial trials involving all the components of the package, any simplification is very risky and therefore no logical strategy can be proposed to extension services. Even after years of experimentation, it is still not possible to quantify the increase in yield at the farm level that would be achieved by the adaptation of each component, or each combination of two or more components. It should also be noted that such quantification is fundamental for planning, especially with simulation techniques.

In conclusion, after more than two decades of agricultural research, many problems have been solved and numerous successes have been achieved, but while impressive, these results can hardly be considered as sufficient. On the technical side alone, some rather elementary questions remain unanswered because the work has focused on crops rather than cropping systems. Even more important has been lack of communication between researchers and the farmers. Herds of experts have studied means to organize the transfer of knowledge from one to the other, but without notable success.

We have attempted to show that the problem must be posed in a different way, at a different level. Even the best extension, backed by strong rural institutions, can never convince farmers to accept innovations over the long term which do not really benefit them. Research has much to contribute, but it will be able to respond to the coming challenge only if it is radically converted along the lines we have suggested above. Further, constant evaluation and self-criticism, not with reference to external or exotic models, but to the reality of the farmer, will ensure that the research remains relevant.

The best way to communicate research results to farmers is through an intermediate level, where a working model is designed to deal with the actual physical, economic, environmental, and organizational problems likely to be encountered in the implementation

of full-scale agricultural development projects. The Mekong Committee has established a system of such models in the Lower Mekong Basin, called “Pioneer Agriculture Projects”.

Pioneer Agriculture Projects

The Mekong Committee’s vision of developing the water resources of the Lower Mekong Basin includes the irrigation of extensive areas of land. Although the basin farmers have for centuries practiced forms of agriculture based upon sophisticated utilization of the monsoon rains, they have very limited experience with modern irrigation. Such experience as they have indicates that they encounter the usual problem in developing new irrigation projects: unexpected and often long delays in the realization of the irrigation benefits. To help solve this problem the pioneer agriculture projects have been designed to remove many of the constraints which are preventing the effective implementation of irrigation. Under this program, one or more pioneer projects has been established in each of the riparian countries, including representation of all the major soil types and cropping patterns found in the lower basin (see Chapter 10).

In view of the close relationship between irrigation and dry farming and in order to provide a basis for judging the increase in agricultural production that can be expected from irrigation, the program also includes some projects in the rainfed (dry farming) agricultural areas. Such projects will also provide a broader basis for planning an efficient utilization of the basin’s agricultural resources.

History of Development of Pioneer Projects

The Bacon and White reports in the 1960s set the stage for the pioneer projects, but several more years of continuing efforts were required before the pioneer projects finally became established. In July 1967, a joint committee of representatives from the Governments of Thailand and Laos, together with technicians from the US Bureau of Reclamation (USBR) Pa Mong team and the Mekong Secretariat, reviewed research requirements. One of the relevant conclusions of this committee reads: “The Committee urges that a pilot irrigation scheme, in the magnitude of 10,000-20,000 ha, be centered in one of the newly constructed irrigation systems in northeast Thailand to consider all aspects of irrigated agriculture. Attention should be given to such key factors as farm size, land clearing and development methods and costs, marketing, farm credit, farmer inputs in crop production cycle, potential of developing agri-business activities, and farmer education processes; further recommends that development channels of governmental, quasi-governmental and commercial nature be exploited to achieve maximum success in the shortest possible time”.

At its 34th session in January 1968, the Mekong Committee approved a statement concerning the general aims of large-scale pioneer projects, which was endorsed by the Committee’s Advisory Board and the representative of FAO. In the Committee’s Annual Report for 1968, pioneer projects were defined as follows: “A pioneer project is the first

sector of development in a large-scale irrigation scheme. Several thousand families are involved in each such project; these pioneer families should receive whatever assistance is necessary. The main function of a pioneer project is to build up the necessary organization for the efficient management and maintenance of the larger developments that are to follow”.

The selection of this first pioneer project in northeast Thailand was made in January 1969, at a meeting held at Khon Kaen, under the chairmanship of Dr Boonrod Binson, Member of the Mekong Committee for Thailand, with representatives from all government agencies concerned as well as the Mekong Secretariat. The project selected at this meeting (Nong Wai) has since been prepared by the Technical Assistance Department of the Asian Development Bank. The general terms of reference of the project cover four main fields:

- a) **Irrigation:** How the headgate units should be organized, including the construction of farm distributaries and farm roads, and the establishment of watering schedules. Land clearing, land development and consolidation of farm units will also be addressed.
- b) **Technical and Scientific Aspects of Agriculture:** Concerning research, how the University of Khon Kaen and the Tha Phra central research station, as well as other experimental stations in the area, would need to be strengthened in order to meet the requirements of the project, particularly in the field of irrigated agriculture.
- c) **The Farmer:** Concerning farmers’ assistance, how the farmers should be assisted and how to increase labor productivity by means of mechanization. This would also include assistance in the field of credit, crop selection, fertilizer use, insecticides, and irrigation practices.
- d) **Marketing:** How to stabilize markets, in particular for crops other than rice.

The meeting recognized that the project should also consider organization (how the Government Departments should be involved in the undertaking, where gaps appear to exist in their present strength, and how the institutions should be strengthened) and resettlement (how any resettlement areas could be improved with a view to improved planning of future resettlement).

At its 38th session in January 1969, the Mekong Committee supported the proposals for the Nong Wai project and decided to sponsor, in addition, one such project in Cambodia (5,000 ha of the Prek Thnot scheme), and one in Lao PDR (5,000 ha of the Casier Sud).

At the 38th session, Sir Arthur Gaitskell, Member of the Advisory Board, strongly supported the proposals and agreed that such pioneer irrigation schemes were essential if the right decisions were to be taken on agricultural development in the region. These pioneer projects would have the primary task of determining programs, size of farm units, farmers’ organizations, and related matters. Unless such pioneer projects could point the way beforehand, the production needed to repay the heavy capital investment might be seriously delayed. He agreed to the use of the term “pilot”, because these projects would truly be breaking new ground and at a fairly large scale. However, in instituting such projects

before completion of adequate research into the technical and economic suitability of crops, the wrong advice might be given to farmers and this could discredit their confidence in the project. He suggested that careful consideration be given to the definition of project objectives, and to the role of private enterprise. Sir Arthur at the same time urged that “the need to strengthen and develop the research work of the experimental and demonstration farms be kept firmly in mind”.

A review of the Mekong program undertaken by the World Bank in 1970 upon the request of the Secretary General of the United Nations, led to the conclusion, among others, that the Committee’s pioneer agricultural programming was an undertaking worthy of being pursued and that a good many advantages might be obtained from a concerted program for pioneer projects in the context of Mekong development, instead of by a country-by-country approach. Advantages included:

- Greater momentum of development can be attained and with more efficient means, such as better training programs, transfer of experience to all four riparian countries, more efficient use of development assistance (e.g., personnel and contracts), and fewer constraints on participation by riparian country personnel.
- Greater appeal to the four riparian countries and effective interest in the individual pilot projects, particularly when associated with development of large scale projects.
- Greater appeal of the regional approach to those cooperating donor countries who are interested in agricultural development of the Mekong and who have expressed on many occasions the priority they attach to pilot projects.

Encouraged by the initial discussion with the World Bank, the Mekong Committee formulated the scope and objectives of a comprehensive Pioneer Agriculture Projects program at its 50th session in January-February 1971. The Committee then submitted a request to UNDP in July 1971, for assistance in the detailed preparation of the agriculture pioneer project program for the Lower Mekong Basin. A US\$ 2 million fund was created for this purpose to which UNDP contributed half, while the United Kingdom, United States, Netherlands, and Japan together provided the other half. The Plan of Operation for this assistance program was signed in September 1971 with the World Bank as Executing Agency, in association with FAO. The duration of the program of preparation was estimated to be about 2 years. Subsequently, it was agreed that part of the studies provided for in the Plan of Operation would be delegated to the ADB with the World Bank retaining the overall responsibility for the program. The agreement provided for participation by ADB in all consultations relevant to the program.

The pioneer projects program, as it has evolved from two years of subsequent studies, by 1977 included 11 projects in addition to projects prepared earlier (Nong Wai in northeast Thailand by ADB). In the seventh progress report of the Executing Agency, the planned pioneer projects program included:

- Three projects to provide pump irrigation facilities for river bank areas in the Lao PDR, northeast Thailand and Cambodia. The projects would explore different infrastructural requirements for the intensification and diversification of land use which

could eventually be applied to an estimated 500,000 ha of fertile levee soils on the Mekong and its tributaries.

- An irrigation rehabilitation project in Cambodia which would restore and improve a gravity irrigation scheme of about 30,000 ha which uses the unregulated flow of a tributary of the Great Lake. The project would focus on changes in systems operation and agricultural regime which would make more efficient utilization of unregulated river flow. It would also provide valuable information about water management and maintenance problems.
- The first stage development of a storage irrigation project in Cambodia would aim at an efficient transition from rainfed subsistence agriculture to intensive diversified land use with the support of year-round irrigation. The 5,000 ha project would establish different degrees of water control at the field level in order to assess the attendant differences in costs and benefits as well as in implementation, operation and maintenance problems. The pioneer project experience would have applications to the design of a full scale gravity irrigation project serving eventually 70,000 ha, and would also contribute essential information to the planning of other irrigation schemes in the Lower Mekong Basin.
- A project which would develop staff, facilities and organizational framework for systematic identification, adaptation, testing and dissemination of cultivation practices and seek technology suited for the improvement of rainfed rice cultivation in northeast Thailand. The project would broaden the activities of two government rice stations in northeast Thailand so that they could function as regional rice development centers with the support of regional rice breeding facilities.
- A pioneer project addressed to the identification, testing and implementation of viable farming systems for sustained rainfed cropping of upland soils in northeast Thailand is currently being prepared in close cooperation with the Directorate General of Agriculture in the Ministry of Agriculture and Cooperatives of Thailand.
- A project in northeast Thailand which would improve irrigated agriculture in the command area of three small reservoirs, giving special consideration to the establishment of project organizations under local control with farmer participation in operation and maintenance of these minor irrigation works. If a viable mode of development for these three reservoirs or “tanks” can be evolved, it could form a model for a large number of the existing tanks in northeast Thailand and might open up the possibility of further developments of this type in other parts of the Mekong basin.
- A project in the coastal zone of the Mekong delta which would establish an integrated salinity control system for a 50,000 ha area. The project would take maximum advantage of existing embankments and salinity control structures. Within this system, the pioneer project would develop modes of system operation to improve surface drainage in the wet season and to extend the period in which freshwater is available in the canals for irrigation by means of low-lift pumps. The project would also improve the agricultural

supporting services in order to enable farmers to take maximum advantage of the changes in the hydrological regime. The project would contribute to the detailed formulation of larger scale projects designed to protect and improve an estimated 1 million ha in the saline coastal belt of the Mekong delta.

- A project in the flood zone of the Mekong delta which would promote low-lift pumping from existing drainage and irrigation canals to introduce dry-season cropping in a 30,000 ha area which is at present primarily devoted to floating rice cultivation. The project would test irrigation techniques and on-farm developments likely to be relevant for the intensification of land use in the flood zone of the delta, which comprises about 300,000 ha.
- A project for year-round irrigation in an area of the delta free from flooding but subject to some salinity intrusion. Irrigation supplies for the 5,000 ha pioneer project would be obtained from a central pumping plant, located on an arm of the Mekong upstream from the location the fresh water-saline water interface during the low flow period. The project would experiment with the design of gravity irrigation systems for delta topography and land tenure and ownership patterns. It is a prerequisite for the detailed preparation of pending larger irrigation projects for as much as 250,000 ha.

Full Project Development

Once research has been applied through the pioneer projects and has been successfully adapted to local conditions, the farmers of the basin are expected to respond quickly to their opportunities. The key to expanding the pioneer projects to other parts of the basin irrigation systems lies in accurately identifying the relative benefit of these projects and conclusively demonstrating these benefits to other farmers (as well as to governments and investors).

It is still much too early to give a meaningful account of project development, but even at this early stage some remarks can be made regarding several major problems encountered. In essence, these problems are related to farm management. We know now that agricultural benefits will in general have a slower start than is assumed in the feasibility reports, with a longer time-lag between implementation and full production level. This applies especially to areas where soil conditions are poor and present farm income level is low. The longer time-lag needed for implementation, the rather uncertain final production levels at the full development stage, and the varying periods of project life combine to account for a rather unreliable estimate of the farm benefits. As a consequence, economic comparison between the various projects should be regarded with the utmost caution.

The planning of production levels and cropping patterns is very difficult and will almost always be inaccurate. Very often projects are judged only on their “bankability”, so that loans can be obtained for implementation. However, once a government wishes to choose among various projects, comparisons between projects must be made. This is possible even with inaccurate production estimates, as long as a consistent method of estimating the benefits is applied; but this has seldom been the case.

A first attempt was therefore made by the Mekong Secretariat to outline a uniform approach for the estimation of farm benefits, which so far is limited to the full development stage. Work is also in progress to include an assessment of the time-lag needed to reach the full production stage. Basically, the methodology for estimating farm benefits is very simple. Crop requirements and preferences or tolerances are evaluated against physical factors prevailing in the field. The latter may prohibit the cultivation of a crop or variety, reduce the yield, have no effect on the yield, or increase the yield.

Although the method is simple in theory, this system nevertheless becomes complicated because of the variety of factors involved. Under normal field conditions many factors cause reductions in yield. The influence of any one such factor is usually studied by keeping all factors constant except the one under investigation. But even where the relationship between the yield of a crop and each relevant factor is known, it is still impossible to predict the yield because of interdependency between the factors. Yield-influencing factors are seldom independent of each other, but neither are all factors interdependent. In fact, groups or clusters of closely-related factors are formed, with each cluster having a more or less independent impact on the yield. Within a cluster, the most limiting factor governs the effect on yield.

Three clusters of yield-influencing factors are distinguished for purposes of model evaluation:

- a. Drought:** The main factors concerned are rainfall evapotranspiration, water-holding capacity of the root zone (dry-foot crops), and depth of ponding and percolation (paddy).
- b. Excessive Water:** The main factors concerned are rainfall, storage capacity, drainage capacity, deep percolation, and flooding. Both drought and excessive water clusters are variable from year to year, though some factors (e.g., rainfall) vary more than others (e.g., evaporation).
- c. Soil Properties:** The main factors concerned are texture, organic matter content, air capacity, cation exchange capacity (CEC), pH, soil depth, and soil permeability. These properties are usually more or less independent of farming practices, though a high level of farm management may improve many of them.

Additional factors are also taken into account as absolute constraints, such as the existence of a paddy layer, ESP (exchangeable sodium percentage), relief, humidity, depth of ground water table, and secondary and micro-nutrients. The macro-nutrients, nitrogen, phosphate, and potash, are treated separately since their utility is largely governed by economic considerations.

A productivity rating is calculated for each cluster in accordance with the degree to which the crop requirement and preferences or tolerances are satisfied. Each cluster receives a rating from 1.0 (no yield reduction) to 0.1, but in practice the productivity ratings vary from 0.4 to 1.0. The cluster ratings are then multiplied, resulting in the relative productivity level that can be expected from a given land unit.

Yields and Fertilizers

The relation between yields and fertilizer inputs is very complex, involving many factors. For planning, it is necessary to formulate as precisely as possible how yields and import requirements are estimated in order to be able to compare different projects. Classes of yields distinguished for the purposes of analysis include: theoretically possible, potential, maximum, obtainable, and actual.

Theoretically possible yield is limited by incoming radiant energy, temperature, relative humidity, and day length. It has been calculated that, at latitudes of 10 to 20 degrees, the average theoretically possible yields of modern rice varieties vary between 11 and 13 tons per hectare when the sky is clear throughout the growing period and between 5 and 6 tons per hectare when the sky is continuously overcast (van Ittersum, 1972). In view of the normal cloudiness in the Mekong basin, this would roughly mean an average theoretically possible yield of about 10 to 12 tons per hectare in the dry season and 8 to 9 tons per hectare in the wet season. Such yields are never reached, mostly due to inadequate disease and pest control, especially in the later stages of the growing period when discoloring of the leaves renders photosynthesis less efficient.

Potential yield (or the approximate research yield) is limited by the stage of technology and structural organization in agriculture. Four different stages of development have been distinguished regarding rice cultivation in Japan, which are compared with the present condition in various countries in Asia (Takase and Kanoh, 1975):

- a. Primitive farming with yields of about one ton per hectare or less (now widespread in the basin);
- b. Irrigated cultivation, where yields increase to 2.5 tons per ha;
- c. Technical innovations, with yields increasing to about 4 tons per ha;
- d. Structural reform, with yields increasing to about 6 tons per ha, while possibly 8 tons per ha can be reached.

Comparisons with Japan's rice-growing history suggest that it will be quite an achievement by the Mekong farmer to reach a yield of half the theoretically possible yield, or 5 to 6 tons per ha in the dry season and 4 to 5 tons per ha in the wet season. These yields are taken to be the potential yields for modern rice varieties in the present study, and are considered realistically obtainable.

The maximum yield is derived from the potential yield by multiplying the productivity rates for drought, excessive water, and soil properties by the potential yield. This varies with each crop and land unit.

The obtainable yield is derived from the maximum yield by taking the supply of macronutrients (N/P/K) into account. Crop responses to fertilizer applications require a lot of experimental work, but since such work is often lacking in the basin, experimental data from outside the area have been used as basic data for the calculations.

The actual **yield** is the average farm yield in a project area. While this yield is directly measured in the field, it can be theoretically related to the maximum obtainable yield discussed above. The actual yield equals the maximum yield if the limiting constraints is one or more physical factors and not the supply of macro-nutrients. It equals the obtainable yield without fertilizer applications when such applications do not show economic benefits; and it equals the obtainable yield with fertilizer applications up to the point where the marginal costs equal the marginal benefits. It is less than the obtainable yield when fertilizers (or suitable credit for buying fertilizers) are in short supply.

The most economical fertilizer application and corresponding yield are calculated from crop response curves, showing the relationship between yield increases and additions of fertilizers. These response curves are mostly derived from results obtained from demonstrations on farmers' fields, and are complemented by specific data from research trials, including requirements for pesticides.

The analysis of the variables discussed above, as well as data on labor requirements and the alternative use of animal or mechanical power, is carried out by means of three main independent, consecutive computer programs, in which the output of the first program is necessary as input for the next, with some additional data.

Program I consists of a main computer program ("RATINGS") in which the relative yields of individual crops are determined for the different land units recognized in a project area, and two auxiliary programs for calculating the rainfall exceedence frequencies ("REXFREX") and the drought ratings ("DROUGHT") by means of a water balance model.

Program II ("YIELDS") translates the relative yields from program I into actual yields which are or can be obtained under specific conditions. Prices play an important role in determining the most economic yield.

Program III ("LP") contains both agronomic and economic aspects and determines, by means of linear programming, optimal farmer behavior (i.e., cropping patterns and production techniques) under different stages of development: the present situation; the projected future without the development project; and the full development model with the project.

The best choice for a farmer, tentatively taken to be maximum income (within the limits of the farmer's capability to invest and take risks), can be calculated with this model, taking into consideration the farmer's resources in terms of land, labor, and working capital. These resources are obviously quite variable in a project area, so that three farm size groups are distinguished: small (less than 2.5 ha); medium (2.5-4.0 ha); and large (over 4 ha). The estimation of accrued farm benefits can be standardized once the period required for attaining full project development is established.

Such a standardized project evaluation system has many advantages. Comparisons can be made between different projects; basic data collecting can be more efficient and precise, enabling a more detailed description of the work, and the production increases can be

calculated from the realistic point of view of the producers.

While the actual decisions of the basin farmers will be based on their own perceptions of their alternatives, such computer modeling can help project planners to make better decisions, which are then more likely to be accepted by farmers.

Conclusion

The agricultural development model adopted by the Mekong Committee consists of three major components. First, research on new and improved crops and techniques is conducted at experimental and demonstration stations. Second, the results of this research are applied to a full-scale project for testing under practical conditions in Pioneer Agriculture Projects, which are essentially proposals for development strategies. Third, the new agricultural systems which have been shown to be successful on a pioneer level will be applied to appropriate areas in the Lower Mekong Basin. The first two steps have already begun, with results presented in the following two chapters.

Chapter 9

Pioneer Agriculture Projects

Introduction

In order to adapt the findings from the experimental stations and demonstration areas to the existing conditions in the various parts of the basin, a network of eleven pioneer agricultural projects was planned by the Mekong Committee, adapted to the major soil types and cropping patterns in the lower Mekong basin. Lessons learned from these projects can thus be applied to areas of considerably greater extent than the pioneer projects themselves.

While the pioneer projects are highly diverse, they have several common themes. Each project provides for project monitoring and studies of the technical, organizational, agronomic and social aspects of the project, with particular emphasis on the response of the farmers to the project. Findings of the monitoring programmes are used for solving problems encountered in project implementation and serve as a guide for the planning and implementation of additional projects and project extensions.

All pioneer projects give strong emphasis to farmer's organizations. While the exact form and structure of these may vary from country to country, all are based on giving the local people the key roles in implementing the projects. Consultants and high levels of government inputs are typically limited in time since all pioneer projects are designed to have a relatively short life as such.

It should be emphasized that each of the pioneer projects was defined in terms of short-term solutions to immediate needs, based on benefit and cost considerations, current availability of fertilizers and current prices of agricultural products. An individual project is therefore not necessarily related to the long-term needs and the long-term potential of the basin, which may be considerably higher than envisioned by a more constrained study. The long-term potential will be reached only as the experimental and demonstration stations determine optimal strategies for development and as these strategies become widespread in the basin.

The proposals for pioneer projects have been prepared by a wide range of agricultural consulting firms. The Asian Development Bank and the World Bank have reviewed these projects, making some modifications. The project synopses presented in this chapter are derived from the Bank review memoranda, which were in turn based on the consultants' reports. More detailed bibliographic references are presented in illustration 12.

ILLUSTRATION 12: SELECTED CONSULTANTS' REPORTS ON VARIOUS PIONEER PROJECTS	
Muk Kampul:	The Muk Kampul Pioneer Agricultural Project in the Khmer Republic (now Cambodia), Vol. I, II. Nov. 1972/Feb. 1973; Summary Report, Hay, 1973. AGRAR/SATEC/ILACO.
Bovel	The Pioneer Agricultural Project. Oct. 1973. Vol. I, II. ILACO/AHT/SATEC.
Casier Sud	Reports on the Vientiane Plain Pump Irrigation Pioneer Projects. Vol. I, II. Aug. 1972. FAO/IBRD. Feasibility Report on Casier Sud Pioneer Agricultural Project, Vol. I-III. Aug. 1973. Japan Engineering Consultant Co.
Se Bang Fai	Draft Report of the Rural Development Pioneer Project July, 1974. FAO/IBRD.
Pump Irrigation	Studies of Pioneer Pump Irrigation Project, Northeast Thailand, Vol. I-VI. Dec. 1973. Lyon Assoc.
Tank Irrigation	Pioneer Project for Tank Irrigation in Northeast Thailand Vol. I, II. W. Halcrow/Urwick Lugg/Gould.
Rainfed Crop Improvement Nong Wai	Nong Wai Pioneer Project for Irrigated Agriculture. Vol. III. June 1971. AHT/ILACo.
Go Cong	Go Cong Pioneer Agricultural Project. Vol. 1.111. Sept. 1973. ADC.
Tiep Nhut	Final Progress report on the Tiep Nhut Pioneer. Agricultural Project. Sept. 1973. Nippon Koei.
Caisan	Caisan Pioneer Project. Sept. 1973. Hunting/Sir M. McDonald and Partners

For each of the pioneer projects, we have discussed the essential elements in two major categories: background (the project area, soils, and socioeconomic factors); and the project (objectives, project works, cost estimates, organization and management, agricultural production, and benefits and justification). Most of the pioneer projects are concerned with irrigation, and the cost of these varies considerably with the ecological conditions and the type of irrigation to be attempted; hence the costs indicated in the project synopses are not fully comparable since they include different development items. Cost data are included for indicative purposes only, showing total costs of about US\$ 83.2 million for the eleven projects.

Muk Kampul Pump Irrigation Pioneer Project (Cambodia)

Background

Project Area. The project area comprises a narrow strip of land about 17 km long and averaging 3 km wide, situated on the west bank of the Mekong river just north of Phnom Penh. It has an area of 4,500 ha, of which 2,200 ha are irrigable. Land forms and land use, which are typical of some 10,000 ha of land along major rivers in Cambodia, include

- a. The bourrelet – the top of the natural river levees occupied by villages, orchards, and vegetable gardens.
- b. The chamcars – the land sloping inland from the levees, subjected to annual flooding from the Mekong; the upper parts are free of flooding for a sufficient period each year to allow double cropping (the lower areas can only be single cropped).
- c. Ricelands and bungs – the low-lying areas flooded during the rainy season which are devoted to rice grown on the receding flood; some areas are irrigated from shallow ponds called “bungs”.

The project area contains some 220 ha of bourrelet, 660 ha of double cropped chamcars, 300 ha of single-cropped chamcars, and 1020 ha of lowland rice. Cropping intensity is about 200 per cent on the bourrelet, 160 per cent for the double-cropped chamcars, 100 per cent on the single-cropped chamcars, and 100 per cent for the lowland rice-growing area.

On the chamcars the land holdings consist of narrow strips of land extending down the slope, varying in width from under 5 m to over 20 m. Many of the strips are subdivided laterally among several owners and a typical farm holding consists of numerous plots scattered throughout the strips. Increases in production of upland crops and rice would be possible if irrigation water could be provided, but the land tenure pattern, presents a major problem in the design and layout of a conventional canal irrigation system.

Soils. Soil types are alluvial on the levees and clays in the lower areas where the annual floods deposit fine sediments.

Socioeconomic factors. The population of the project area is about 14,000, divided into 2,700 family units of which some 1,600 are farming families (the high percentage of non-farming families is due to the proximity of Phnom Penh and is not representative of most riverain areas). Farm holdings average about 1.5 ha, usually consisting of four or five separate plots. Land rental or leasing usually concerns only separate plots and not whole farms. Garden and fruit plots on the bourrelet do not appear to be leased, while some 9 per cent of the paddy area and 17 per cent of the chamcar area are under rental agreements.

The Project

Objectives. The main objective of the project is to evolve and demonstrate solutions to the technical and managerial problems which have prevented the development of viable pump irrigation on the riverain lands adjacent to the Mekong river. On the technical side, the aim is to develop a water distribution system for the existing land tenure conditions which would be economically and socially workable. On the managerial side, emphasis is on securing closer cooperation between the various agencies providing services to the farmers and developing organizations for collective pumping under local control

The main difficulty in formulating the project has been the design of a distribution system compatible with the land tenure pattern. Not only is the land farmed in narrow strips, but the ownership also changes along the strip. However, providing water in a channel along the top of the chamcar would give considerable incentive for the farmers to cooperate in water distribution in various ways by means of earth channels, portable pipes, or flumes. Given technical advice and assistance in planning the water distribution layouts, it should be possible to evolve workable systems extending 200 to 300 m down the chamcar, to encompass most of the double-cropped chamcar area.

Project Works. The pioneer project would provide pumping units, irrigation distribution facilities for some 730 ha, and improved agricultural services in the project area. Project works include 17 pumping units, each with three 15 hp diesel-powered pumps mounted on rails to allow them to be raised and lowered according to water levels in the Mekong. Main channels will be constructed to take water to the edge of the bourrelet zone running parallel to the river, across the bourrelet, under the highway, and along the top edge of the chamcar zone, again running parallel to the river. The minor distribution systems would be evolved during project implementation and would include channels, earth channels, and possibly portable flumes and pipes.

On the average, each unit would irrigate some 5 ha of high value crops on the bourrelet, about 20-30 ha within the double-cropping chamcar zone and about 10-15 ha of ricelands where the farm plots are wide enough to accommodate a delivery channel through the chamcar. The project also includes buildings, workshops and vehicles for government supporting services, technical assistance for project implementation and the preparation of future riverain pump irrigation projects, village electrification, and project monitoring.

Total project costs are estimated at US\$ 206 million, of which about half is in foreign exchange (illustration 13). The cost estimates assume free access for goods supplied to the project area and do not reflect currently prevailing transportation conditions and costs.

ILLUSTRATION 13: COST SUMMARY, MUK KAMPUL (in US\$ 000)			
Item	Local	Foreign	Total
Pumping units	60	220	280
Irrigation distribution system	490	130	620
Buildings	80	20	100
Vehicles	0	100	100
Engineering and consultants	100	200	300
Surveys and mapping	40	30	70
Village electrification	50	150	200
Base cost estimate	820	850	1,670
Physical contingencies	110	70	180
Price contingencies	330	320	650
Total	1,260	1,240	2,500
Project monitoring	30	20	50
Studies for future development	10	40	50
Total project cost	1,300	1,300	2,600

Organization and Management. One of the major aspects of this project is to evolve a suitable and efficient management structure for the type of riverain area development to be pioneered. Firm proposals for organization and management will require discussions with the Government of Cambodia. For the present, the establishment of a Project Implementation Unit (PIU) within the Ministry of Agriculture is suggested. The PIU would be staffed by the necessary personnel to survey and map the topography and soils of the project area; to plan and design the irrigation system in cooperation with the project farmers; to organize the construction of the works; and to coordinate the supply of agricultural credit, services, and physical inputs in the pump area and to supply extension services to the farmers.

The operation and maintenance of the pumps would initially be the responsibility of the PIU, but this responsibility would be passed to the village group served by an individual pumping unit as soon as the group has been organized and a village based pump operator has been trained. Operation and maintenance of the irrigation distribution system will be the responsibility of the farmers, organized into water user groups. Each group would be headed by a common irrigator who would be responsible for determining water releases for the group and for collecting pumping charges, operation and maintenance fees assessments, and any other necessary fees.

Agricultural Production. As a result of the project, the cultivation of high value vegetables during both wet and dry seasons in the irrigated part of the bourrelet is estimated to increase from 25 ha to 80 ha, with the average yield rising from 7.7 tons per ha to 10 tons per ha. Irrigation of the chamcars (450 ha) would lead to increased yields of vegetables and other crops and to an increase in cropping intensity. Paddy yields on the 200 ha of irrigated ricelands are estimated to increase from 2.0 tons per ha for local varieties to 4.2 tons per ha for high yielding varieties. Estimated annual crop production at present, and for the with-project case, are shown in illustration 14.

ILLUSTRATION 14: ESTIMATED ANNUAL PRODUCTION FOR IRRIGATED AREAS OF MUK KAMPUL PIONEER PROJECT			
Crop	Unit	Present	Future with Project
Bourrelet (80 ha)			
Vegetables	tons	350	1,600
Fruit or other	tons	550	-
Chamcar			
Vegetables	tons	550	1,260
Maize - cobs	cobs (,000)	-	8,400
- grains	tons	-	700
Sesame	tons	42	95
Greengram	tons	41	67
Rice Lands (200 ha)			
Paddy - Local	tons	400	-
- HYV	tons	-	840

Benefits and Justification. The project would evolve and demonstrate technical and managerial approaches to the development of pump irrigation on the riverain lands adjacent to the Mekong river, and other similar areas along major rivers in Cambodia.

The economic rate of return was calculated using farmgate prices derived from projections of 1985 world market prices expressed in 1974 constant terms. Project costs used in the analysis covers pumps and distribution systems of the 17 Muk Kampul units and one-half of the cost of other items (which would serve a wider area), amounting to a cost of US 1,100 per ha for the 730 ha which it is conservatively assumed would benefit from the proposed 17 units. With these assumptions, a 13 per cent economic rate of return is indicated. Present and projected production values for the project are summarized in illustration 15.

ILLUSTRATION 15: MUK KAMPUL PIONEER IRRIGATION PROJECT
Present and Projected Production Values

	Cropped Area (ha) (W.S.+D.S.)¹	Yield (ton/ha) (US\$/ton)	Farm-gate Price Production	Gross Value of (US\$/ha)	Production Costs² Production Area	Net Value of Project thousand)	Net return from (US\$
Present Production							
Bourrelet							
Vegetables	50 (25+25)	7	120	840	82	758	37.9
Other ³	110 (55+55)	5	120	600	50	550	60.5
Sub-total	160 (80+80)						98.4
Cropping Intensity (%)	200						
Double-Cropped Chamcar							
Vegetables	110 (30+80)	5	120	600	56	544	59.8
Maize (cobs) ⁴	350 (220+130)	17	8	136	22	114	39.9
Sesame	140 (100+40)	0.3	250	75	14	61	8.5
Greengram	120 (80+40)	0.34	250	85	24	61	7.3
Sub-total	720 (430+290)						115.5
Cropping Intensity (%)	160						
Riceland							
Local Rice Varieties	200 (0+200)	2	110	220	19	201	40.2
Cropping Intensity (%)	100						
Total Present Return							254.1
Bourrelet							
Vegetables	160 (80+80)	10	120	1,200	168	1,032	165.1
Cropping Intensity (%)	200						
Double-Cropped Chamcar							
Vegetables	140 (40+100)	9	120	1,080	162	918	128.5
Maize (cobs)	240 (150+90)	35	8	280	132	148	35.5
Maize (grain)	200 (100+100)	3.5	70	245	134	111	22.2
Sesame	135 (90+45)	0.7	250	175	72	103	13.9
Greengram	95 (50+45)	0.7	250	175	81	94	8.9
Sub-total	810 (430+380)						209
Cropping Intensity (%)	180						
Ricelands							
Local Rice Varieties	25 (0+25)	3	110	330	76	254	6.4
HYV Wet Season	25 (25+0)	3.5	110	385	69	316	7.9
HYV Dry Season	175 (0+175)	4.3	110	473	135	338	59.2
Sub-total	225 (25+200)						73.5
Cropping Intensity (%)	113						
TOTAL PROJECT RETURN							447.6

¹ (Wet Season + Dry Season)

² All production costs exclude payments for labour.

³ Comprising annual fruits, lesser value vegetables resulting from inefficient irrigation.

⁴ Maize (cobs) yield and farm-gate price are expressed in a unit of 1,000 cobs
(i.e. 1,000 cobs/ha and US\$/1,000 cobs).

Bovel Gravity Irrigation Pioneer Project: (Cambodia)

Background

Project Area. This is a run-of-river irrigation project in Battambang Province on the right bank of the Mongkol Borey river, a tributary of the Great Lake. The existing works of the Bovel project include a diversion dam on the Hongkol Borey, a natural channel serving as the main canal, 61 km of secondary canals, and 15 km of tertiary canals. The works were constructed to divert the waters of the Mongkol Borey over an area of about 46,000 ha for irrigation of a wet-season crop. The river has no storage and the divertible flows show wide seasonal variation. Dry season flows are very low but in the high rainfall months the river rises sharply and overflows its banks, flooding the area to depths of about 50 cm. Mid-season droughts in July and August are common, and the project was planned to provide supplementary irrigation to the rice crop during these drought periods.

Most of the area is planted to long-straw local rice variety, which are broadcast in May and June and harvested in December and January. The area receiving supplementary irrigation is limited to narrow strips of land bordering the canals, in total about 10,000–15,000 ha. Even in these areas the water supply is unreliable, because in its present form the system is difficult to operate and maintain. The chief deficiencies are the lack of tertiary canals to distribute water from the main and secondary canals and inadequacies in the design and construction of the canals and control structures.

Soils. The soils in the area are heavy clay, including 90 per cent brown hydromorphics and 10 per cent lacustrine alluvials.

Socio-economic factors. About 8,500 families (68,500 people) are engaged in agriculture on the area of 46,000 ha. At one time most of the land was farmed as rice plantations, but these have been broken into smaller units over the years. Still, about 30 per cent of the area is taken up by large blocks of over 100 ha (in 1974; this has certainly changed now), with these larger holdings being leased or rented under various arrangements in units of about 5 ha. Landlords with large holdings typically farmed only about one-third of their land. Probably more than 80 per cent of the project areas is farmed in units of 5 ha or less.

The Project

Objectives. The primary objective of the project is to develop techniques for effective utilization of the waters of the Mongkol Borey by improvements in canal systems and drains an area of over 14,000 ha. The experience gained in engineering, operation, and maintenance, the strengthening of agricultural supporting services, and experimentation with different rice varieties and crop calendars, would apply to future development in the Bovel Project area and to other potential developments in Cambodia.

Project Works. The project would include construction of a 10 km primary canal, a 5 km secondary canal, and 25 km of main and secondary drains; the rehabilitation of 25 km of existing secondary canals; and the construction of tertiary canals and drains for an area of about 14,000 ha. Consultant services would also be supplied to assist the government in the design and construction of the irrigation works, the establishment of operation and maintenance procedures, and technical advice in setting up improved agricultural supporting services. Vehicles, equipment, and buildings for project implementation and management would also be supplied.

A net area of 40,000 ha would be developed, with a proposed cropping intensity of 125 per cent. Crop varieties and crop calendars would be selected to match the water requirements as far as possible to the seasonal variation in water supply. Long-duration rice varieties (180 days) would be planted in May on 10,000 ha; a short-duration rice variety (120 days) would be planted on 10,000 ha of the area in July and on 20,000 ha early in August. Half the area planted in August would also be used to grow an early season crop of mung beans, maize, or peanuts

This level of agricultural development would require much better water control than currently exists, so irrigation systems servicing blocks of about 100 ha will be constructed over a period of 12 years. The pioneer project would be the first phase of an overall development plan comprising a 4,000 ha unit with intensive development. It would also include some major features of the ultimate project plan, such as sections of the main and secondary canals and drains.

Realization of the proposed cropping pattern would depend on the ability of the proposed drainage works to reduce flooding. Rice planted in August, as opposed to the current practice of planting in May and June, would be much more susceptible to damage. In September or October from inundation due to heavy rains and overbank flow from the rivers; the technical and economic feasibility of preventing seasonal inundation of the project area has yet to be established. Uncertainty on this account does not, however, preclude a phased program of canal system improvements and drain construction which, in conjunction with improved agricultural practices, could lead to higher yields for the wet-season rice crop. Drainage works constructed in the early stages would be designed to allow for future extension and modification when the drainage problem is better understood. To develop such an understanding, a data collection programme would be carried out as a part of the pioneer project.

Cost Estimates. Total project costs are estimated at US\$ 8.1 million of which the foreign exchange component is estimated at US\$3.7 million. Project costs are summarized in illustration 16.

ILLUSTRATION 16: COST ESTIMATES, BOVEL PROJECT			
	Local	Foreign US\$ (thousand)	Total
Main, secondary canals and drains	1,240	950	2,190
Tertiary canals and drains	880	520	1,400
Roads and bridges	160	201	80
Buildings	130	201	50
Vehicles and equipment	0	150	150
Surveys and mapping	20	30	50
Government supporting services	150	0	150
Consultants:			
Project implementation	80	340	420
Further studies	60	240	300
Base cost estimates	2,720	2,270	4,990
Physical contingencies	400	350	750
Price contingencies	1,200	1,080	2,280
Project monitoring	60	20	80
Total project cost	4,380	3,720	8,100

Organization and Management. One of the major pioneering aspects of this project will be to evolve a suitable and efficient management structure for the implementation and operation of project works and for agricultural supporting services. Firm proposals for organization and management of the project will require discussions with the government of Cambodia, but for the present the establishment of a Project Implementation Unit (PIU) is suggested, with similar duties to those of the PIU suggested for Muk Kampul.

The operation and maintenance of the diversion dam, main canals, and secondary canals, would initially be the responsibility of the PIU, while operation and maintenance of the tertiary systems will be the responsibility of the farmers, organized into water user groups. The PIU would encourage the formation of these groups and initially supervise their activities. Each water user group would be headed by a common irrigator who would be responsible for determining water releases for the group and for collecting operation and maintenance assessments and any other necessary fees. Steps would be taken by the government to impose and collect water charges to cover the costs of operation and maintenance and a reasonable proportion of capital costs. This charge would be imposed within about one year of the first delivery of a reliable water supply.

Agricultural Production. As a result of improved water control and more effective agricultural supporting services, local variety wet season paddy yields on the 14,000 ha project are expected to increase from 1.4 ton per hectare to around 2.5 tons per hectare at full project development. No significant increase in yields is likely without the project;

with the project, rice yields are expected to improve due to better water control and the use of a small amount of cash inputs and some improved farming practices, resulting from the sustained extension effort under the project. Total paddy production at full project development is estimated to increase from 20,000 tons without the project to 35,000 tons, while net farm income from a 4 ha farm is estimated to increase from some US\$ 330 at present to around US\$ 600 at full development.

Benefits and Justification. The proposed project is justified on its own merits, and as a pioneer project demonstrating technical and organizational approaches applicable to other run-of-river irrigation projects in Cambodia. The economic rate of return is estimated to be 12 per cent, based on conservative assumptions for cropping patterns and crop calendars. No account was taken of the possible introduction of shorter-duration rice varieties which would allow early or late season crops of maize, mung beans, groundnuts, and others; such developments would greatly increase the rate of return. The net return of the project is projected to be about US\$ 2.45 million per year, as opposed to US\$ 1.33 million per year without the project (illustration 17).

ILLUSTRATION 17: BOVEL PIONEER IRRIGATION PROJECT Present and Projected Production Values		
	<u>Wet Season</u> Present Production	<u>Wet Season</u> Project Production at Full Development
	Local Variety Paddy	Irrigated Local Variety Paddy
Cropped Area ha	14,000	14,000
Yield (ton/ha)	1.3	2.5
Farm-gate Price US\$/ton	100	100
Gross Value of Production	130	250
Production Costs ¹ US\$/ha	35	75
Net Value of Production	95	175
Net Return from Project Area US\$ million	1.33	2.45

¹ All production costs exclude payments for labour

Casier Sud Pump Irrigation Pioneer Project (Lao PDR)

Background

Project Area. The project area comprises 5,100 ha, of which 3,170 ha are cultivable, lying on the left bank within a large loop of the Mekong river about 7 km downstream

from Vientiane. This area is strongly affected by floods, with serious damage occurring as often as every two years. Even during years when overbank flooding from the Mekong is not severe, its relative high levels often significantly impede surface drainage from local rainfall, resulting in crop damage. During the dry season, although discharges in the Mekong are plentiful, water levels drop to as much as 10 meters below the project area, too low to permit use for irrigation. In the absence of other sources of water, crops in the project area often suffer from drought.

Present cropping in the area is predominantly rainfed wet season glutinous rice, with significant amounts of green maize, tobacco, sugar cane, fruits, and vegetables grown during both wet and dry seasons, albeit with recurrent damages from flood and drought. Cropping intensity is estimated at 112 per cent. Small amounts of sericulture and livestock production are also practiced, with the nearby and easily accessible Vientiane area providing a ready market for the area's produce.

Soils. The area contains primarily clayey flood plain soils and lighter, more free-draining natural levee soils; both groups of soils are potentially quite productive assuming flooding can be controlled.

Socio-economic Factors. The present population in the project area is estimated to be about 23,000 of which some 17,500 are agriculturally employed or supported. The project area has an estimated 2,580 agricultural households, each of which operates a farm averaging about 1.5 ha. The land tenure situation is unclear and requires further investigation.

The Project

Objectives. The aim of the project is to increase agricultural production from an area of about 3,170 ha currently subject to prolonged wet season flooding from the adjacent Mekong river or from impeded local drainage, and also subject to dry season drought conditions. These conditions combine to significantly limit agricultural output from an otherwise potentially productive area. As a pioneer project, it will serve to demonstrate viable solutions to these physical problems which are typical for large areas of agricultural lands along both banks of the Mekong river upstream of the delta.

The proposed project would include construction of a flood control dike along the left bank of the Mekong, protecting an area of some 5,000 ha from overbank flooding; construction of four irrigation pumping plants; the required associated electrical transmission facilities; construction of an integrated irrigation and drainage system and farm road network; rehabilitation of the existing Hat Dok Keo experimental station; and provision of consulting services for project design and construction supervision and for assistance in initiating agricultural development programmes under the project.

The project diking scheme provides a ten-year level of flood protection as giving adequate yet economically justifiable protection for the project's predominantly agricultural areas, without adversely affecting flood conditions upstream or on the opposite bank of the river. The project proposal includes provision for controlled egress into the project area of flows in excess of this level of protection, to eliminate catastrophic damages which would otherwise be expected from overtopping or breaking of the levee.

Project Works. The project includes four irrigation pumping plants spaced along the project periphery, each serving 600-1,000 ha, rather than a single large plant serving the entire 3,170 ha project area. This proposal is justified by system operation and maintenance considerations and the difficulty of achieving cooperation among large groups of farmers. While multiple pumping stations would be preferable to a large single station in this case, it will still be necessary to examine use of simpler and less costly types of pumps better suited to local conditions, in place of the vertical shaft submerged pump facilities currently planned.

For disposal of surface drainage from inside the project area during periods when gravity drainage is not possible because of high water levels in the Mekong, a drainage pumping facility is proposed. This would have capacity to evacuate the drainage flood within a three-day period while limiting areas affected by inundation to 1,000 ha; damaging inundation is not expected to exceed 180 ha in a normal year.

Cost Estimates. Total project costs are estimated at US\$ 9.0 million, including a foreign exchange component of US\$ 5.3 million. Project costs are summarized in illustration 18.

ILLUSTRATION 18: COST SUMMARY OF CASIER SUD PIONEER PROJECT			
Item	Cost (US\$ equivalent)		
	Total	Local	Foreign
Land acquisition	359	359	-
Construction			
Preparatory works	90	49	41
Dike	1,183	624	559
Pumping Plants and transmission lines	1,096	144	952
Irrigation and drainage system	1,374	554	820
On-farm works	100	100	-
Agricultural facilities	137	91	46
Other			
Consulting services, surveys and mapping	936	164	772
Operation and maintenance equipment	130	-	130
Administration	216	216	-
Basic cost estimate	5,621	2,301	3,320
Physical contingencies	1,124	460	664
Price contingencies	2,226	911	1,315
Project monitoring	70	50	50
Total Project cost	9,041	3,722	5,319

Organization and Management. The project will be implemented under the Ministry of Economy and Planning. During project implementation farmers' cooperative organizations will be set up to provide for cooperative marketing, credit and input supply, and farm mechanization efforts in association with the project. Concurrently, water users' associations will also be formed to assist with system operation, maintenance, and collection of water fees.

Agricultural production. The primary direct effect of the project will be increased agricultural production. The improved water control and availability of necessary inputs, credit, and agricultural supporting services will permit widespread use of high-yielding varieties and large-scale double cropping. Cropping intensities in the project-area are expected to increase from present levels of 112 per cent to 190 per cent with the project. Under the project, paddy yields are expected to increase from 0.9 to 4.0 tons per ha in the wet season and from 3.0 to 5.0 tons per ha in the dry season; corresponding increases in yields would take place for other products.

Benefits and justification. The project would directly benefit some 2,600 farm-families, raising net farm income of a typical 1.5 ha operation very substantially from present levels of US\$ 119. The project would, under full development, produce an incremental 12,000 tons of rice and some 9,000 tons of various other crops yearly. All these products are currently in short supply in the Vientiane area and are being imported; thus in addition to directly enhancing farm incomes, the project would contribute substantially to improvement of the country's balance of payments position.

The project's economic internal rate of return is estimated at about 11 per cent, considering only direct crop production benefits and using a 25-year evaluation period.

Se Bang Fai Flood Protection and Pump Irrigation Pioneer Project (Lao PDR)

Background

Project Area. In the Thakhek, Savannakhet, and Pakse provinces, as in the rest of the Laotian part of the Mekong basin, agricultural production is mainly concentrated in the flood plains of the Mekong tributaries such as the Se Bang Fai, which forms the boundary between the two provinces of Savannakhet and Khammouane. The Se Bang Fai flood plain extends from both banks of the river, over some 30,000 ha; the plain is some 15 km wide and is bounded by the Mekong river to the west and by upland open forest to the east. It consists of low-lying basins separated by higher grounds at an elevation of a few metres above the basins. The Se Bang Fai river levees are high enough to prevent overtopping by rainy season flooding except at points or in years of exceptionally high floods.

Within the area, the road system consists of two dirt tracks on the river banks and of access tracks to the flood plains which are bad during the dry season and impassable during the rainy season.

Soils. The soils in the project area are mainly alluvial deposits whose characteristics are determined by their elevation (which affects texture, internal drainage, and leaching). Old alluvial terraces at higher elevation surrounding the flood basins are never flooded and are covered by open forest. These are shallow, acidic soils of very low fertility. Soils of the annually flooded plain have a fine texture in surface and sub-surface horizons which impedes drainage; because of their physical characteristics, these soils are suitable only for paddy if permanently flooded during the whole growing cycle. Levee soils (about 10 to 20 per cent of the flood basin area) along the river bank which are flooded only at peak water level have a coarser texture. They are well-drained because of their higher permeability and elevation; they are fertile and can be used for all upland crops.

Socio-economic Factors. In the flood plains of the southern provinces of Laos, most farms are of comparable size and characteristics. A typical farm includes some 2.5 ha of permanent rice fields and 1.0 ha of upland fields on levees and river banks where vegetable and tobacco are grown when the floods recede in the dry season. A typical farm keeps 4 to 5 head of cattle and buffalo, 2 to 3 pigs, and 10 to 20 chickens. The family usually is engaged in non-farming activities as well, such as fishing, weaving, and handicrafts.

While farm economy is still largely at subsistence level, the traditional standard of farming is high. In the low-lying areas, rice fields are leveled and bunded; buffaloes are used as draft animals with locally made implements; supplementary feed is supplied to working animals and household waste to pigs and poultry. Manure is used for river bank cultivation, but modern inputs are rarely applied because of high risks of crop failure in the flood plain.

Farmers live in villages which are the centres of all agricultural activity. The village headman and the elders elected by the village inhabitants play a prominent role in village life. Mutual help between families is common for rice cultivation and traditional farmer groups operate and maintain small gravity irrigation schemes built by the community as well as village rice mills. Such groups also contribute to the construction of flood protection or of small water storage works by providing voluntary unpaid labour.

The Project

Objectives. The Se Bang Fai flood plains appear quite suitable for a pioneer project, since soils, population density, farming techniques, cropping patterns, and crop returns are similar to those of most other Mekong tributaries above the reach of the Mekong flood, not to mention of many other small river basins scattered all over Laos. The first phase of the pioneer project would gradually improve traditional agriculture without modifying present cropping patterns; it would increase returns by avoiding periodic crop failures. Such project targets are tailored to the concepts and traditional activity and organization of the Lao farmer, and should therefore meet his full cooperation inasmuch as this does not imply intensive cultivation practices.

Over a period of five years the project would develop some 22,000 ha of rice land plus 8,000 ha of uplands and river bank levees, involving about 8,000 farms or 55,000 persons living in 86 villages. On the basis of hydrological data, the project would ensure partial control of flooding, entailing rice crop security 9 years out of 10 instead of 2 out of 3 as at present. In addition, small dams and diversion canals on three perennial streams would allow double cropping of rice on some 700 ha. On the levees, in addition to tobacco, maize would be grown for improved pig and egg production.

Project Works. Project works would include flood control works, involving the construction of compacted earth-filled concrete dams with flap-gates at the mouth of 17 of the Se Bang Fai tributaries, three earth-filled concrete dams with sluice gates on three perennial streams and 8 km of unlined main canals to small gravity irrigation perimeter ditches to be hand built by farmers; supply of 200 small individual pumps to water vegetables, tobacco, and maize on the upland and levee soils; supply of 7 village rice-mills of 1 ton per hour capacity; and rural and social infrastructure including 60 km of road, 100 village wells, and 6 village health dispensaries.

Other components of the project will include farmer training, extension services at a ratio of one field extension officer for 250 farmers; a small animal breeding unit to supply improved piglets and ducklings; technical assistance; and further studies to prepare a full flood control/irrigation second phase project over the whole project area.

Cost Estimates. Total project costs are estimated at US\$ 8.3 million, of which foreign exchange component is 4.7 million. Project cost estimates are presented in illustration 19.

Organization and Management. An important part of the pioneer project will be the progressive establishment of farmer groups, based on traditional village organizations, in order to cooperate in construction and maintenance of flood protection works and small supplementary irrigation networks, organizing marketing of surplus production, processing rice at village level, and ensuring distribution of improved piglets and breeding poultry to be supplied by a project breeding unit.

The project would be implemented by the provincial services of the Department of Agriculture in the two concerned provinces under the supervision of the Ministry of Agriculture and its specialized sections. Technical assistance provided under the project would include one rural engineer, two extension specialists, and a training specialist.

Agricultural Production. As stated previously, the project would increase production of glutinous rice under traditional farming practices by improving crop security. It would also introduce (on a minor scale) dry season rice and would upgrade pig and poultry production through progressive introduction of improved breeds.

In the low-lying rice fields, traditional rice varieties would continue to be grown, with no use of fertilizers and pesticides. Through better crop security, average yields would increase from 1 t/ha to 1.4 t/ha of paddy over some 22,000 ha; this is an increase of 9,000 tons/year. Total pig live weight would increase from 640 tons to 2,000 tons, while egg production

ILLUSTRATION 19: LAO PDR, SE BANG FAI PIONEER PROJECT Project Cost Estimates in 000 US\$			
Item	Quantity	Unit	Total
Flood Protection			
Cut-off dams	17	No.	535
Levee dykes	10	km	
Irrigation			
Derivation dams	3	No.	165
Main distribution canals	5	km	10
Communication Network			
Roads	60	km	600
Construction on dykes and dams	30	Km	240
Bridge	1	No.	10
Motor ferry	2	No.	70
Social Infrastructure			
Borehole's	100	No.	500
Dispensaries	6	No.	60
Ambulances	6	No.	60
Credit Requirements			
Rice-mills	7	No.	140
Individual pumps	200	No.	100
Ducks and pigs ('000)	40	No.	40
Extension and Training			
Extension officers	3	No.	30
Extension workers	32	No.	250
4-wheel drive vehicles	3	No.	15
Motorbikes ('000)	80	No.	24
Running costs of vehicles			100
Training and demonstration equipment			5
Project Management			
Managerial staff	10	No.	120
Clerks, accountants, drivers	30	No.	225
Rural engineers	3	No.	20
Draftsmen, topographers	20	No.	90
Office and houses including equipment			100
Trucks	4	No.	40
4-wheel drive vehicle	15	No.	15
Motor bikes ('000)	50	No.	150
Running costs of vehicles			
Technical Assistance			
Foreign technical staff	5		1,150
Housing	50		50
4-wheel drive vehicle	9	No.	45
Running. costs of vehicles			50
Animal breeding unit	1	No.	25
Further studies			200
Maintenance of works (item 1 to 4)			650
Total			7,214
15% Contingencies			1,100
GRAND TOTAL			8,314

would increase from 320,000 to 1,830,000/year. Net project incremental income would reach about US\$ 1.3 million annually at full development.

Benefits and Justification. The project would benefit some 8,000 families, solving their main agricultural problem (high risk of crop failure due to flooding and drought). Net farm incomes would increase by an average of 70 per cent. Added health facilities and village water supplies together with improved transport networks would improve living conditions in the area and ease marketing difficulties. Incremental rice and animal production would substitute for imports and entail foreign exchange savings of about US\$ 600,000 yearly. The economic rate of return is estimated at about 12 per cent.

Lam Pao Irrigation Improvement Pioneer Project (Thailand)

Background

Project Area. The Lam Pao project area is located in Kalasin province in the central part of northeast Thailand. Most of the main irrigation works have already been completed, including a large storage reservoir and main canals. The project area includes 4,500 to 5,600 ha, with a total of 50,000 ha available for wet-season irrigation and 25,000 ha available for dry season irrigation at full development. The local relief within the project area is varied; in the northern part, the ground falls from the main canal in a series of terraces to the river plain, but in the southern portion the land slopes more regularly at about 1 per cent from the laterals to the river. Areas of slightly higher ground are occupied by villages but these cover relatively small areas.

Soils. The area selected for the pioneer project lies mainly on the leached sandy loams which have been classified as suitable for both rice and upland crops, if provided with adequate irrigation and drainage facilities. These soils total about 85 per cent of the pioneer project area. The lower areas adjacent to the river have small areas of heavy clay soil covering approximately 15 per cent of the pioneer area; these soils are also suitable for rice cultivation. Both of the soil types are typical of the soils in the other drainage basins of the northeast where major irrigation schemes are sited.

Socio-economic Factors. About 1,500 farm families live in the project area in 15 villages; the average family size is about 6, including 2 to 3 adult farm workers. The cultivated area per farm is usually between 2 and 4 ha, virtually all of which is devoted to paddy cultivation on leveled and bunded plots. Plot sizes normally vary from about one-sixth to about one-half of a ha, depending on the land slope. On many of the farms, the plots held by individual farmers are continuous, although often small and irregular because of the need to collect and control rainfall. The problems arising from scattered and fragmented landholdings have not been fully evaluated; the inheritance system is such that a farm is taken over by only one offspring while the others clear and level new land elsewhere. The freehold

ownership of land (through a title deed) is rare although almost all the land is holder-operated. Most land is held under customary occupation and user-rights and is not negotiable for sale or for mortgage.

The Project

Objectives. The pioneer project is to be a prototype model for accelerating development of large areas in northeast Thailand which are currently served by major irrigation works which have been completed or are under construction. The main problem on these projects is the gap between the construction of major irrigation facilities and the utilization of water for farm development. The primary objective of the pioneer project is to show how this gap can best be bridged and to ensure the supply of irrigation water, in both the wet and dry season, from the existing main irrigation facilities to the individual farm unit within the service area of 40 ha (which comprises a farm turnout). The use of irrigation water in quantities and at the time required by the crops would by itself bring substantial increases in agricultural production in both seasons. Because of the general lack of firm information about such development in the area and the need to build new procedures, evaluation, and flexibility into the initial project, its costs will likely be substantially higher per hectare than those anticipated for the expansion based on its results. The increase in agricultural production from irrigation, however, would justify the project in strictly economic terms and the experience gained can be applied and repeated to other projects where major irrigation facilities already exist. The project would, in short, determine the extent to which irrigation and drainage facilities must be brought to the farmers' fields before the farmers themselves can be expected to operate and maintain the new field channels.

Project Works. The Lam Pao reservoir was filled in 1969; the main canal and four lateral canals are concrete-lined and were completed in 1970-1971. New project works will include the rehabilitation of that portion of the existing main canal and four laterals which extend through the project area; this would include the designing of the canals and the stabilization of the deep-cut and spoil banks lining the canals. Channel structures and additional outlets from the existing main canal and laterals will be provided, along with better cross-channel drainage on the laterals. Distribution canals and drains would be constructed to serve about 5,000 ha. The operation and maintenance equipment required for the project are estimated to comprise three small excavators, vehicles, tractors, and small tools. The project includes provision for improving existing roads in the area and levelling the tops of canals so that they are useful for transport.

Cost Estimates. The preliminary total cost of the pioneer project is estimated at US\$ 3.8 million, of which the foreign exchange component is about 28 per cent. Project costs are summarized in illustration 20.

Organization and Management. Since no public agency is as yet experienced in the type of integrated organization and management which the Lam Pao project would require, exclusive responsibility within the project area should be vested in an independent project

**ILLUSTRATION 20: SUMMARY OF COSTS FOR LAM PAO
PIONEER PROJECT**

	US\$ (thousand)		
	Local	Foreign Exchange	Total
Land acquisition	80	-	80
Offices, camps, housing	100	-	100
Project equipment	-	92	92
Construction	1,237	413	1,650
Consulting services	245	367	612
Farm credit	150	-	150
Maintenance equipment	-	116	116
Project staff	165	-	165
Field crop promotion	100	-	100
Sub-total	2,077	988	3,065
Contingencies 25 per cent	519	247	766
Total	2,596	1,235	3,831

management authority, which would be directly responsible to the authority designated by the Government of Thailand, perhaps in the Prime Minister's office. The provision of farm services would also be financed under the project, consisting of three categories of assistance to the farmers:

- a. A programme carried out to construct, operate, and maintain field channels through the service units would require that the designs, supervision of works, and guidance in operation and maintenance be given to the farmers by the project staff.
- b. Project agricultural officers would advise farmers on irrigated agricultural development and farm intensification. These officers would work in close cooperation with the credit services.
- c. A special programme would be financed for demonstrating efficient field crop production. Works under this programme will include the rationalization of field layouts and the installation of better field irrigation and drainage on a selected number of the larger farms in the area located on suitable soils. A small number of service units would be involved during the first year, with the number subsequently expanding as necessary.

A short and medium-term credit programme could be linked with the farm extension activities in the project area. In some cases, loan terms would need to be tailored to the special needs of the project. A special project credit fund would be made available through the auspices of the Bank of Agriculture and Agricultural Cooperatives.

Agricultural Production. The yields of rice, which is now grown under rainfed conditions only are expected to be doubled from the present level of under one t/ha to over two t/ha. It is believed that with a secure supply of water in the fields during the dry season, the farmers would begin to grow rice and field crops during that period. After three or four years, about half the farm area would be sown in the dry season.

Benefits and Justification. Given a reliable and adequate supply of irrigation water, the farmers should have a compelling income incentive to diversify their cropping patterns and adapt new inputs. On the average it appears that a farm family should be able to increase cash income at an annual rate of not less than 15 to 20 percent. The average gross value output of a hectare of farm land under these assumptions would almost quadruple. Labour returns would more than double, although most of the increased farm returns would, in fact, result from the more efficient use of labour through its better distribution over the year.

As the Lam Pao dam and main canal are already in existence, they have been treated as sunk costs and only supplementary expenditures required to bring the water from the main canals to the farms are included. The economic rate of return has been calculated on the basis of assumed net additional income generated per family within the context of the entire Lam Pao basin development, varying from 13 to 16 per cent, depending on the speed with which the irrigated innovations are applied.

Northeast Thailand Pump Irrigation Pioneer Project (Thailand)

Background

Project Area. The project covers three existing pump irrigation projects on the right bank of the Mekong river in Nong Khai, including Tha Bo (cultivable area 1,500 ha, irrigation area, 700 ha), Ban Mo (cultivable area 1,400 ha, irrigation area 1,000 ha), and Muang Mee (cultivable area 1,000 ha, irrigation area 500 ha).

Service areas of the projects consist of river levees, old river channels, and poorly drained swamp areas. The slope is generally away from the main river and poor drainage limits the area to which intensive irrigation can be introduced. With the exception of Muang Mee, the project areas are subject to flooding from the Mekong at about 10-year intervals. The existing irrigation systems require extensive rehabilitation and reconstruction. Except at Muang Mee, the electric pumps are operable but require overhauling. The canal systems, however, will largely need to be replaced, due to poor design and neglect of maintenance. The existing systems have no tertiary ditches, drains, or access roads. Agricultural supporting services are generally weak, with a ratio of extension worker farmers at least 1 to 4,000.

Soils. Soils are alluvial and are generally fairly heavy silty clays. About 10 per cent of the soils are suitable for upland crops, which include sugar cane, maize, tobacco, and vegetables.

Socio-economic Factors. About 2,000 farm families (14,000 persons) own land in the project area. The average farm size varies from 1.3 ha at Muang Nee to 3.6 ha at Tha Bo. Less than 10 per cent of the farmers are tenants. Less than 10 per cent of the cultivable area is cropped in the dry season, but some revenue is being collected from the farmers to cover part of the operating costs of the rainy-season irrigation system

The Project

Objectives. The main objective of the project is to demonstrate solutions to the technical and managerial problems that have prevented the full realization of benefits from the existing pump irrigation projects in northeast Thailand. On the technical side, the aim is to rehabilitate and improve the pumping and conveyance systems to a level permitting a high cropping intensity and the use of high-yielding rice varieties. This will require on-farm development in and the establishment of proper operation and maintenance procedures. On the managerial side, an attempt will be made to develop close cooperation between the various disciplines providing services to agriculture and to provide a transition between government control and eventual local control of the systems. Finding the best method of doing this will be one of the major pioneering aspects of the project.

Project works. The project includes the following works on the pumping and conveyance systems:

- a. **The Bo:** rehabilitation of the pumping station; intake protection works; replacement of delivery pipeline; reconstruction of the canal system, with service roads and telephone links; construction of on-farm and main drains; and minor leveling;
- b. **Ban Ho:** similar to Tha Bobut, but also includes re-siting of the pumping station;
- c. **Muan Mee:** similar to Tha Bo but also includes purchase of two turbine pumps and construction of a pumping station.

On-farm works will be provided to ensure proper water control at the farm level and adequate access to the farms. This would comprise a system of territory ditches and drains, spaced at about 150-250 m, with farm roads along 40 per cent of the ditches. In order to achieve a rational and economical layout, some boundary realignment would be required, together with a moderate amount of land levelling.

Cost Estimates. Total project costs are estimated at US\$ 2.9 million, of which US\$ 293,000 would be for project monitoring and further studies. The foreign exchange component is estimated at US\$ 1.5 million. Project costs are summarized in illustration 21.

Organization and Management. The Royal Irrigation Department (RID) would be responsible for implementing all civil works, including on-farm development. The RID would be assisted by a firm of consulting engineers in rehabilitation and improvement of existing works and in establishing operation and maintenance procedures.

**ILLUSTRATION 21: SUMMARY OF PROJECT COSTS OF THE
NORTHEAST THAILAND PUMP IRRIGATION PIONEER PROJECT**

Item	Local	Foreign	Total	Local	Foreign	Total
	Baht million			US\$ (thousand)		
Pumping stations	3.4	5.0	8.4	170	250	420
Irrigation and drainage	4.6	2.5	7.1	230	125	355
On-farm development	5.0	3.3	8.3	250	165	415
Miscellaneous items	3.5	2.8	6.3	175	140	315
Engineering and consultants	3.3	4.0	7.3	200	340	365
Sub-total	19.8	17.6	37.4	990	880	1,870
Physical contingencies	1.7	1.9	3.6	85	95	180
Price contingencies	5.4	6.0	11.4	270	300	570
Total	26.2	25.5	52.4	1,345	1,275	2,620
Project monitoring and evaluation	1.3	0.3	1.6	65	15	80
Studies for future projects	0.6	3.4	4.0	30	170	200
Grand total	28.8	29.2	58.0	1,440	1,460	2,900

The RID would be responsible, at least in the early years, for operating the pumps and primary canals, but operation and maintenance of the secondary and tertiary systems serving units of about 100 ha would be the responsibility of the farmers themselves, organized into water-user groups. Each group would be headed by a common irrigator who would be responsible for determining water releases within the service unit and for collecting water charges.

Improvements in the quality and intensity of extension in the three concerned areas would emphasize the training of leading project farmers to serve as extension workers. The Department of Agricultural Extension would appoint a Project Extension Officer for each command area to develop and manage a small agricultural unit. Selected farmers would be recruited and trained as extension personnel. Credit would either be provided by strengthening the cooperatives in the project area, or by stationing a credit agent in each pump area from the Bank for Agriculture and Agricultural Cooperatives, which would be the main institutional credit source. Multi-purpose cooperatives would also be established, as for the tank irrigation project.

Agricultural Production. As a result of the project, the cropping intensity of the 2,100 ha to receive irrigation would increase from 110 per cent at present to about 190 per cent. There would be a shift from broadcast to transplanted long-strawed, early-maturing native varieties in the lower-lying areas and high-yielding varieties would be introduced in the flood-free irrigated area. The area devoted to upland crops, such as sugarcane and tobacco, would remain small but cropping intensity would increase. Rice yields would improve due to greater water control and a greater use of cash inputs and improved farming practices,

resulting from the sustained extension effort under the project. Paddy yields would increase from the present 1.6 t/ha to about 3.3 t/ha for native varieties, and 4.7 t/ha for high-yielding varieties, at full project development. Net farm income for a 1 ha farm would be approximately doubled.

Benefits and Justification. Although the sub-projects are not ideal sites for pumped irrigation, due to flooding and drainage constraints, rehabilitation and completion of the existing projects, together with the provision of strong supporting services, would be justified. Successful pioneer project would provide a model for similar rehabilitation or new pump irrigation projects along the banks of the Mekong. The project would approximately double employment opportunities and farm income in an area of about 4,000 ha.

The economic rate of return was found to vary from 13 per cent for Muang Mee to 19 per cent for Ban Mo to 21 per cent for Tha Bo, with the whole project returning 18 per cent. It was found that projects of this kind should give an acceptable rate of return (15 per cent) with total investment costs, at 1974 prices, of up to about US\$ 1,400/ha. Present and projected production values are presented in illustration 22.

ILLUSTRATION 22: NORTHEAST THAILAND PIONEER PUMP IRRIGATION PROJECT
Present and Projected Production Values

	Cropped Area (ha)	Yield (ton/ha)	Farm-gate Price (Bht/ton)	Gross Value of Production	Production Costs ⁵ (Bht/ha)	Net Value of Production	Net Return from Project Area (Bht/million)
Wet Season (irrigated)	<u>Present Production</u>						
Broadcast paddy ⁶	250	1.1	2,200	2,400	600	1,800	0.5
Transplanted paddy ⁶	1,600	1.6	2,200	3,500	800	2,700	4.3
Others ⁷	350	-	-	5,000	2,000	3,000	1.1
Sub-total	2,200						5.9
Dry Season (irrigated)							
HYV paddy	160	2.8	2,200	6,100	1,700	4,400	0.7
Others ⁷	200	-	-	5,000	2,000	3,000	0.6
Sub-total	360	1.3					
Total	2,560						7.2
Cropping Intensity	116						
Wet Season (irrigated)	<u>Projected Production at Full Development</u>						
Transplanted paddy ⁶	1,100	3.2	2,200	7,000	2,000	5,000	5.5
HYV paddy	600	4.6	2,200	10,100	2,400	7,700	4.6
Others ⁷	500	-	-	6,000	2,500	3,500	1.8
Sub-total	2,200						11.9
Dry Season (irrigated)							
HYV paddy	1,700	4.7	2,200	10,300	2,500	7,800	13.3
Others ⁷	300	-	-	6,000	2,500	3,500	1.1
Sub-total	2,000						14.4
Total	4,200						26.3
Cropping Intensity (%)							

⁵ All Production costs exclude payments for labour

⁶ Local varieties

⁷ Comprising sugar cane, tobacco beans and vegetables

Northeast Thailand Tank Irrigation Pioneer Project (Thailand)

Background

Project Area. The project area consists of the command areas of three tanks, each taking the name of the stream on which it is constructed: Huai Chorakne Mak, Huai Talat, and Phok. The total drainage area is 1,040 km²; live storage capacity is 52 million m³; the area to be irrigated includes 3,440 ha in the wet season and 890 ha in the dry season.

The service areas consist of narrow strips of flood plains and low river terraces bordering the rivers on which the dams have been constructed. Most of the land has been cleared and is used for cultivation of a wet-season crop of long-duration transplanted native rice varieties. About 100 ha is used for upland crops in the dry season with irrigation from the canals or the rivers. Paddy yields are about 1.1 t/ha. The average farm size in the service areas is about 2.5 ha and the land holdings are often fragmented.

Each of the dams consists of a low earth embankment, about 5 m high, with a concrete spillway and reservoir outlet. The dams have deteriorated since construction, due to inadequate compaction and excess seepage, erosion and insufficient freeboard. In one case, extension of the spillway is required. The main canals, which total 93 km in length, have mostly been excavated to their design lengths, but due to neglect or maintenance, only one is operative for its full length. About 30 percent of the canals are lined, but the lining has deteriorated in many places.

Soils. Soils are low humic gleys derived from alluvial material; they are characterized by a high content of fine sand and silt in the upper horizons, with underlying horizons of hard impermeable silty clay.

The Project

Objectives. The aim of the project is to develop engineering and organizational solutions to the problems which have heretofore stood in the way of effective utilization of existing tanks in northeast Thailand. The project will include repairs and improvements to the dams and spillways; rehabilitation and improvement of about 80 km of primary canals; construction of secondary and tertiary canals and drains serving about 3,400 ha; improved standards of operation and maintenance; improvements in agricultural extension and other supporting services; and technical assistance in project implementation and the identification and preparation of similar projects for tank rehabilitation.

Following a review of 30 of the largest existing tanks, three tanks were selected because the records showed that they filled in most years and hence would provide a reliable supply of irrigation water.

Project Works. The three existing earth embankments forming the tanks would be repaired and improved to ensure their stability and safety from wave action and seepage. The crest level of the Lam Phok dam would be raised by 0.5 m to provide adequate freeboard. New gates would be installed at the Huai Talat spillway. At Lam Phok, the flood discharge capacity would be increased by construction of an additional spillway structure.

Of the 83 km of existing primary canals, some 27 km have been lined with concrete. In the lined sections, the damaged portions of the concrete lining would be repaired or replaced. The unlined sections, of which parts are badly deteriorated, would be restored to their design profiles and cross-section. Service roads with a 3.5 meter-wide laterite surface would be provided along the main canals and some of the larger secondary canals. The secondary canals and drains would be provided to serve blocks of about 100 ha, and tertiary canals and drains would be provided to serve units of about 5 ha. Land consolidation and land leveling would be carried out on about 100 ha in each tank service area, to provide experience for more intensive development in the future.

Cost Estimates. Total project costs are estimated at US\$ 3.9 million, of which the foreign exchange component is estimated at US\$ 1.8 million. Costs are summarized in illustration 23.

ILLUSTRATION 23: COST SUMMARY FOR THE TANK IRRIGATION PIONEER PROJECT						
Item	Local	Foreign	Total	Local	Foreign	Total
	(Bht.	Million)		(US\$ thousand)		
Dam rehabilitation						
Canals and drains:	4.6	3.2	7.8	230	160	390
Primary	6.6	4.2	10.8	330	210	540
Secondary	6.3	4.0	10.3	315	200	515
Tertiary	1.7	1.0	2.7	85	50	135
Buildings	1.5	0.5	2.0	75	25	100
Vehicles and equipment	0	1.5	1.5	0	75	75
Surveys and mapping	0.6	0.4	1.0	30	20	50
Consulting services	1.2	4.2	5.4	60	210	270
Engineering and Supervision	2.5	0	2.5	125	0	125
Sub-total	25.0	19.0	44.0	1,250	950	2,200
Physical contingencies	4.8	3.2	8.0	240	160	400
Price contingencies	10.2	7.8	18.0	510	390	900
Total	40.0	30.0	70.0	2,000	1,500	3,500
Studies for future projects	1.5	5.0	6.5	75	250	325
Project monitoring	1.1	0.4	1.5	55	20	75
Grand total	42.6	35.4	78.0	2,130	1,770	3,900

Organization and management. Following an organization pattern which has been adopted by the government for other irrigation projects, the Royal Irrigation Department (RID) would be responsible for implementing all civil works including on-farm development. The RID is planning to set up a special unit for on-farm development, whose staff could receive the necessary in-service training for the layout and design of the secondary and tertiary systems.

The RID would be responsible, at least in the early years, for operating the dams and primary canals. Operating and maintenance of the secondary and tertiary systems serving blocks of about 100 ha would be the responsibility of the farmers themselves, organized into water user groups. The RID would encourage the formation of these groups and supervise the standards of their work. Each group would be headed by a common irrigator who would be responsible for determining water releases within the service unit and for collecting water charges.

One of the major purposes of this project would be to develop organizations capable of managing small irrigation projects, with a minimum of assistance from the government.

The type of organization required would be a multipurpose cooperative which would eventually assume responsibility for operation and maintenance of all civil works including the dam as well as the provision of agricultural extension, credit, and marketing services. Arrangements would therefore be made with the Department of Cooperative Promotion to design the organizational framework of three multipurpose cooperatives for each tank, in close cooperation with the RID and the Department of Extension.

Agricultural Production. As a result of the project, the cropping intensity on the area to receive irrigation (3,400 ha) would increase to about 125 per cent. In the wet season, traditional varieties would gradually be replaced by high-yielding varieties on about 75 per cent of the irrigated areas. Dry season cropping would be divided almost equally between high-yielding rice, legumes (groundnuts and mung beans), and vegetables. Paddy yields would improve due to greater water control and a greater use of cash inputs and improved farming practices, resulting from the sustained extension effort under the project. Paddy yields are estimated to increase from 1.1 t/ha at present to 2.8 t/ha for traditional varieties and to about 4.2 t/ha for high yielding varieties. Vegetable yields would approximately double, to 9 t/ha, and legume yields would increase from 0.8 to about 1.3 t/ha. Paddy production would increase from about 6,000 to 14,000 tons as a result of the project.

Benefits and Justification. The proposed project would be justified both on its own merits and as a pioneer project leading the way to further irrigation development of this type in northeast Thailand. The economic rate of return would be 16 per cent. The individual sub-projects show little difference in the rates of return, as differences in their investment costs per hectare are mainly attributable to the costs of repairing the dams and spillways. These costs were omitted in the economic analysis, as it was considered that these expenditures would in any case be essential to safeguard life and property.

Nong Wai Irrigated Agriculture Pioneer Project (Thailand)

Background

Project Area. The project area is located on the right bank of the Nam Pong river, stretching for more than 30 km in the vicinity of Khon Kaen. The total area covers some 15,000 ha, of which about 13,500 ha can be irrigated. Well over 50 per cent of the project area is under rice cultivation. Due to soil conditions, the cultivation of vegetables and garden areas is restricted to the levees of the major water courses. The project area is typical of rice-growing areas in northeast Thailand

Soils. Most of the soils are of alluvial origin. About 43 per cent of the soils are alluvial hydromorphics and consist of clay throughout the profile. Low humic gley soils cover about 26 per cent of the area and are located on low and middle terrace formations. The top soil consists of loamy sand in the higher sections and clay loam in the lower sections. Soils located on the high and upper levels of the middle terrace consist mainly of grey podzolics and cover about 7 per cent of the area; the topsoil texture ranges from sand to sandy loam over a sandy loam to sandy clay loam subsoil. Several other soil types are currently covered in forest.

Socio-economic Factors. The present farm population is about 25,000, occupying 33 villages within the project area; the 3,500 farms have an average household size of 6.9 and an average labour force of 4. Based on the fact that not all the land is currently being cultivated and the population growth trends, it can be assumed that with full implementation of the project area, a total of 36,000 people with 5,200 farms of 16 rai each will occupy the area in 1985.

The Project

Objectives. The main objective is to develop a complete programme for effective and efficient water-use, which, based on existing water resources and facilities to be developed, would ensure adequate support for the accelerated development of agriculture in the project area. This would include improvement and extension of existing irrigation and drainage facilities, with special emphasis on the provision of necessary terminal facilities; on-farm development and land leveling and consolidation; establishment of feeder roads and market structures; institutional development, including farmers' organizations, credit support research, training, and extension activities; livestock development; and agriculture-related industries which will emphasize the utilization of local raw materials and manpower resources.

The programme for the development of agriculture takes into account the human factor and the present situation of the farmers. The change from subsistence to surplus farming can only be achieved gradually, requiring intensive assistance to the farmer.

From a purely economic point of view, based on net production values, it would appear that the farmers should concentrate their attention on one or two crops, but restrictions imposed by labour availability, risk factors, and the need for home consumption preclude such development. On the other hand, a strong case can be made for crop diversification; activities such as the organization of marketing or the establishment of processing facilities for tomatoes, the canning of fruits and vegetables, oil extraction and animal feed milling in the project area will give farmers the market assurance they need for a diversified pattern of crop production.

The implementation of the irrigation system requires certain land consolidation measures to assure that all farms receive an equitable share of irrigation water. Because Thailand does not yet have a law governing land consolidation, the project can only be initiated when the farmers agree on a voluntary basis, though cooperation with the Land Department is essential for the legal aspects.

Project Works. After the construction of the main irrigation system, a network of tertiary canals was built in the project area to bring irrigation water into the service area. However, only a few of these ditches were in use during the last irrigation season and in most of the project area the minor irrigation system is completely inoperable. The layout for the majority of ditches, based on inadequate topographic maps, needs improvement. Long and cross sections are very irregular; the flow is too large and cannot be properly controlled. Many plots are out of command and others receive water only through adjacent fields (a situation fraught with potential conflicts among the farmers).

Project works include a 50 ha demonstration plot, a main irrigation and drainage system, flood control measures, works on the farm level, the establishment of a training centre and provisions for maintenance of the irrigation scheme. For adequate operation and maintenance, roads will be required along all major irrigation canals and also along the flood embankments. This does not apply so much for the major drainage channels. Siltation will not be a problem here and the channels do not carry any structures which require regular attendance. The roads will also improve the communication between the most important villages, opening the entire area to improved markets. Together with a network of farm roads, each giving access to individual holdings, they will link up with the major road system and thus constitute an integral part of the infrastructural requirements for accelerated development of the Nong Wai project.

Cost Estimates. The estimated investment for the construction works in the minor systems and at farm level will amount to about US\$ 3.4 million, an investment of about US\$ 240/ha. If the future construction costs for the main systems, flood control and half the main road investment are related to the irrigable area, the anticipated investment increases to US\$ 345/ha.

Organization and management. Implementation and operation of the irrigation scheme is coordinated around the establishment of a single, multi-purpose cooperative society. This cooperative will serve the total project area, assuming the responsibility for irrigation services, supply of credit, marketing of products and extension services; it would also ensure the full participation by the farmers. With a minimum membership of 3,500 (the present number of farm units), the cooperative could become a self-sustaining organization. While the project area stretches over more than 30 km from north to south, one association should be able to render services to all members, provided that the facilities are adequately distributed over the total area.

The cooperative will work with government authorities throughout the implementation and operation of the pioneer project. However, during the initial phase, substantial support from the government will be necessary. It will be difficult for any of the existing government departments to handle the promotion of the association alone; its success will depend on the coordinated efforts of all governmental agencies concerned. Therefore, a promotion committee should be organized under the Ministry of National Development, including officials from the Department of Land Cooperatives, Royal Irrigation Department, Department of Credit and Marketing Cooperatives, Department of Agriculture Extension, and the Bank for Agriculture and Agricultural Cooperatives as committee members.

Due to the administrative organization of the governmental agencies with fragmentation of power and competences, several different ministerial departments are currently handling the services and projects within the Nong Wai area. The overlapping of works and resources have caused subsequent results to be unsatisfactory. The maximum effectiveness of a pioneer project in irrigated agriculture, such as the Nong Wai project is intended to be, can only be achieved if all governmental efforts and international aid programmes are concentrated and well coordinated. This approach must be applied during both the planning and implementation period, as well as during the following period of agricultural operation.

It would be unrealistic to have the project solely implemented by one of the existing government departments, since none of the departments is currently able to render all the required services with an optimum of effectiveness; the integrated programme calls for interdepartmental contributions from all agencies involved. In order to avoid the present lack of cooperation, it is necessary to entrust the overall responsibility for the total programme to one agency, which might be named "Nong Wong Wai Project Office". While this agency should be a temporary institution, it should be maintained during the entire implementation period (1971-1984).

Agricultural Production. Current production at Nong Wai is to a large extent subsistence production which never reaches the market. An agricultural development programme under irrigated conditions, however, would mean a large increase in agricultural production, most of which will be marketed. In illustration 24 the net production value of agricultural production in Nong Wei has been calculated under full development (1984), when all farms in the area will have reached their maximum production potential. Total annual income from both crops and animals will amount to more than US\$ 20 million. Income

from off-farm activities will no longer significantly contribute to the farmers' income; it must be assumed that under irrigation conditions with two crops, the farmers will be fully employed in agriculture.

On the basis of an average farm at 16 rai, with 6.9 household members, farm income and per capita income will reach 20,000 baht and 2,900 baht respectively, an income increase of 340 per cent over present levels

ILLUSTRATION 24: ANNUAL INCOME FROM AGRICULTURE IN NONG WAI UNDER FULL DEVELOPMENT			
Production	Total annual income Bht million	Per farm Bht	Per capita. Bht
Crop husbandry	83,780	15,995	2,318
Livestock	<u>21,753</u>	<u>4,153</u>	<u>602</u>
Total	105,534	20,148	2,920

Benefits and Justification. A cost/benefit analysis of the investments over a period of 30 years, disregarding the expenditures for the main canals, indicates an internal rate of return of 31 per cent. The exclusion of the investments on the main canals is justified because they are past expenditures and play no role in the decision concerning the profitability and desirability of the envisioned supplementary irrigation works. With the inclusion of the canal construction costs, the internal rate of return falls to 10.7 per cent.

Go Cong Gravity Irrigation Pioneer Project (Viet-Nam)

Background

Project Area. The project is situated in the northeastern corner of the Mekong delta about 40 km southwest of Ho Chi Minh City and is accessible by excellent paved roads. Access to this area is also possible by navigation via the Cho Gao canal or the nearby Rach Go Cong river. The project area, covering 7,400 ha including 6,300 ha of net irrigable area, was selected as representative of the eastern coastal region of the Mekong delta in Vietnam and its problems typify that region's major physical constraints: non-availability of fresh water supplies during much of the dry season; periodic flooding; sluggish surface drainage; saline soils; and inadequate farm-to-market roadway access. The project will demonstrate effective water control and adequate utilization of land and water resources through complete and closely coordinated systems of gravity irrigation, drainage, flood, and salinity control, and a suitable programme of agricultural services to support intensive crop production and integrated agricultural development in a 6,300 ha service area.

Soils. The soils are uniformly heavy-textured, poorly-drained, and slowly permeable. Soils situated in depression areas along streams, which turn highly saline early in the dry season, are the least productive soils in this area, with an especially high groundwater table and excessive salt content. These soils are currently planted only in the wet season to certain salt-tolerant but low-yielding rice varieties. Most of the other soils are generally suitable for intensive year-round paddy cultivation.

Socio-economic Factors. The project area is one of the most densely populated areas in the basin. The 6,700 ha of land cultivated at present are used by 4,270 farm households averaging about 6.6 persons per household for a total population of about 28,000. Most farmers in the project area operate 1-2 ha of cultivated land, but one third of the farmers cultivate less than 1 ha each; nearly all of the farmers own their land.

Despite the salinity problems in the low-lying area, single crop paddy production has been maintained on around 6,540 ha, or nearly 98 per cent of cultivated land area, supplemented by only small patches of dryland crops, vegetables and banana and coconut trees near the houses. Only 340 ha of paddy land was double-cropped in 1972.

About 35 per cent of the local rice production entered the commercial market. Most farmers sold their produce to village paddy collectors and local mills and the rice was ultimately collected and transported to Ho Chi Minh City by the larger dealers.

The Project

Objectives. Water quality is the over-riding consideration in any proposal to provide irrigation in the project area. Most of the available water in the dry season has salinity levels much in excess of value considered to be acceptable for irrigation of rice, but measurements show that the Song My Tho canal in the vicinity of the project area normally exhibits salinities higher than acceptable only about 2 to 3 months each year (March-May). Irrigation from this source would thus permit large-scale double cropping in the project area, though planting dates for the second crop would still have to be rigidly scheduled to permit cessation of irrigation before the late dry season period.

The objectives of the project include supplying the 6,300 ha pioneer project with irrigation water, averaging 140 million m³/year; improvement of the present sluggish drainage conditions; prevention of the recurrent flooding of the low-lying lands during high tides; and improvement of the road system. Another primary objective of the project is to achieve, on the basis of full irrigation and effective water control, a system of farming that promises maximum area-wide increases of farm production and income, a system which is stable and enduring, insensitive to weather fluctuations and conducive to continuous improvement of land productivity. Detailed investigations have established the fact that for the Go Cong area and for the eastern coastal region as a whole, the best alternative is intensive double-cropping of rice for the foreseeable future.

Project works. Project works will include construction of intake structures on the Song My Tho, along with feeder canals; enlargement of existing pumping installation; enlargement

of existing main canal and construction of a new irrigation network, including 14 irrigation laterals totaling 53 km in length, which in turn will feed on a rotating basis some 350 km of sub-laterals, each serving about 50 ha; improvement of the drainage conditions by means of construction of about 400 km of new lateral and sublateral drains and by providing egress of drainage flows from this system through five major sluices; prevention of recurrent flooding by means of a dike averaging one metre in height along the northern edge of the project area; and strengthening of agricultural services and improvement of local farmers' organizations.

Cost Estimates. The project cost will be about US\$ 7.1 million (\$ 2.1 million in foreign exchange). Illustration 25 gives a summary of the project cost.

Organization and Management. The executing agency for the proposal project will be the Ministry of Agriculture. For the execution of the construction works under the project, a Go Cong pioneer project office would be established to prepare budgets, supervise consultants, initiate procurement procedures, prepare tender documents and evaluate bids for procurement of goods and services, supervise project construction, prepare the initial operation of the agricultural development programme, and serve as liaison with other concerned government agencies. Operation and maintenance of newly completed project facilities during the construction period will be the responsibility of an operation and maintenance section in the project office.

ILLUSTRATION 25: COST ESTIMATE: GO CONG (VIET-NAM) ADB Appraisal Mission's Cost Estimate for the Project a/ US\$ (thousand)			
Item	Total	Local	Foreign
A. Land Acquisition	423		423
B. Civil works			
Construction	3,830		3,830
Materials	710	-	710
Machinery and Equipment	225	-	225
Sub-total	4,765	3,830	935
C. Consulting Services	634	150	484
D. Operation and Maintenance, Agricultural Equipment	299	-	299
E. Contingencies	979	597	382
Total (A through E)	7,100	5,000	2,100

A Go Cong agricultural development centre will be established in the third year of project implementation to provide and coordinate agricultural development services to the farmers in the project area. A farmers' education and technical services section will be responsible

for seed multiplication, farmer education and training, advisory services on the proper use and maintenance of farm machinery, demonstration and extension of suitable types of paddy driers, technical services for improving livestock production, and provision of marketing assistance and other extension services. A rural credit and farm supplies section will be responsible for the assessment of the credit and farm input needs and for assistance in their timely procurement and distribution. A farmers' organization section will be responsible for providing necessary assistance to existing farmers' associations to enable more active participation of local farmers in the agricultural development process. The Agricultural Development Centre will continue to operate until about 1984, the target year for the full realization of the projected productivity improvement. It is envisaged that after 1984, a major part of the agricultural extension work would be taken over by the then-revitalized local farmers' organizations.

Agricultural Production. As the project is implemented, more intensive and productive agricultural use of the land and water resources in the project area would be made possible. With the project, nearly 90 per cent of the project area would be double-cropped compared with only about 15 per cent at present. The yield of paddy would increase from the current level of 2.1 t/ha to about 3.8 t/ha by 1984. Annual incremental production attributable to the project would then amount to about 24,000 tons of paddy and some 6,000 tons of various crops. The annual incremental production net value, allowing for cultivation and harvesting costs, is estimated at about US \$ 1.35 million in 1984, expressed in 1974 dollars.

Farmers will be the direct beneficiaries of the project, with present average annual farm income rising from US\$ 265 to US\$ 747 by 1980 and US\$ 833 by 1984. The average net farm revenue per ha will increase from US\$ 117 in 1973 to US\$ 413 in 1980.

Benefits and Justification. The project would create new permanent jobs for the operation and maintenance of project facilities and for the provision of strengthened institutional services. In addition, an annual supply of about 450,000 additional man-days of farm labour would be needed to meet labour requirements for intensive double-cropping of paddy, which would provide further on-farm employment to benefit particularly the smaller farmers in this area.

After the completion of project facilities, the period during which river water of acceptable quality for domestic use can be reliably delivered to Go Cong city by the Tham Thu canal will be extended from about 7 months to about 9 months in a year, with corresponding reduction of the period over which trucking of water at high cost is necessary. Construction of new roads under the project will also bring benefits to the farmers in the area. Further, the success of this pioneer project would be of great significance to the overall rural development in the Go Cong area, and, because of its demonstration effect, would facilitate large-scale development efforts throughout the delta region.

On the basis of projections of economic benefits and costs, the project's internal rate of return is estimated at 13.4 per cent.

Tiep Nhut Salinity Control Pioneer Project

Background

Project Area. The project area is in the Mekong delta, about 140 km from Ho Chi Minh City. It comprises 50,000 ha along the right bank of the Bassac river, near its outlet to the South China sea. The project lies entirely in Ba Xuyen province and is in many ways typical of large areas in the coastal region of the delta subject to seawater intrusion, soil salinity, and inadequate surface drainage. The project area is roughly in the shape of a triangle with its apex at the town of Dai Ngai. Its eastern side flanks the Bassac and South China sea; its base along the south is defined by the My Thanh river; and its western side fronts the Saintard canal from Dai Ngai to Bai Xau and the Du Tho river from Bai Xau south to Hoa Tu. About 46,000 ha are potentially cultivable, and the remainder is taken up by villages, roads, canals, rivers and swamps.

The project area is fairly flat, rising from sea level to about 2.5 m MSL. The general slope of the land runs down from the eastern portion along the Bassac river to depressions in the south and west. Low-lying zones are adjacent to some of the natural channels intersecting the area, but there is also a network of canals. Relatively high areas are well-drained, but lower land is subject to flooding by excessive ponding of runoff or by seawater inundation. For these reasons about 16 per cent of the total area is not farmed, especially along the coastline. Deeply ponded areas will not be suitable for improved rice varieties without better drainage, nor could the saline coastal areas be cultivated without protection.

Both drainage and seawater intrusion are related to tidal variations in the Bassac and My Thanh rivers. The maximum tidal range of the South China sea in the project area is about 3 m, though at Tan Thanh and Bai Xau, this range is reduced to 2 m owing to the damping effect of the Saintard canal and the My Thanh and Du Tho rivers. Within the project area, tidal fluctuations are further reduced by existing control structures; for example, the water surface of the Cai Xe canal varies by no more than 20 cm as a result of the Ngan Ro dam.

As river discharges decrease in the dry season, the fresh/salt water interface moves inland from the South China sea. Along the Bassac, the salt front reaches Long Phu in February and Dai Ngai in late March. Movement up the My Thanh and Du Tho rivers is more rapid, with salt water reaching Hoa Tu in December, only one month after the end of the rainy season. Farmers report that the reach of the Saintard canal between Chau Khanh and Sung Dinh never becomes saline; this is probably due to the fresh water carried toward the project area by the Maspero canal, which draws water from the Bassac at An Ninh, some 30 km upstream of Dai Ngai and well above the northern-most front of seawater intrusion. As flows rise with the onset of the wet season, the fresh/salt water interface withdraws from the project area. Most points within the project area, have a period during which the local waterway is too saline for irrigation purposes.

Soils. The soils are predominantly heavy alluvial clays, with the exception of some young river and marine sediments along the banks of the Bassac and some locally disturbed soils

along the main canals. Land capability studies have shown that about 20,000 ha have no serious limitations for agricultural production other than a lack of freshwater for irrigation. The remaining area suffers in varying degrees from periodic saltwater inundation and flooding from storm runoff. Soil salinity is reduced by heavy rainfall during the rainy season to levels where paddy can be cultivated, but the growing season is often curtailed if the rains are late or end too early.

Socio-economic Factors. Villages are scattered through the area, with a total population of 860,000 (1973). Khanh Hung is the provincial capital, and with Bai Xau provides administrative, agricultural processing and marketing centres. Land reform in recent years has resulted in most of the land being cultivated by farmer owners. The average farm size is 2.8 ha, with about 70 per cent of the holdings ranging between 1.5 and 3.0 ha in area. Of the 46,000 ha of potentially cultivable land, about 37,000 ha is currently being cultivated. The uncultivated land is mostly along the My Thanh river and is inundated by seawater for most of the year through breaches in the dikes which formerly protected the area. A single wet-season crop of transplanted rice is the main crop in the project area. High yielding varieties are grown on about 4 per cent of the area and only 100 ha produce rice in the dry season. Yields average 2.3 t/ha overall, with ranges from 1.4 to 2.5 t/ha for local varieties and 3.5 to 4.0 t/ha for HYV.

Objectives. The main objective of the project is to improve conditions for rice cultivation in an area of about 46,000 ha which is currently subject to salinity intrusion and inadequate surface drainage. It would provide facilities for low-lift pump irrigation on 15,000 ha of which about 7,000 ha would receive a water supply for a dry season crop. The project is intended as a forerunner of similar large-scale projects in the coastal areas of the Mekong delta subject to salinity intrusion and inadequate drainage.

The proposed project would include: (a) construction of a system of small dams, sluices, and dikes to control water levels and water quality over an area of about 46,000 ha; (b) rehabilitation of existing laterals and construction of new lateral canals; (c) bulk procurement of about 2,500 low-lift irrigation pumps and on-farm works for an area of about 15 ha; (d) establishment and operation of a development centre to provide agricultural supporting services; (e) construction of buildings and procurement of vehicles and equipment for project implementation and operation; (f) consulting services to assist the government in project implementation, operation, and maintenance of the works, including data collection and analysis; (g) technical assistance in the establishment and operation of the development centre; and (h) consulting services for studies of future developments within and adjacent to the project areas.

The approach to salinity control adopted for this project involves construction of 11 small dams on the canals and rivers subject to salinity intrusion; this approach is advantageous over building a single large dam on the My Thanh river in that it poses no major engineering problems requiring lengthy investigation and would provide salinity control and drainage works for a large area at a relatively low cost.

The proposed works will lengthen the period in which freshwater would be available in the canals and rivers of the project area. The number of points where freshwater could be introduced to the project area would depend on the time of the year. An ample supply of freshwater would be available from July through December for the 5,000 ha which it is assumed would receive supplementary irrigation (HYV are already cultivated on about 2,800 ha in the wet season), because the water would be fresh on more than half of the periphery of the project. By the end of March, however, the advance of the salt water front would leave the intake points on the Saintar canal as the only sources of freshwater, and withdrawals in March and April might cause a further advance of the salt water front in the canal. In view of this, a conservative estimate of 7,000 ha has been adopted as the area which could be supplied with fresh water in March and April. This would support a dry-season cropping intensity of just under 50 per cent.

Project Works. The proposed water control system includes 11 dams. Four have already been built, and some preliminary work has begun on the Long Phu dam. The project will include completion of the Long Phu dam and the construction of six new dams, with a typical dam consisting of an earth embankment across the canal or river and a control structure in an adjacent by-pass channel. The control structure is usually in the form of several culverts which can be closed by slide gates on the upstream or landward end, and flap gates on the downstream end.

The purpose of the dams is to control water levels and water quality in the project area. In addition to preventing salinity intrusion, the dams will be operated to drain the project area; during the periods of high rainfall, the slide gates would be opened and water flow out of the project area through the flap gates on the falling tide. As the tide rises, the flap gates would close and prevent return flow. During the dry months of the rainy season, the dams would be operated to raise water levels in the canals by opening the slide gates and flap gates at high tides, and closing them on the falling tide. This would be done only where the outside water quality is suitable for irrigation.

A perimeter dike system to exclude tidal overflow into the project area will extend from Dai Ngai in the north, down to the right bank of the Bassac, up the left bank of the My Thanh, and along the Du Tho river to Bai Xau. The dike would be a simple earthfill embankment with a 4 m crest width and a parallel collector drainage ditch running along its inner side. The excavation of 83 km of collector ditches along existing and new dikes would provide the fill material required for these low embankments.

Canal construction required for the project would consist of deepening of 35 km of existing canals and constructing 65 km of new laterals. The three existing major east-west canals would not require improvement, but the Long Phu-Lich Hoi Thuong canal would be deepened along its entire length of 14 km, and 5 km of new main canal would be excavated east from Long Phu and down along the Bassac river. Twenty-one lateral canals would traverse the intensive cultivation area, with the new ones about 1 km apart. With this spacing the conveyance distance for irrigation would not exceed 500 m and would be less in many areas where low-lift pumping is practiced. Of these 21 laterals, 10 exist but will require enlargement, and 11 will be new.

The project works would create conditions suitable for low-lift pump irrigation on about 15,000 ha where high yielding varieties would be introduced. Experience in the Mekong delta has shown that farmers with access to a freshwater canal or river will cooperate in low-lift pumping and water distribution. The development centre established in the project area would advise the farmers in land levelling and in the layout and construction of their distribution ditches and drains. Most of the land to be irrigated has been devoted to paddy cultivation for many years and only light levelling would be needed. The development centre would advise the farmers on grouping together to contract for land levelling and credit would be provided for acceptable proposals. Demonstration of suitable land levelling attachments for the locally available tractors would be carried out under the project.

Cost Estimates. Total project costs are estimated at US\$ 15.2 million, including a foreign exchange component of US\$ 7.7 million. Costs are summarized in illustration 26.

ILLUSTRATION 26: COST SUMMARY FOR TIEP NHUT PIONEER PROJECT			
	Local	Foreign US\$ (thousand)	Total
Civil works			
Structures	390	590	980
Dikes	420	280	700
Canals	1,170	780	1,950
Roads and Bridges	300	200	500
Buildings	140	60	200
Telephone System	50	110	160
Miscellaneous works	130	120	250
Surveys	100	50	150
Sub-Total	2,700	2,190	4,890
Land acquisition	150	-	150
On-farm development			
Pumps	-	1,250	1,250
On-farm works	1,240	410	1,650
Consulting Services			
Project implementation	70	410	480
Development Centre	50	250	300
Further studies	50	310	360
Development Centre operation	610	50	660
Vehicles and equipment	-	200	200
Project monitoring	80	40	120
Base cost estimate	4,950	5,110	10,060
Physical contingencies	540	440	980
Expected price increase	2,010	2,150	4,160
Total	7,500	7,700	15,200

Organization and Management. The project would be implemented by departments of the Ministry of Agriculture. The Ministry would appoint a senior official as Project Coordinator who would be responsible for ensuring coordination between the departments and with other government agencies.

The development centre would provide general extension services to the farmers in the project area; prepare, in conjunction with individuals or groups of farmers, farm plans with cost estimates and expenditure schedules; draft contracts for contractor's services; prepare design standards for irrigation and drainage ditches, bunds, and land leveling; supervise and approve contractual services for on-farm works; promote among the farmers the concepts of pump irrigation development; and maintain up-to-date records on agricultural development in the project area. Arrangements would be made to provide medium-term credit for the purchase of low-lift pumps, motors and spares by project farmers and for the execution of on-farm works comprising land leveling and construction of distribution ditches.

Agricultural Production. Yields of local varieties in the more favourably located areas are now about 2.5 t/ha, but in less favoured areas the yields range from 1.4 to 2.2 t/ha (depending on the severity of the salinity and drainage problem). A shift to HYV with yields of 4.2 t/ha is expected on 15,000 ha, but in the less favourable areas average yields with the project are expected to reach 2.6 t/ha. Total production is expected to rise from the current 87,500 tons to 166,000 tons (illustration 27). Cultivation of fruits and vegetables, which is now carried out on a fairly large scale in the freshwater zones of the delta, is likely to increase with the availability of freshwater in the dry season, but no attempt has been made to quantify this.

ILLUSTRATION 27: TIEP NHUT PIONEER PROJECT							
Present and Projected Production Values							
	Cropped Area	Yield	Farm-gate Price	Gross Value of Production	Production Costs⁸	Net Value of Production	Net Return from Project Area
	(ha)	(ton/ha)	(US\$/ton)		(US\$/ha)		(US\$ million)
<u>Present Production</u>							
Wet Season							
Local varieties	33,700	2.2	96	212	38	174	5.85
HYV rice	3,200	3.5	96	337	114	223	0.72
Dry Season							
HYV rice	100	4.0	96	385	184	201	0.02
Total							<u>6.59</u>
<u>Future Production</u>							
Wet Season							
Local varieties	27,000	2.6	96	251	48	203	5.48
HYV rice	15,000	4.2	96	404	103	301	4.51
Dry Season							
HYV rice	7,000	4.7	96	452	133	320	2.24
Total							<u>12.23</u>

⁸ Lower production costs for HYV (future) reflect more efficient diesel-powered pumps and improved water management.

Benefits and Justification. A successful pioneer project in the Tiep Nhut area would pave the way for broadly similar development of a large part of the coastal belt where development is constrained by salinity and drainage problems, a potential area amounting to over 30,000 ha. Implementation of the pioneer project would help to establish the availability of freshwater for further development in the project area, the most appropriate cropping calendar for its use, and the need for additional work to augment the supply of freshwater.

The internal rate of return has been calculated as 24 percent, using 1974 prices for estimated capital and production costs.

The caisan Low-Lift Pumping Irrigation Pioneer Project (Viet-Nam)

Background

Project Area. The Caisan Settlement Area covers about 30,000 ha in the lower Mekong delta 170 km southwest of Ho Chi Minh City, between the Bassac river and the Gulf of Thailand. The area lies in the southern part of the seasonally flooded land stretching from the border with Cambodia to include about 500,000 ha where floating rice is by far the dominant crop. Dry season cropping of rice and other crops is limited to small areas along the canals and rivers. Before the settlement, the region was accessible only along the main road between Rach Gia and Long Xuyen and along the major navigation canals (Rach Gia/ Long Xuyen, Rach Soi/Vam Cong and the interconnecting Tan Hiep canal).

To provide access to the hinterland of these main channels, 19 lateral canals have since been excavated and the villages of the settlers were located on the canal banks.

Each year the Bassac and other arms of the Mekong inundate large areas in the upper parts of the delta. The land starts to flood in July or August and the area and depth increases irregularly up to October; the floods then recede to within-bank stages in November or December. The change in water level over the flooded area is gradual and outside the rivers or canals the velocity of flow is very low. In the project area, the maximum depth of inundation averages about 1.5 m, but varies with local topography. During the dry season, water levels in the canals which intersect the project area are influenced by tides in the Gulf of Thailand. Salt water intrusion is limited to small areas near the coast.

The wet season rice crop is broadcast in May and harvested in December, with yields ranging from 1.2 to 1.5 t/ha. Virtually all of the land preparation is handled by tractor operators who contract with the farmers. Dry season irrigation is now limited at about 2,500 ha, or 10 per cent of the net project area. This is in contrast to the high cropping intensities achieved in those parts of the delta free from annual flooding but having a year-round supply of freshwater. The relatively slow growth of dry season irrigation in the area is due to several factors. A second crop can only be grown in the period January-May

and hence virtually all the crop water requirements have to be met by pumping, whereas in the areas of extensive double cropping in the delta, the farmers can arrange their crop calendar so that the major part of the water requirements are met by rainfall. Delivery of so large a quantity of water leads to high pumping costs because of the low efficiency of the locally-made pumps commonly used. Furthermore, while the farmers are generally progressive, few of them have developed on-farm channels which enable them to irrigate their 1 km-long holdings. Given technical advice on constructing farm ditches, and with the availability of more efficient pumps, farmers could be expected to achieve much higher cropping densities. Finally, many of the canals become dry at low tides, and these need to be deepened to ensure a reliable dry-season water supply.

Soils. The soils are derived from recent fine textured alluvium of the delta with some marine influence. They are acid and very fine textured with small areas of acid sulphate soils, presumably a reflection of marine influence at some time in the recent past. The soils of the project area are well suited to rice cultivation because of their low permeability. On the ground and from the air the land appears uniformly flat to the horizon, but as the flood recedes, local depressions, about 10-20 meters in diameter and up to 50 cm deep, become noticeable. These small areas contain highly acidic soils and can be seen from the air as slightly darker patches where rice does not flourish.

Socio-economic Factors. The settlement area has a population of about 50,000. The land is owner-operated in holdings of 2.6 to 3.5 ha. The holding size is governed by the farm distribution pattern established at the time the area was settled. The lateral canals are about 2 km apart and each farmer has a 26-to 35-metre-wide frontage on a canal from which his holding extends 1,000 m.

The Project

Objectives. The primary objective of the project is to promote dry season cropping in the flood zone of the Mekong delta. It will aim at promoting irrigation from existing canals by farmer-owned, low-lift pumps, through supervised credit, technical assistance in on-farm works, deployment of more efficient pumps, and advice on higher standards of farm management. The project would include: (a) deepening of existing lateral canals to ensure a reliable water supply at low tides; (b) construction of new lateral canals and farm boundary realignment for a 6,000 ha extension of the settlement area; (c) bulk procurement of about 2,500 low-lift pumps, and provision of medium-term credit to the farmer for purchase of pumps and for other on-farm improvements; (d) establishment and operation of a development centre in the project area; (e) technical assistance in the establishment and operation of the development centre, the design and construction of the extension area, and the identification and preparation of a further “Caisan type” development on about 50,000 ha; and (f) rural electrification works along selected lateral canals.

The basic approach used in the Caisan project is to raise agricultural productivity in the flood free period, which extends from December through July. The conditions which have to be met for the successful introduction of farmer-owned low-lift pump irrigation are: ready access to a canal with a reliable dry season water supply; well-defined farm holdings; and farmers living on or close to their land. All of these conditions are satisfied in the Caisan Settlement Area, but they are not currently met in other parts of the flood zone. The location of the pioneer project in an area with these advantages provides an opportunity for pioneering modes of development applicable to future settlement projects implemented elsewhere in the region. The project includes a 6,000 ha extension area which would provide experience in planning and implementation of new settlement projects.

Project Works. Of the 190 km of existing lateral canals, about 100 km need to be deepened by an average of half a metre to ensure an adequate water supply at low tides. About 27 km of new laterals would be constructed to form a 6,000 ha extension to the settlement area. These would be similar to the existing laterals which have surface widths averaging 15 m. Farm boundaries would be realigned to give each farmer a frontage on the new laterals..

The project would provide technical advice and supervised credit to allow the farmers to undertake the on-farm development needed to expand dry-season rice cultivation. This would include more efficient low-lift pumps, irrigation ditches, and land leveling. On-farm development on about 6,000 farms, or approximately 18,000 ha, would be carried out during the six-year implementation period. Electric-powered pumps would be provided for the areas covered by rural electrification. In other areas the choice will be between gasoline and diesel motors, depending on the area served and the presence of the farmers (though in the long run, the spread of rural electrification and increasing petroleum prices will encourage electric-powered pumps).

A development centre would be established to provide advice to the farmers about the construction and layout of the irrigation ditches and land leveling. The development centre would establish standards for land leveling and assist the farmers in negotiating contracts. Credit would be provided for acceptable proposals. Demonstration of suitable land leveling and ditching attachments for locally available tractors would be carried out under the project.

Cost Estimates. Total project costs are estimated at US\$ 10.9 million, of which the foreign exchange component would be US\$ 5.2 million. Project costs are summarized in illustration 28.

ILLUSTRATION 28: SUMMARY OF CAISAN PROJECT COSTS	
	Total US\$(thousand) US\$'000
Civil works:	
Existing laterals	800
New laterals	700
Footbridges	40
Land acquisition	80
Surveys	100
Buildings	160
On-Farm Development:	
Pumps	1,500
On-farm works	1,620
Technical assistance:	
Project implementation	480
Further studies	300
Project monitoring:	100
Vehicles and equipment	150
Engineering and supervision	200
Development centre operation	350
Rural electrification	250
Sub-total	6,830
Physical contingencies	930
Price contingencies	3,140
Total	10,900

Organization and Management. The project will be implemented by the Ministry of Agriculture. The Ministry would appoint a senior official as Project Coordinator who would be responsible for assuring coordination between the departments and with other Government agencies.

Arrangements will be made to provide medium-term credit for the purchase of low-lift pumps, motors, and spares by project farmers and for the execution of on-farm works comprising land leveling and construction of distribution ditches.

Agricultural production. Projected production is based on the assumption that about 6,000 farmers in the settlement area and in the extension area would benefit directly from the project. The total area farmed will be about 18,000 ha, and it is expected that dry-season cropping will take place on 14,000 ha. Higher yields for floating rice would result from some supplementary irrigation in the early period of growth (May to July) when the crop often suffers from drought. High-yielding varieties would give higher yields with a

more reliable water supply, and with improved extension services. Current production is about 56,000 tons of paddy, of which 46,000 are floating and 10,000 are HYV. Future production will total about 117,000 tons, including 51,000 tons of floating rice and 66,000 tons of HYV.

Benefits and Justification. The project will directly benefit about 6,000 farmers in the Caisan area and demonstrate techniques likely to be adopted by several thousand other farmers in the vicinity of the project. As a pioneer project, it would serve as a guide to more intensive land use in the flood zone of the delta. The internal rate of return is estimated at 27 per cent with all labour priced at the current rate for hired labour in peak periods (1974). Present and projected production values are presented in illustration 29.

ILLUSTRATION 29: CAISAN PIONEER PROJECT
Present and Projected Production Values

	Cropped Area (ha)	Yield (ton/ha)	Farm-gate Price (US\$/ton)	Gross Value of Production	Production Costs (US\$/ha)	Net Value of Production	Net Return from Project Area (US\$/million)
<u>Present Production</u>							
Wet Season							
Floating Rice	31,000	1.5	96	144	36	108	3.35
Dry Season							
HYV Rice	2,500	4.0	96	385	191	194	0.48
Total	33,500						3.83
Wet Season							
Floating Rice	18,000	1.8	96	173	47	126	2.26
Floating Rice	13,000	1.5	96	144	36	108	1.40
Dry Season							
HYV Rice	14,000	4.7	96	452	140	312	4.37
Total	45,000						8.03

Chapter 10

Summary and Conclusions

Introduction

This report has summarized the research and development programme carried out under Mekong Committee auspices from 1967 to 1977 in the field of agriculture. The objective has been to assist the riparian countries to make the best possible use of water resource development potentials to meet food needs and to contribute to the national economies both from the short-term and the long-term point of view.

Agriculture in the lower Mekong basin is complex because of the wide diversity in climatic, soil and water supply conditions and because of the rapid technological changes that have been taking place in many parts of the lower basin over the past several decades, including changes in agricultural methods and practices.

To obtain a proper understanding of this complex situation and to devise feasible improvement measures, the agricultural programme of the Mekong Committee has included a great number of studies and projects, especially over the past decades focused on all known significant problem areas. The importance of this programme, as one of the key elements of the Mekong Committee effort, was underlined by the establishment of a special Agricultural Sub-Committee of the Mekong Committee in 1971. The Sub-Committee held nine meetings devoted to furnishing guidance to the Mekong Committee for formulating the most effective agricultural research and development programme.

The result of these efforts has been a voluminous series of reports and papers, each contributing to the overall programme. The objective of this book has been to bring all of those contributions together in as systematic a manner as possible and thus to make the information readily accessible.

Agricultural Resources of Lower Basin

Conventional analyses of the agricultural potentials of a basin or region generally are limited to consideration of climatic, soil, and water supply, with some attention to fisheries, which in fact do represent the major parametres of concern from the short-term point of view. The numerous studies of the lower basin included in the Mekong programme, which have addressed all of the natural resources of the basin, have revealed a close relationship between agricultural development and conservation of forests and wildlife, and demonstrate

that to achieve optimal long-range sustained yields the overall plan for agricultural development must integrate forest/wildlife conservation together with rainfed and irrigated cropping and aquaculture. Thus our discussion has encompassed all natural resources which contribute to a broader concept of agricultural management, which could be called “environmental agriculture”.

- (i) **The Northern Highlands** comprise northern Laos and parts of northern and northeast Thailand, and is a mountainous region of steep and rugged topography containing only a few sizeable upland plains. It is covered at higher elevations mostly with hill evergreen forest (much of it now replaced by grasslands because of swidden agriculture), and at lower elevations with dry evergreen and deciduous forests with only the valley floors usable for cropping (mainly rice).
- (ii) **The Annamite Chain** is a mountainous zone from 50 to 300 km wide which extends some 800 km within the basin in a northwest-southeast direction, and separates the Mekong drainage on the west from the South China sea drainage on the east. While previously heavily forested, many areas have been cleared for swidden agriculture and are now covered with unproductive grasslands. The opportunities for agricultural development are mostly limited to the tributary valleys.
- (iii) **The Southern uplands** comprise the Elephant and Cardamom mountains separating the Cambodian part of the Mekong Plain from the Gulf of Thailand. The rainfall is very heavy and the region has little scope for agricultural development.
- (iv) **The Korat Plateau** comprises northeast Thailand and adjacent parts of the Lao PDR from Pa Mong south to Pakse. The plateau is a large (550km x 500 km) inter-mountain basin, with thick underlying salt deposits, and much of the area is somewhat dry due to the rainshadow effect of the surrounding mountains. Deforestation is extensive and many areas are covered with scrub or grassland. While an important agricultural area historically, its present productivity barely matches local food needs; but protection can be significantly increased through water control including irrigation, flood control, and drainage.
- (v) **The Mekong Plain** is the vast low-lying area (800 km N/S by 600 km E/W) traversed by the Mekong river, comprising most of Cambodia, the Mekong delta and a small part of the Lao PDR. The Vietnamese portion of the Mekong Plain is the most densely populated part of the basin, with rice grown on 50 per cent of the land area. While the present agricultural productivity is substantial, the potentials are excellent for much higher production by means of water resource development.

Climate is a major constraint governing productive agriculture. While little can be done to increase rainfall or to decrease droughts, much can be done to develop the natural water resources and this is the major thrust of the Mekong programme. Although the entire lower basin is in a tropical monsoon region, the basin actually has three distinct major climatic zones: (i) a tropical monsoon zone with heavy rainfall most of the year (most of the Southern Uplands much of the Annamite Chain, and the southern tip of the delta); (ii) a

tropical wet and dry zone with a distinct dry season (December-May), which is the dominant climate in the more populated areas of the basin (Korat Plateau and Mekong Plain); and (iii) a subtropical wet and dry zone, with cooler temperatures than the tropical wet and dry zone (upland parts of Northern Highlands and highest parts of the Annamite Chain).

The total annual rainfall is sufficient to grow wet-season crops anywhere in the lowlands and uplands of the basin, but the distribution of rainfall during the growing season tends to be erratic with many years experiencing serious damage from floods or droughts. This has fostered land leveling and use of bunds for holding rainwater on the fields. An analysis of the effect of rainfall variations on paddy production in northeast Thailand by model simulation showed that drought conditions are especially deleterious in reducing the planting area if occurring in the early part of the growing season and in reducing harvests when occurring during the flowering season.

Climate is a serious limiting factor in many parts of the lower basin and particularly in northeast Thailand. Rainfed paddy production is erratic, and the supplemental water needed for sustained yields amounts to 280 mm in an average year and 800 mm in a drought year (1 in 10 years). The total demand in northeast Thailand for supplemental water on irrigable paddies is about 14,000 million m³ in drought years, as compared with the total local water supply potential of 15,000 million m³ assuming it can be stored and distributed. For diversified and double cropping the total requirement is estimated at 41,000 million m³. The deficit (27,000 million m³) can only come from the Mekong river. A deficit of even larger magnitude exists for the agricultural development of the Mekong delta in Viet-Nam.

The peak flows in the Mekong river and its tributaries are spectacular and can be destructive, but the low flow levels are usually the limiting factor now influencing the scale of potential agricultural development. The great variations in flow between seasons and between years necessitate substantial storage to provide more uniform flows for purposes of irrigation as well as for water supply for community and industrial purposes, power generation, and to provide for flood control and navigation improvement.

Sediments in the Mekong river are often said to be essential for annual replenishment of nutrients in delta soils, but an evaluation of these deposits in the Vietnamese portion of the delta indicates that the annual deposits can hardly influence the fertility of delta soils. However, in Cambodia, fresh silt deposits in areas adjacent to the natural river levees and in the flood lands around the Great Lake do contribute to maintaining soil productivity, and alternative provision of fertilizers may be required if these deposits are diminished by upstream developments.

The water of the Mekong river and its tributaries is of low mineral content and may be considered excellent for both irrigation and community and industrial purposes. The only exceptions occur in the delta during the dry season where about 0.5 million ha are affected by saltwater intrusion from the sea, sufficient to place limitations on agricultural development there.

Other than for the low flows in the dry season, the greatest constraint on agriculture in the lower basin is rainy season flooding, especially in Cambodia. Toward the end of the rainy season in September-October, some 10 million ha of lowlands are flooded, including the entire delta, the Great Lake lowlands, the portions of the Korat Plateau along the major streams, large parts the Vientiane and Se Bang Fai plains, and parts of northern Thailand. Although rice growing has been adapted to flood conditions in various ways, this represents only a very partial use of the floodwaters. While tributary reservoirs play an important role in alleviating floods, in the long run control of the mainstream itself offers the best potential for enhancement of agriculture through reducing flood frequencies efficiently to bring large areas of now unusable land into production.

The fundamental constraint for agricultural development in the lowlands is water management, including management of the tributaries and the mainstream.

Assuming flood and drought conditions are improved through water control, soil characteristics become the major constraint for agriculture. Both a soil map and a land capability map (scale 1:1,000,000) have been prepared, and with the aid of satellite imagery the soil map has been improved to serve also as a pedo-morphological map. While the maps are small scale they are very useful for preliminary agricultural planning.

The soils of the lower basin have been formed as a function of parent material, climate, hydrology, and age. Vegetation, as in most tropical zones, has been of minor importance in that humus formation is usually not a permanent characteristic. Although 29 major soil types have been recognized in the lower Mekong basin, for agricultural development purposes these can best be grouped into eight categories including four lowland and four upland soils.

The lowland soils, which occupy most of the Korat Plateau and the Mekong plain (the areas with the greatest agricultural potential), are primarily hydromorphic with gley characteristics at shallow depths. The lowland soil types include:

- (i) **The Coastal Complex Soils** (along the sea coast in the delta) are generally clayey, low in organic matter and with high cation exchange capacity (CEC). They are usually slightly acidic and suited for paddy; however saline intrusion is the dominant factor governing use.
- (ii) **The Delta Land Soils** are clayey soils with varying quantities of organic matter depending on drainage, and often with accumulated sulfides which can cause extreme acidity and other toxicity problems when reclaimed. These soils can grow paddy to a limited extent but their full potentials can be realized only by water control and irrigation by which acid conditions can be minimized or even eliminated.
- (iii) **The Flood Plains Soils** formed from flood-deposited sediments, are mostly clayey with low organic matter, medium CEC, and are slightly acid and moderately fertile. The levee soils are well suited to dry-foot crops depending upon flood conditions, whereas the backswamp soils are generally suited to paddy. The productivity of the flood plains would be greatly enhanced by flood control or flood regulation

- (iv) **The Groundwater Complex Soils** are typically silt loam with low surface clay content, low organic matter, low pH, and low CEC. These soils are suited for paddy, but for upland crops both drainage and fertility level must be improved. Some of the groundwater soils on the Korat Plateau contain salts originating from remnants of ancient salt formations; these saline soils can be reclaimed by improvement of drainage.

In upland areas the groundwater is well below the surface. The soils are grouped into four categories: **The Podzolic soils:** These are low in clay and organic matter, have a low CEC and low, water-holding capacity but, with use of fertilizers. They are generally capable of sustained yields of upland crops. **Red and Black Soils** suited to upland rainfed agriculture including rubber, corn, and cotton. **Lateritic Soils** occur in the uplands of the Karat Plateau and Mekong Plain, have laterite at shallow depths, and are generally not suited for field crops. And the **Mountain Soils** are mostly shallow soils derived from sandstones and acid igneous rocks with limited potential for agriculture, even for shifting cultivation.

The soils of the lower basin have many limitations for agriculture, typically including low CEC, low organic matter, and high acidity and moreover sizeable areas are affected by saline intrusion, floods, and droughts. Mitigation of these constraints is one of the key objectives of the Mekong programme.

The natural vegetation is the floristic expression of the physical complex described above, including the geology, climate hydrology, and soil characteristics. The term “natural vegetation” is used to mean any non-agricultural vegetation, the term “climax vegetation” being the idealized mature forest (or grassland in some seasonally inundated areas) that would occur in the absence of human influence. Practically all of the forests of the lower basin have undergone periodic clearing and regrowth due to swidden agricultural practices over many generations.

The basin forests comprise two major groups, evergreen and deciduous, and each of these groups is further divided into types. Among the evergreens, tropical rainforests occur from sea level up to 1,000 m in areas having an annual rainfall of 2,000 mm or more distributed throughout most of the year; hill evergreens grow in areas of heavy evenly-distributed rainfall above about 1,000 m; coniferous forests are typically found in sandy soil areas at 600-1,400 m; mangroves occur only in the Mekong delta; and freshwater swamp forests occur mostly in Cambodia around the Great Lake and in the backwater areas of the Mekong. Deciduous forests include monsoon forests, confined to areas with pronounced wet and dry seasons, generally at elevations below 1,000 m with 1,250 to 2,000 mm annual rainfall; and dry dipterocarps, also called “savanna forests” replace the deciduous monsoon species when the rainfall drops below about 1,250 mm and where the dry season is as long as six months. In addition to these forest types, many lower basin areas formerly forested are now covered with savannas or grasslands especially Imperata. The scarcity of natural grasslands (except in seasonally inundated areas) indicates that the edible grasses (the main food for grazing animals) depend on human intervention to maintain appropriate growing conditions especially by the use of fire.

The forests of the lower basin have played a vital role in human societies over the centuries in furnishing food, fuel, construction materials, medicines, resins, gums, and pesticides, and in supporting wildlife. It appears that, while production is adequate in the Lao PDR and in Cambodia, the requirements are from 4 to 10 times production in Thailand and Viet-Nam. Moreover requirements are increasing while the operable forest areas are being reduced by 5 to 10 per cent per year.

In the long run the forests may play an even more important role in human economies through their capacity to produce fuel and raw materials, on a sustained-yield basis. The forests should be regarded as a renewable resource which with proper management can give sustained yields for the indefinite future. Unfortunately, over-exploitation and lack of management have drastically depleted the forests in recent decades. The indications are that the rate of depletion, due to population pressures and modern technology, is far beyond the natural ability of the forests to replenish themselves. Much forested land is thus being converted to grasslands which are permanent in time scales relevant to land use planning, and a major effort is required to conserve existing forests, re-establish depleted forests, and establish plantation forests to meet the basin's requirements for forest resources.

Studies sponsored by the Mekong Committee have produced some fairly comprehensive assessments of the terrestrial and aquatic fauna of the lower basin. These show that the basin supports at least 212 species of mammals, 696 species of birds, 213 species of reptiles and amphibians, and 800 species of fish. Ten of the mammals are endemic (found only in the Mekong basin) and 14 are considered rare and endangered. The basin birds include about 500 resident species (at least 12 endemic) and another 160 as winter migrants. Generally, the faunal diversity closely follows floral diversity, with the greatest diversity in the evergreen forests and the least in agricultural areas.

From the point of view of agricultural development, the main significance of wildlife is its role in controlling agricultural pests and as a genetic resource. The role of wildlife in controlling pests in the lower basin cannot be quantified because of lack of data but generally this role is indicated to be a major one and merits careful evaluation and assessment. With regard to genetics, the domestic animals of the lower basin not only were derived from wild species but the wild species represent in some cases raw material for developing new breeds with superior disease resistance and ability to adapt to the available food supplies.

Because terrestrial wildlife is closely associated with forest habitat, the wildlife resources of the lower basin are being rapidly depleted, and the continuing pressure of population growth will tend to intensify such depletion.

Increasing population pressure means that the prospects for protecting the remaining forests and wildlife appear dim indeed considering the many other problems facing the governments of the riparian countries. The Mekong programme has therefore focused on designing practicable remedial measures that are realistic in terms of implementation, particularly if considered and implemented as an integral part of overall water resource development planning. It appears that the only conservation measure which can be successfully implemented at this time would be to establish a series of protected areas to preserve

typical portions of the forest habitats throughout the basin, integrated with modernization and intensification of agriculture in the lowlands. A system of 29 national protected areas has been suggested, totalling about 7 per cent of the lower basin land area, including existing protected areas. Moreover, these protected areas should be considered as “core areas” for larger land and resource management areas surrounding and incorporating them. Within the core area, no extractive activities would be allowed but the forest resources of the surrounding buffer zones would be subject to limited utilization for the benefit of the local residents.

Other complementary wildlife conservation measures envisioned in the Mekong programme include livestock development in agricultural areas for improved production of domestic animals, thus decreasing reliance on wild animals for protein, and captive breeding of wildlife to satisfy the demand for zoos and laboratories and possibly even for human consumption, both for reducing pressures on the natural wildlife and also for reintroducing these species in defaunated forest areas.

An important aspect of the Mekong programme’s approach to conservation of natural resources is recognition that comprehensive water resource development in the lower basin affords an opportunity to protect forest habitats and wildlife. It is clear, by projecting current trends, that unless something is done, the forest habitats and wildlife are doomed to near-extinction. However, incorporating the conservation measures noted above into water resource project development, as an integral part of these projects, and thereby ensuring the means for financing and administering these measures, provides hope for some very real progress even in the short period of the next few decades.

The aquatic fauna of the lower basin is very rich indeed, including over 800 species of finfish alone. The finfish, together with shellfish, are by far the most important animal species economically and nutritionally and also offer the greatest potential for continuing development through aquaculture. In view of the importance of fisheries to the economy, the Mekong programme included a basinwide fisheries study, the first such comprehensive attempt to quantify the complex fisheries of the basin, including those associated with riverine, lacustrine, inundation, and estuarine habitats. While by no means representing a complete assessment of the basin fisheries, nevertheless the study furnishes sufficient hard data to provide the basis for basin water resource planning at this stage of the Mekong programme. Salient findings of the basin wide study include:

- (i) While construction of a dam and reservoir may well result in decreased riverine fish production, this loss may be small in comparison with the new fishery created in the reservoir (the value of which in the case of some tributary projects has approached that of the power generated);
- (ii) Despite the richness of the reservoir fishery, without proper management over-fishing is probably unavoidable. Conversely, if properly managed to take advantage of all natural food sources in the reservoir (by filling the available ecological niches), the reservoir fish productivity can be very substantially increased;

- (iii) By furnishing a reliable supply of freshwater throughout the dry season, the reservoirs create the opportunity for large-scale downstream aquaculture including fish farming in the vicinity of irrigation canals and fish cage culture directly in the water courses;
- (iv) Aquaculture must be carefully planned and integrated with irrigation development planning so that modernization of agriculture does not lead to constraints in aquaculture and conversely modernization of aquaculture does not constrain high productivity agriculture. Water supply systems can be designed so as to satisfy this basic condition.

In general the fishery gains in the new reservoirs and from downstream aquaculture can both compensate eventual losses and, with good management, result in a large net increase in fish production both on a regional basis and from locality to locality. The Mekong programme therefore is focused on (i) achieving optimal production of desirable species from reservoirs, and management of the reservoir fisheries to ensure optimal socio-economic gains (such management would include licensing of fishermen, establishment of fishermen's associations and facilitating provision of infrastructure such as storage and marketing facilities), and (ii) developing the most appropriate technologies for construction and operation of aquaculture systems including improvements in fish food supplies and developing the apparently lucrative aquaculture potentials in brackish water delta areas.

Agricultural Development Programme

The long-term agricultural development model is driven by two immediate, compelling concerns. First, the population of the basin is increasing at a rate of 2.2 to 2.7 per cent per year, so the basin must increase its agricultural production if only to feed its own population. At this growth rate, which assumes use of birth control methods to the extent possible, the current basin population of about 35 million will increase to over 65 million by the year 2000 (illustration 30).

ILLUSTRATION 30: POPULATION PROJECTIONS: MEKONG BASIN AREAS OF THE FOUR RIPARIAN COUNTRIES (In Thousands)						
	1975	1980	1985	1990	1995	2000
Cambodia	7,308	8,309	9,539	10,951	12,450	14,155
PDR Lao	3,134	3,563	4,090	4,696	5,391	6,131
Thailand	15,936	18,385	20,902	23,764	26,756	30,124
Viet-Nam	8,285	9,505	10,705	11,990	13,370	14,830
Totals	34,663	39,762	45,236	51,401	57,967	65,240

(Note: Country estimates for the period up to 1985 have been guided by the UN Population Division Projections, 1965-1985 (provisional). Projections for the years from 1985 to 2000 are based on an extrapolation procedure following earlier trends. From Mekong Committee Annual Statistical Bulletin, 1970).

Second, the natural resources for agriculture are being heavily over-exploited, leading to the destruction of potentially renewable resources such as soil, forests, and wildlife. These resources must be used at a level which does not exceed their sustainable yield, but the over-exploitation of the basin resources can only be controlled if basin residents have a viable alternative. The long-term model therefore presents such an alternative, whereby optimal use of the available resources will both feed the increasing population and conserve the natural resources of the basin. Conservation and development thus cannot be separated, but are complementary products of the same policy.

A third, perhaps less immediately compelling, reason for the long-term model is that many years of agricultural instigations now indicate that the basin is capable of producing considerably more than the requirements of the basin itself, if the conditions for sustained yield agriculture are met. This surplus productivity can earn considerable revenue for the basin.

Existing Agricultural Systems

Because of the wide diversity in the natural resources of the lower basin contributing to agricultural development, many agricultural systems have been applied in various parts of the basin, each adapted to local conditions. These existing systems may be divided into three broad categories: traditional agriculture, extractive agriculture, and modern agriculture.

- (a) **Traditional Systems** include swidden agriculture in the uplands, and banded field agriculture in the lowlands, and for both categories the most important crop is rice. Both are permanent subsistence farming systems, suitable for indefinite use for closed agricultural economies with low population pressure. For upland farmers about 30 ha per family are needed for subsistence, whereas 1.5 to 3 ha is sufficient with banded fields.

Swidden agriculture ("slash and burn") is a rotational system which requires very few tools or draft animals and produces high yields (1 to 4 tons/ha), but after 1 or 2 crops the fields must be left to fallow for at least 10 years. Swidden agriculture has come under wide attack as being destructive of forests and watersheds but, when properly performed with suitable fallow periods and in areas having soils with suitable regenerative capacity, it appears to be an effective way of cultivating areas where "conventional" farming would be impracticable. In recent decades, however, population pressures have led to shortening of fallow periods and to use of non-regenerative soil areas, resulting in extraction of fertility and conversion of the areas to grassland of no significant agricultural value. Because swidden farming can be a viable economic base for hill tribes, it is important to introduce measures for restoring swidden agriculture on a self-sustaining basis.

Banded farming is a self-regenerating cycle including annual ploughing, fallowing during the dry season, and nitrogen fixation when the field is flooded during the growing period. The dominant parameter is the hydrologic regime, and many rice varieties

have come into use (with growing periods from 100 to 270 days) to suit these variations. Three types of bunded fields are in common use in the lower basin: the Khmer system using small square fields; the Vietnamese system using long narrow fields; and the Thai system in which the field sizes and shapes are adapted to the local conditions.

Total rice production from bunded fields has been greatly increased since 1860 in the lowland areas of the basin, not only for local use but for export, primarily by expanding the area under cultivation, through provision of canals. It is clear, however, that traditional agriculture has now reached its limits in the lowlands and that it has already exceeded these limits in northeast Thailand. An estimated 50 per cent of the bunded fields in the northeast are marginal, experiencing unpredictable yields and not infrequent failures. The production ceiling in the northeast from bunded field farming is estimated to be about 6 million tons/year. Further intensification of rainfed bunded field farming, although attempted for many years, has not yet met with success. Yields per hectare have not been raised appreciably.

- (b) In recent decades (since the 1950s) population pressure in the basin has led to **extractive commercial agriculture**, particularly in northeast Thailand, through clearing of the lower-lying forests to produce crops saleable on the international market, especially feed crops. In this period such upland commercial crop production in Thailand (cotton, sugar, kenaf, cassava, and maize) has earned over US\$ 4,000 million in foreign exchange. Unfortunately, this type of farming leads to progressive depletion of soil fertility so that in the long-term part of the cleared lands may progressively degenerate into wastelands.
- (c) **Modern agriculture systems** include ranching, integrated live-stock farming, forest farming, commercial plantations, and irrigated farm systems. While all of these might be feasible for application in the lower basin, all except irrigated farming have quite limited potentials. In fact, only irrigated farming has the potential for the wide application needed to meet anticipated food needs. With full development of ongoing and proposed irrigation schemes (including mainstream development), nearly 4 million ha will be benefited, making up about half of the existing planted areas. Apart from certain areas to be reclaimed through flood control, the irrigation schemes will serve the existing farmlands. Such irrigation will permit a wide diversity in choice of crops, markedly increase yields (for example, from 0.7-1.6 to 3-4 tons/ha for rice), and promote rural employment.

Because irrigated agricultural systems require drastic change in traditional farming methods, many problems have been experienced in attempting to establish irrigated farming in the lower basin. Such problem areas include:

- provision of adequate water and control of the water supply so that the farmer has it when he needs it, which means provision of an adequate distribution system;
- soil problems resulting from numerous sources including low fertility for crops other than rice, water logging, and subsoil exposure;
- drainage to remove excess water and to enable timely harvesting to clear the land for the next crop;

- ensuring adequate operation and maintenance of irrigation facilities;
- land consolidation and field re-levelling to adapt the existing bunded field patterns to irrigation;
- farm management (choice of crops, fertilizers, etc.);
- farm equipment, credit for financing farm improvements, equipment, and agricultural chemicals; and
- community public services as a basis for sustaining an acceptable quality of life in the new agricultural community (which may be characterized as “hectic” compared to the easy-going attitudes of traditional farming practices).

More intensive use of lowland areas through modern sustained-yield agriculture is the only alternative which will meet the basin’s critical food needs over the next several decades. Such irrigation on a wide scale will also require large supplies of water to be made available by an adequate system of dams and reservoirs.

Agricultural Development Model

Agricultural development in the lower Mekong basin, while depending heavily on increasing irrigated agriculture, would encompass all types of cropping from dry farming to gravity irrigation with full water control, including numerous intermediate alternatives such as irrigation with partial water control, drainage only, use of groundwater, and pumping directly from streams. In all cases traditional farming practices are to be modernized, achieving the transition through a step-by-step process proceeding from research and development to controlled application on a relatively small scale to limited full-scale application, and finally to regional application.

- (a) Agricultural Research** in the lower basin over the past 20 years may be likened to the proverbial broth being made by too many cooks. Despite a great number of studies and projects, the overall impact on the agricultural economy in the basin, while significant, has hardly been the bonanza anticipated in the concept of the “green revolution”. This is not to belittle the very real achievements attained; in fact it is estimated that the gains represent an internal rate of return on the research investment of about 47 per cent. However, in terms of total production, these gains have represented an increase of only about 15-20 per cent, less than the population growth.

Substantial progress has been made recently in development of varieties, both for rice and upland crops, which exhibit better quality, improved resistance to drought, floods, pests and diseases, and improved timing of flowering to suit hydrologic variations, with potentials for widescale application throughout the region. However, relatively little progress has been made in developing improved methods for using fertilizers and farm machinery, controlling pests, and improving livestock production. The large gap between the crop yields obtained at research stations compared with actual farmer yields is noteworthy, and analysis of the reasons for this has led to some fundamental understandings which guide the continuing Mekong research programme.

These include the following:

- (i) The objective is not maximum yield but optimal yields consistent with the farmers' constraints including limited availability of production input and the time requirements for transition to new methods; thus, for example, the objective in rice cropping could be not to achieve 10 tons/ha, but to go from 1-2 to 2-4 tons/ha and eventually to 6 tons/ha. A significant finding thus far is that small doses of fertilizer, especially if used as top dressing, are affordable by the farmer and give excellent extra yields for the limited investment.
 - (ii) A key step in proceeding from the research station to practical application requires more field trials, including outreach trials as distinguished from unreplicated trials for proving the practical aspects of the new methods before proceeding to unreplicated trials and to larger scale applications.
 - (iii) While international, regional, and national research stations are important in the overall research complex, the most important element is the local station. Only one of these, together with its outreach activities, is needed for each type of ecological zone, and through operation of these as an organized network the results can be widely applied throughout the basin.
 - (iv) Emphasis of the research should be directed also to developing optimal cropping systems, because land preparation, inputs, fertilizing methods, etc., for multicrop cultivation offer considerably greater potential for increase in total productivity than single purpose research, such as selection of varieties.
- (b) **Pioneer Agriculture Projects** cover all major soil types and cropping patterns found in the basin of sufficient magnitude both in area and in number of farm families to demonstrate the feasibility of the basic irrigation practices selected for the project and to identify and correct any significant problems, before proceeding to overall regional application. The pioneer programme includes a total of 11 projects, including three projects in pump irrigation, an irrigation rehabilitation project to permit more efficient use of unregulated river flow, a storage irrigation project with different degrees of water control, a project for improving lowland rainfed rice cultivation a project for improving upland rainfed cropping, a project making use of a group of small reservoirs or tanks, and three projects in the Mekong delta involving provisions for salinity intrusion control, using low-lift pumping for dry season cropping, and featuring year round irrigation.
- (c) **Full Project Development** building on successful pioneer projects will require systematic estimation of farm benefits for identification of areas to be developed and of appropriate farming methods. This is a complex analysis which can best be approached in terms of "clusters" of parameters affecting yields, with each cluster representing a group of parameters having a more-or-less independent impact on the yield. For purposes of model evaluation, three clusters are distinguished: (i) Drought (including the factors of rainfall, evapo-transpiration, root zone water holding capacity,

and depth of ponding and percolation); (ii) excessive water (rainfall, storage capacity, drainage, deep percolation, and flooding); and (iii) soil properties. A productivity rating is calculated for each cluster (usually from 0.4 to 1.0); these cluster ratings are then multiplied to give the overall relative productivity level for the land unit under consideration.

The next step is evaluation of the impact of fertilizers on yields. Classes of yields include the theoretical possible yield, the potential (research) yield, the maximum yield (potential yield multiplied by the cluster productivity rating), the obtainable yield (less than the potential yield to the extent fertilizer inputs are less than ideal) and the actual yield (average farm yield in the area under consideration). Thus the actual yield will equal the maximum yield if the limiting constraint is not fertilizer, and it equals the obtainable yield with fertilizer applications up to the point where marginal costs equal marginal benefits. It is less than the obtainable yield when fertilizers are in short supply. The optimal fertilizer application and corresponding yield are obtained from crop response curves (mostly derived from farmer field demonstrations) showing the yield increase versus fertilizer application.

The analysis discussed above is carried out by means of three independent consecutive computer programmes, in which the output from one programme is the input for the next. The best choice for the farmer is that which yields the maximum income (within the limits of the farmer's capability to invest and take risks), taking into consideration his resources in land, labour, and working capital. Such computer modelling should help project planners to make more rational decisions, i.e., decisions likely to be accepted by the farmers.

The network of experimental and demonstration stations established under the aegis of the Mekong Committee together with the riparian countries includes five stations located throughout the lower basin: the Korat Plateau of northeast Thailand at Kalasin, the Vientiane Plain of the Lao PDR at Hat Dok Keo, the Central Delta Highlands of the Socialist Republic of Vietnam at Eak Mat; the lowlands of Cambodia at Prek Thnot; and the Battambang Plain at Banan in Cambodia. Significant findings as of 1977 include:

- (a) **Eak Mat:** Trials featuring upland crops showed the Juma 1 variety of upland rice to be by far the best of those tested, yielding 4 tons/ha. Sweet potatoes were found to be the only vegetable suited to the rainy season. Yields for other crops were 2.5 tons/ha (dry season) for soybeans, 2.5-3.0 tons/ha (rainy season) for maize, and 1.7 ton/ha for groundnuts when sown in mid-April.
- (b) **Prek Thnot:** The trials, mostly on wet rice, indicated that water requirements are 6,500 m³/ha in the rainy season and 10,000 m³/ha in the dry season, a spacing of 20 x 15 cm was found optimal, a tilling depth of 10 cm produced 1 ton/ha more than tilling at 15-25 cm, the best fertilizer formula of those tested was 60-60-0 N/P/K allowing yields of over 5 tons/ha, and direct sowing saves 15 per cent on labour over transplanting.
- (c) **Banan:** Trials indicated that 60-30-0 N/P/K may be the best fertilizer combination for rice but yields were only 2.2 to 3.0 tons/ha, hence more studies are in order.

Introduced varieties of groundnuts yielded 4 tons/ha compared to 1-2 tons/ha for local varieties. Several new agricultural implements were produced including a seed drill which greatly reduces labour input.

- (d) **Hat Dok Keo:** Variety trials on vegetables gave yields up to 60 tons/ha for tomatoes, 40 tons/ha for onions, 40 tons/ha for potatoes, and 7 tons/ha for watermelons. Several farm implements were improved including a rice transplanting platform, a threshing apparatus, a rice-mowing device, a weeding tool, and a furrower.
- (e) **Kalasin:** Tests have focused on improving the soil fertility and physical characteristics of the poor soils typical of northeast Thailand. Irrigated rice land can hardly be used for other crops because an impermeable layer forms at shallow depth in paddy fields. With 45-60-60 N/P/K fertilizer application and a spacing of 30 x 25 cm, irrigated rice yields were 6-8 tons/ha. Water requirements were determined for a variety of upland crops, and improved farming tools were developed. Jute fibre yields were highest (1.4 ton/ha) with narrow spacing (25 x 10 cm). Ground nut yields were 3 tons/ha with 30-60-40 N/P/K, and soybean yields reached up to 3.9 tons/ha at 20-120-20. Potatoes yield up to 39 tons/ha but the crop may fail in abnormally hot years.

Pioneer Agricultural Projects

In order to facilitate the transformation of agricultural practices in the lower basin from traditional concepts to applicable modern methods, including application of the findings from the experimental and demonstration stations, the Mekong Committee has made established a network of 11 pioneer agriculture projects, selected and located to correspond to the major soil types and cropping patterns found in the lower basin. An important feature of the entire programme is provision for continuing project monitoring and assessment of the technical, agronomic, and institutional aspects of each project with partial attention to farmer response and to establishment of appropriate farmer organizations in which local leaders have key roles for project implementation. Each of the projects is designed as a relatively short-term operation to develop solutions applicable to the current situation based on current availability of fertilizers and current agricultural prices; hence the pioneer projects do not necessarily indicate the long-term potentials of the basin but serve as an important step towards this goal. The total costs for all 11 projects would be about US\$ 83 million (at 1975 prices).

The 11 projects with respect to type and objective can be summarized as follows:

(a) **Muk Kampul (C) (pump irrigation):**

- (i) **Agricultural land type:** Typical riverine lands along Mekong river including bourrelets (river levee), chamcars (back slopes of the river levee), and adjacent low-lying ricelands.
- (ii) **Soil types:** Silt loam on levees and clays in lower areas.

- (iii) **Objectives:** To demonstrate practicability of pump irrigation for riverine lands, including effective water distribution and management.

(b) Bovel (DK) (gravity irrigation):

- (i) **Agricultural land type:** Typical lowlands in Mekong Plain.
- (ii) **Soil type:** Heavy clay (90 per cent brown hydromorphics and 10 per cent lacustrine alluvials).
- (iii) **Objectives:** To demonstrate effective use of river water for gravity irrigation of 14,000 ha including strengthening of agricultural supporting services and development of appropriate rice varieties and crop calendars.

(c) Casier Sud (L) (flood protection/pump irrigation):

- (i) **Agricultural land type:** Mekong plain area characterized by annual flooding and drought.
- (ii) **Soil types:** Clayey floodplain soils and lighter levee soils (both potentially productive, assuming flooding can be controlled).
- (iii) **Objectives:** To increase agricultural production by flood control and irrigation.

(d) Se Bang Fai (L) (flood protection/pump irrigation):

- (i) **Agricultural land type:** Typical flood plains of Mekong tributaries.
- (ii) **Soil types:** Alluvial deposits with characteristics determined by elevation including low fertility older alluvium, fine texture floodplain soils, and fertile levee soils.
- (iii) **Objectives:** To increase agricultural productivity by minimizing or avoiding periodic crop failures (entails rice crop security 9 years of 10 instead of 2 of 3 as at present) and by dry season irrigation.

(e) LamPao (TH) (irrigation improvement)

- (i) **Agricultural land type:** Korat Plateau.
- (ii) **Soil types:** Leached sandy loams suitable, for both rice and upland crops (85 per cent) with some heavy clay in lower areas (15 percent).
- (iii) **Objectives:** To demonstrate how to accelerate utilization by the farmers of water made available by major irrigation works.

(f) Northeast Thailand (pump irrigation)

- (i) **Agricultural land type:** Riverine lands along the Mekong including river levees, old river channels, and poorly drained swamp areas.
- (ii) **Soil types:** Alluvial, usually fairly heavy silty clays (about 10 per cent suitable for upland crops).
- (iii) **Objectives:** To demonstrate solutions to technical and managerial problems of pump irrigation including use of higher cropping intensity and high-yielding rice varieties.

(g) Northeast Thailand (tank irrigation)

- (i) Agricultural land type:** Riverine floodplains and low terraces.
- (ii) Soil types:** Low humic gleys derived from alluvial materials with high content of fine sand and silt in upper horizon with underlying horizons of hard impermeable clay.
- (iii) Objectives:** To demonstrate solution of technical and managerial problems in tank irrigation including irrigation system rehabilitation.

(h) Nong Wai (TH): (irrigated agriculture)

- (i) Agricultural land type:** Typical rice growing area of Korat Plateau.
- (ii) Soil types:** Alluvial clay hydromorphics (43 per cent), low humic gley soils (26 per cent) on low and middle terraces.
- (iii) Objectives:** To demonstrate comprehensive and accelerated development of gravity irrigation with emphasis on provision of terminal facilities, on-farm improvements including levelling and consolidation, and development of farmer organizations.

(i) Go Cong (VN): (gravity irrigation)

- (i) Agricultural land type:** Eastern coastal region of Mekong delta featuring lack of-freshwater in dry season, periodic flooding, poor drainage, saline soils, and inadequate roadway access.
- (ii) Soil types:** Mostly heavy-textured, poorly-drained, with low permeability including saline soils in lower areas along streams.
- (iii) Objectives:** To furnish dry season irrigation water from local canal, improve drainage, prevent tidal flooding, and improve the road system.

(j) Tiep Nhut (VN): (salinity control)

- (i) Agricultural land type:** Typical of large areas in delta coastal region subject to seawater intrusion, soil salinity, and poor drainage.
- (ii) Soil types:** Heavy alluvial clays with some young river and marine sediments and locally disturbed soils along the streams and canals.
- (iii) Objectives:** To improve conditions to permit rice cultivation in 46,000 ha now subject to salinity intrusion and poor drainage including low-lift pump irrigation.

(k) Caisan (VN): (low-lift pump irrigation)

- (i) Agricultural land type:** Typical of upper delta, featured by seasonal flooding where floating rice is the dominant crop.
- (ii) Soil types:** Recent fine-textured delta alluvium with some marine influences, mostly acid with some acid sulfate areas, well suited to rice cultivation because of their low permeability.

- (iii) **Objectives:** To promote dry season cropping in the delta flood zone, based on irrigation from existing canals with low-lift pumps including provision of credit and technical assistance.

Conclusions: Long-Term Prospects for Agricultural Development

Long-term prospects for Agricultural Development

It has been calculated that each cubic meter of irrigation water results in an average production of 0.32 kg of paddy with present irrigation techniques, and that basin residents annually consume about 300 kg of paddy equivalent per capita (Pa Mong Phase II report); thus the irrigation water drawn from the Mekong can yield enough rice to feed about 65 million people (illustration 31). Assuming that these production levels could be attained by the time that the basin population would have increased by about 30 million people, then sufficient rice (or equivalent crops) will be available for export to feed about 50 million people. At the 1974 export price of US\$ 460/ton, the approximately 14 million tons of surplus rice for export will earn about US\$ 6,440 million per year.

ILLUSTRATION 31: ULTIMATE WATER RESOURCES DEVELOPMENT FOR AGRICULTURE					
	Cambodia	PDR Lao	Northeast Thailand	Mekong Delta	Totals
Irrigated area (1)	100	102	1,300	1,028	2,530
Water requirements (2)	1,690	1,7102	7,000	27,000	57,400
Increase in production (3)	0.54	0.54	8.01	9.18	18.27
Area reclaimed by flood control (1)	151	59	42	-	252
Increase in production (3)	0.30	0.12	0.84	-	1.26
Increase in people fed (4)	2.8	2.22	9.53	0.66	5.1

1. Thousand hectares
2. Million m³/yr
3. Million tons paddy equivalent/yr over rainfed conditions at realistically attainable yields; includes double cropping.
4. Millions of people, consuming 300 kg paddy equivalent/capita/yr.

Another important point in the overall agricultural plan relates to conservation in the watersheds, in that agricultural development of the magnitude envisioned will be sufficient to draw the extractive agriculturalists to the lowlands, thus reducing the current levels of exploitation in the uplands and allowing these areas to be used for their optimal long-term conservation purposes.

The primary purpose of the Mekong agricultural programme is to produce enough food to meet its own needs, based on the likely rate of population growth and considering that the traditional agricultural methods have already exploited and in some cases over-exploited the readily available lands. The objective of the model is to replace traditional agricultural and extractive agricultural methods with more modern methods to the degree needed to achieve the necessary levels of production on a sustained yield basis. A secondary objective is to use this same approach to achieve a further increase in production, thus realising a very sizeable surplus for export.

The various studies and evaluations described in this book show clearly that the climatic, soil, and water resources of the basin are sufficient to realize the desired objectives, through use of modern agricultural methods, especially irrigation, and that the key feature is water control. The natural water resources of the basin (rainfall, tributaries, and the Mekong mainstream) are more than ample on a total annual basis, but the distribution within the year is such that most areas have too little water in the dry season and substantial areas are too frequently flooded in the rainy season to be suitable for cropping. To remedy this, development of tributary water resource projects is the first step, and this is well underway. A total of 10 tributary dams have been built (nine of these in northeast Thailand), and many more tributary projects sites have been identified throughout the basin as probably feasible. Unfortunately, the two most important heavily populated areas of the lower Basin are lacking in sites for further tributary development, namely northeast Thailand (the Korat Plateau), where practically all tributary sites have already been utilized, and the Vietnamese delta, where no tributary sites are available.

Northeast Thailand, with full development of its tributaries, will need to import another 27,000 million m³ of water for irrigation annually in order meet anticipated minimum food needs over the next several decades, and the only feasible source is the Mekong mainstream.

With respect to the Vietnamese delta, while no food crisis is anticipated, this region has large-scale potentials for agricultural development capable of supporting a much higher density of population, as well as producing for other parts of the country and for export. It would seem to be of considerable importance to the economy of Viet-Nam to utilize this potential, but for that purpose a substantial dry season flow in the Mekong must be maintained. It is estimated that the shortage in the water balance for growing crops in the delta at full development would be about 50,000 million m³ per year.

The agricultural development programme for the basin, with respect to water management, must first pursue tributary projects to the full extent that these can meet the needs.

Under the Mekong Committee's programme in agricultural development, virtually every aspect of agriculture in the lower Mekong basin has been evaluated. This has included study of each of the major agro-ecological zones in the basin, with different cropping patterns stemming primarily from differences in the soil and hydrologic characteristics.

The analysis has covered all of the natural resources of the basin which can contribute to agriculture, and an assessment of their potential use in producing rice and other crops, employing both traditional and modern agricultural methods.

The analysis shows clearly that the lower Mekong basin, despite the fact that its soils are generally not very fertile, does have ample land areas with suitable soils sufficient to increase agricultural production. But to achieve this, adequate water control will be needed. Utilization of water from the Mekong would not only be a great blessing to the people concerned but, considered over a longer time-span, it could lead the way in transforming the lower Mekong basin into a sustaining yield surplus food production area, perhaps to become one of the food baskets of the world.

Chapter 11

The Lower Mekong Basin Today

This book has described agricultural development and natural resource management in the Lower Mekong Basin as of 1977-1980. So why is the book still relevant today? It would have been interesting to revisit the various pioneer sites to see how conditions have evolved, and to review the social, economic, cultural, and political changes that have affected the region since the book was first written. We would have had to review a vast literature, carry out at least a few months of field work, and compare our various projections with the actual conditions on the ground. But the relevant current data are not aggregated in a way that would readily enable the necessary comparisons to be made, and both of us have other responsibilities that have prevented us from writing what would, in essence, be a second book. We hope that some enterprising researchers will take on this challenge, reveal how accurate were our projections, and indicate how new technologies have rendered our hopes for development into realities on the ground.

But we also believe that the book is still highly relevant, beyond its historical interest. In reviewing our work from several decades ago, we were intrigued by how many of the basic ideas developed under the Mekong Committee's Agriculture Programme remain topical today. Initiatives such as involving farmers as part of the agricultural research team, diversifying crops on farms so that the various species complement each other, considering individual farms as part of a larger landscape that includes other land uses as well (today sometimes called "ecosystem management" or "landscape management"), the importance of wild species in contributing to rural livelihoods, and the role of protected areas in contributing to the welfare of rural people, all remain high on the agenda of those seeking to support sustainable rural development. Indeed, the agricultural strategy of the Mekong Project contains the essential elements of "sustainable development", which became the dominant development paradigm of the decades immediately following the completion of our work.

IUCN – The World Conservation Union has been contributing to these forms of development in the Mekong Basin for several decades. Southeast Asia became one of the focal areas for IUCN when Dr. Boonsong Lekagul was an IUCN Vice-President and convened an international meeting in Bangkok in 1965. Entitled "Conservation in Tropical South East Asia", the meeting brought some of the world's leading scientists and resource managers to the region. Their deliberations contributed to many of the ideas that have been further elaborated in this book and that have been applied to both working landscapes and areas designated for conservation. IUCN's recent efforts to design protected areas in Lao PDR, Cambodia, and Vietnam so that they provide benefits to the rural population have helped to change the relationship between sustainable agriculture and nature conservation, making them now part of the same human enterprise

Of course, much has changed since the late 1970s. The word “biodiversity”, now the subject of an international convention with 188 Parties, was not even coined until the mid-1980s. But many of the principles of the Convention on Biological Diversity are already evident in this book. The objective of providing equitable shares of benefits to the rural populations, the essential role that pollinators play in supporting agriculture, the importance of predators in controlling agricultural pests, the value of conserving upstream watersheds that provide the high-quality water that supports the irrigation systems that enable agriculture to be intensified, the concept of sustainable use, the role of incentives in promoting conservation, and even the recognition that a wide diversity of plants and animals enhance human well-being have all been discussed in some detail in this book. Such approaches did not originate with the Mekong Project, but rather are part of the evolution of thinking beyond single high-yielding varieties of crops. Many more scientists now recognize that a diversity of approaches to development is just as important as diversity in nature, and that people who have lived long on the land have knowledge that is as deserving of respect as modern scientific knowledge in solving the complex challenges of earning a living in difficult environments. We believe that our work represented in this book brought these multiple ideas together for the first time in Southeast Asia and applied them on a landscape scale that extended well beyond single farms. Donor countries such as the Netherlands, Australia, and the USA saw the value in these approaches, and the European Union provided its first major technical support to the agricultural programme of the Mekong Committee.

War disrupted many of the promising initiatives recorded in this book, but new innovations are building on the foundations laid by the Mekong Committee. Of course, many challenges still remain. While the new protected area systems in the Lower Mekong Basin look excellent on a map, the reality on the ground still leaves much to be desired. Poverty remains a stark reality for many farmers, and globalization is offering new hurdles to be jumped. Farm mechanization has become widespread, but as the price of oil inexorably increases, what is this going to do to the profitability of farms that have become dependent on machines? How is the balance of power and welfare between urban and rural people going to be resolved? What role will biotechnology play (a topic that was hardly imaginable in the 1970s)? Such issues go far beyond the specific agricultural development issues we have discussed in this book, but provide the background against which further developments will need to take place.

The Lower Mekong Basin remains a special place for us, a diverse landscape united by one of the world’s great rivers, the home of some of the world’s most fascinating civilizations, supporting forests where amazing new species of wildlife are still being discovered, and offering a richness of opportunities to the farmers who are constantly seeking new ways to earn a living from the soil. We hope that our early contribution to developing agriculture in ways that also conserve natural resources at a landscape scale will stimulate further research and development that will enable the rich cultures of the Basin to live in peace, harmony, and well-being.

References

- Backer, M. de, and K. Openshaw. 1972. **Timber Trends Study-Thailand**. FAO, Rome.
- Bardach, J.E. 1960. **Report on Fisheries in Cambodia**. UDOM, Phnom Penh.
- Brinkman, R. 1970. Ferrollysis, a Hydromorphic Soil Forming Process. **Geoderma** 3:199-206
- Brink, R. 1974. Surface-water Gley Soils in Bangladesh, ms.
- Brink, R. and P. Dieleman. 1977. Problem Hydromorphic Soils in northeast Thailand. **Proc. Conf. on Classif. and Mgmt. of Tropical Soils**, Kuala Lumpur, Malaysia.
- Bradfield, R. 1972. Maximizing Food Production through a Multiple Cropping System Based on **Rice**. **Rice, Science, and Man**. IRRI, Los Banos, Philippines.
- Carbonel, J P. 1972. Le quaternaire Cambodgien. **Mémoire Orstom** 60: 1-248.
- Ceruse, J. 1977a. **Land Use Map of the Lower Mekong Basin**. Mekong Secretariat, Bangkok.
- Ceruse, J. 1977b. **Pedo-morphological Map of the Lower Mekong Basin**. Mekong Secretariat, Bangkok.
- Champion, H. 1962. **Memorandum on Forest Management in Thailand**. FAO, Rome.
- Chapman, E.C. 1970. Shifting Cultivation and Economic Development in the Lowlands of northern Thailand. pp. 226-250, in **International Seminar on Shifting Cultivation and Economic Development in Northern Thailand**.
- Crump, D. and S. Sanoh. 1972. **The Dairy Industry in Thailand**. New Zealand Colombo Plan Notes.
- Davidson, A. 1975. **Fish and Fish Dishes of Laos**. Imprimerie Nationale, Vientiane.
- Exell, R. and K. Saricali. 1976. **The Availability of Solar Energy in Thailand**. AIT, Bangkok.
- FAO. 1959. **A Survey of the Investigations Required for Planning Development of Agriculture, Forests, and Fisheries in the Lower Mekong Basin**. FAO, Rome.

FAO. 1974. **Coordination of Plant Production Research in Thailand**. Working Papers 1-6, FAO, Bangkok.

FAO. 1975. **Country Tables of Production, Trade, and Consumption of Forest Products**. FAO, Rome.

FAO. 1976. **Forest Resources in Asia and Far East Region**. FAO, Rome.

Hafner, J.L. 1973. Man and Environment in Rural Thailand. **Journal of the Siam Society** 61(2): 129-138.

Hagen, T. 1960. **Nepal**. Oxford, New Delhi.

Hinton, P. 1970. Swidden Cultivation Among the Pwo Karen of Northern Thailand, pp. 1-24 in **International Seminar on Shifting Cultivation and Economic Development in Northern Thailand**.

Janzen, D. 1972. Tropical agroecosystems. **Science** 182:1212-1219.

Karlberg, S. 1976. Draft paper on Timber/Kenaf farming.

Kawai, T. 1974. Analysis of Rainfall in Northeast Thailand as a Basis for the Planning of Irrigated Agriculture. Working Paper, Mekong Secretariat, Bangkok.

Kreuze, J. 1975. Summary of Monthly and Yearly Hydro-Meteorological Data in the Thai Part of the Lower Mekong Basin. Mekong Committee, Bangkok.

Kunstadter, R. 1970. Subsistence Agricultural Economies of Lua and Karen Hill Farmers of Mae Sariang District. pp. 47-144. **International Seminar on Shifting Cultivation and Economic Development in Northern Thailand**.

MBFS. 1976. **Fisheries and Integrated Mekong River Basin Development**. School of Natural Resources, University of Michigan, USA.

Mekong Committee. Annual Statistical Bulletins (since 1966). Lower Mekong Hydrologic Yearbooks (since 1962).

Mekong Committee. 1970. Report on Indicative Basin Plan, A Proposed Framework for the Development of Water and Related Resources of the Lower Mekong Basin.

Mekong Secretariat. Agricultural Periodicals I, II, III.

Mekong Secretariat. 1977. **Agricultural Research Efficiency in Thailand: Vol. I Rice; Vol II Maize**. Mekong Secretariat, Bangkok.

Mekong Secretariat. 1977. **Pa Mong Optimization and Downstream Effects Study.** Mekong Secretariat, Bangkok.

Menasveta, D., S. Shindo, and S. Chullasorn. 1973. **Pelagic Fishery Resources of the South China Sea and Prospects for Their Development.** FAO, Rome.

McNeely, J.A. 1975. **Draft Report on Wildlife and National Parks in the Lower Mekong Basin.** Mekong Committee, Bangkok.

Molnar, P. and P. Tapponier. 1975. Cenozoic Tectonics of Asia: Effects of a Continental Collision. **Science** 189: 419-426.

Mouhot, H. 1864. **Travels in the Central Part of Indochina.** John Murray, London.

NDDT (Netherlands Development Team), 1974. **Recommendations Concerning Agricultural Development with Improved Water Control in the Mekong Delta,** Mekong Committee, Bangkok.

Netherlands Economic Institute. 1974. **Long-Term Projections of Domestic and Foreign Demand for Agricultural Products for the Countries of the Lower Mekong Basin.** Mekong Secretariat, Bangkok.

Owen, N.G. 1971. The Rice Industry of Mainland Southeast Asia, 1850-1914, **Journal of the Siam Society** 59(2): 75-143.

Pa Mong Phase II Report. 1972. Main Report. Mekong Committee, Bangkok.

Pavie, A. 1904. Mission Pavie Indochine 1879-1895. Poisson, Paris.

Powell, J.H. 1971. Crocodiles and Public Relations. **IUCN Publication 1 (N.S.) 32:** 149-151.

Shuyler, S. and S. Ratanaworabhan. 1970. **Rodent Damage to Rice in Thailand.** ASRCT, Bangkok.

Singh, G. 1972. **Draft Report on the Watershed Conditions in the Lower Mekong Basin.** Mekong Committee, Bangkok.

Spencer, J.E. 1966. **Shifting Cultivation in Southeast Asia.** University of California Press, Berkeley, CA.

Takaya, Y. 1974. **Salinity in northeast Thailand.** Mekong Committee, Bangkok,

Talbot, L.M. 1973. **Effects of Mekong Development on Wildlife, Parks, and Reserves.** SEADAG, Washington D.C.

Tan, V.N. 1968. The Raising of Deer: An Attractive Resource for Developing Countries in Southeast Asia. **IUCN Publication 1. (N.S.)** 10:132-134.

Thompson, L.M. 1975. **Weather Variability, Climatic Change and World Food Supplies.** American Association of Agricultural Science, New York.

Tomosugi, T. 1965. Historical Development of Irrigation and Drainage in the Chao Phya Delta. pp. 165-176 in **Water Resource Utilization in Southeast Asia.** OTCA, Tokyo.

Turbang, 1973. Unpublished report, FAO, Bangkok.

Uehara, G., M. Nishina, and G. Tsugi. 1974. **The Composition of the Mekong River Silt and its Possible Role as a Source of Plant Nutrients in Delta Soils.** University of Hawaii, Honolulu.

UNDP/FAO. 1969. **Report of the Joint UNDP/FAO Mission on the Review of the Agricultural Work of the Mekong Committee with Special Reference to the Agricultural Research, Demonstration and Training Programmes.** UNDP, New York.

van der Kevie, W. 1972. Land Capability Map, Mekong Committee, Bangkok.

Van Liere, W.J. 1977. Southeast Asia, potential world food basket. **Netherlands Journal of Agricultural Science** 25: 14-25

Wharton, C. 1968. Man, Fire, and Wild Cattle in Southeast Asia. **Tall Timbers Fire. Ecol. Conf.** 8:107-167.

White, G. et al., 1962. **Economic and Social Aspect of Lower Mekong Development.** Mekong Committee, Bangkok.

Whitemore, T.C. 1975. **Tropical Rainforests of the Far East.** Clarendon Press, Oxford.

Whyte, R. 1972. The Graminae, Wild and Cultivated, of Monsoonal and Equatorial Asia, I. Southeast Asia. **Asian Perspectives** 15:127-151.

Yantasast, A. et al., 1970. Breeding Dwarf Varieties of Rice for Tolerance to Deep Water. **Thai Journal of Agricultural Science** 3-19.

IUCN – The World Conservation Union

Founded in 1948, The World Conservation Union brings together States, government agencies and a diverse range of non-governmental organizations in a unique world partnership: over 1000 members in all, spread across some 150 countries.

As a Union, IUCN seeks to influence, encourage and assist societies throughout the world to conserve the integrity and diversity of nature and to ensure that any use of natural resources is equitable and ecologically sustainable. A central Secretariat coordinates the IUCN Programme and serves the Union membership, representing their views on the world stage and providing them with the strategies, services, scientific knowledge and technical support they need to achieve their goals. Through its six Commissions, IUCN draws together over 10,000 expert volunteers in project teams and action groups, focusing in particular on species and biodiversity conservation and the management of habitats and natural resources. The Union has helped many countries to prepare National Conservation Strategies, and demonstrates the application of its knowledge through the field projects it supervises. Operations are increasingly decentralized and are carried forward by an expanding network of regional and country offices, located principally in developing countries.

The World Conservation Union builds on the strengths of its members, networks and partners to enhance their capacity and to support global alliances to safeguard natural resources at local, regional and global levels.

Chief Scientist's Office

Rue Mauverney 28
CH-1196 Gland
Switzerland
Tel: +41 22 999 02 84
Fax: +41 22 999 00 25
E-mail: JAM@iucn.org
www.iucn.org

IUCN Publications Services Unit

219c Huntingdon Road
Cambridge CB3 0DL, UK
Tel: +44 1223 277894
Fax: +44 1223 277175
E-mail: books@iucn.org
www.iucn.org/bookstore